



Molecular characterization and population dynamics of lactic acid bacteria during the fermentation of sorghum

by

EVELYN MADOROBA

Submitted in partial fulfilment of the requirements for the degree of

Philosophiae Doctor

in the Faculty of Natural and Agricultural Sciences

University of Pretoria

Pretoria

March 2009

TABLE OF CONTENTS

DECLARATION.....	iii
ACKNOWLEDGEMENTS.....	iv
DEDICATION.....	v
SUMMARY.....	vi
LIST OF FIGURES.....	viii
LIST OF TABLES.....	ix
LIST OF ABBREVIATIONS.....	x
RESEARCH COMMUNICATIONS.....	xii
CHAPTER ONE.....	1
LITERATURE REVIEW	
CHAPTER TWO.....	55
POLYPHASIC TAXONOMIC CHARACTERIZATION OF LACTIC ACID BACTERIA ISOLATED FROM SPONTANEOUS SORGHUM FERMENTATIONS USED TO PRODUCE <i>TING</i> , A TRADITIONAL SOUTH AFRICAN FOOD	
CHAPTER THREE.....	70
DIVERSITY AND DYNAMICS OF BACTERIAL POPULATIONS DURING SPONATNEOUS FERMENTATIONS USED TO PRODUCE <i>TING</i> , A SOUTH AFRICAN FOOD	
CHAPTER FOUR.....	103
USE OF STARTER CULTURES OF LACTIC ACID BACTERIA IN THE PRODUCTION OF <i>TING</i> , A SOUTH AFRICAN FERMENTED FOOD	
CHAPTER FIVE.....	139
GENERAL DISCUSSION AND CONCLUSIONS	
APPENDICES.....	143

DECLARATION

I declare that the thesis, which I hereby submit for the degree, Philosophiae Doctor (Microbiology) at the University of Pretoria, is my own work and has not previously been submitted by me for a degree at another University.

Signed:

Date:

ACKNOWLEDGEMENTS

I am greatly indebted to Prof. T. E. Cloete for awarding me an opportunity to work in his laboratory just when my academic career was at the crossroads. Prof. Cloete supplied crucial advice and inspiration throughout my studies.

I am also grateful to Prof. J. Theron and Dr. E. T. Steenkamp, who have been my co-promoters, for their expert advice and academic passion in my PhD studies. Both co-promoters supplied a plethora of significant information that improved this thesis.

Dr. Geert Huys, Dr. Ilse Scheircklinck and Ann Van Schorr are thanked for assistance with polymerase chain reaction denaturing gradient gel electrophoresis (PCR-DGGE), which gave a refreshing dimension to this thesis.

I sincerely wish to express special thanks to Prof. Gert Marais for assistance with identification of fungi. I also wish to express my heartiest appreciation to Dr. H. L. De Kock for expert advice regarding sensory evaluation of *ting*.

This project was made possible by financial support from the National Research Foundation, South Africa. I am also indebted to the Third World Organisation for Women in Science for awarding me a PhD fellowship.

My sincere gratitude is extended to my family and friends for their unwavering support and encouragement during times of distress. I shall never forget the patience and prayers of Chawapiwanashe Chipso and Masimba Michael Madoroba. My husband is thanked for his support during my absence from home.

Above all, I would like to thank the invisible force behind the success of all my experiments, and this is none other than the Almighty God.

DEDICATION

This thesis is dedicated to my parents, the late Mr. Henry Mahembe and Mrs. Grace Mahembe for emphasizing the significance of a sound education.

SUMMARY

Molecular characterization and population dynamics of lactic acid bacteria during the fermentation of sorghum

by

Evelyn Madoroba

Promoter: Prof. T.E. Cloete

Co-promoters: Prof. J. Theron and Dr E.T. Steenkamp

Department of Microbiology and Plant Pathology

University of Pretoria

for the degree Ph.D

Ting is a cooked fermented sorghum food that is popular amongst southern Africans for its sour taste and unique flavour. However, major challenges are associated with large-scale production and marketing of this spontaneously fermented food due to inconsistent microbiological and sensory quality. The use of starter cultures may circumvent these limitations. Prior to engaging starter cultures, detailed knowledge of the microbial diversity and dynamics during fermentation is important. Therefore, the aim of this study was to investigate microbial diversity and dynamics during sorghum fermentations, and to clarify the role of starter cultures regarding the microbiological safety and consumer acceptance of sensory characteristics of fermented *ting*.

A culture-independent approach, based on the use of PCR-denaturing gradient gel electrophoresis (DGGE), revealed that *Lactococcus lactis*, *Lactobacillus curvatus*, *Weissella cibaria* and some *Enterobacteriaceae* were predominant at the end of spontaneous sorghum fermentations. Culture-dependent methods indicated that *Lb. fermentum*, *Lb. plantarum*, *Lb. rhamnosus*, *E. faecalis*, *E. mundtii*, *W. cibaria* and *L. lactis* were predominant at the end of fermentation. These results not only indicated the predominant bacteria during sorghum fermentation, but also indicated that a combined approach is required to reveal microbial diversity and dynamics during spontaneous sorghum fermentations.

Based on the above results, *L. lactis*, *Lb. fermentum*, *Lb. plantarum* and *Lb. rhamnosus* were evaluated as starter cultures for production of *ting*. All the starter cultures were able to ferment sorghum, but the lowest pH and highest lactic acid was produced in naturally fermented sorghum inoculated with *L. lactis*. This fermentation showed an increase in the number of lactic acid bacteria and yeasts, whilst pathogen counts decreased. *Ting* from this fermented gruel, in contrast to naturally fermented sorghum, had sensory properties preferred by panelists. The results indicated that the use of *L. lactis* in starter cultures may result in *ting* with consistent and acceptable attributes.

LIST OF FIGURES

Fig. 1.1	Generalized scheme for the formation of important metabolic products from pyruvate by lactic acid bacteria.....	8
Fig. 1.2	Neighbor-joining phylogenetic tree, based on 16S rRNA gene sequence analysis, showing the relationships among type species of the representative genera from the families of <i>Lactobacillales</i>	17
Fig. 1.3	Main steps involved in DNA-based methods for characterization of LAB.....	23
Fig. 1.4	Main steps involved in DNA-based methods for studying microbial population dynamics of LAB.....	28
Fig. 2.1	Maximum likelihood (ML) phylogeny for the <i>Lb. plantarum</i> , <i>Lb. casei</i> and <i>Lb. reuteri</i> Groups based on 16S rRNA and <i>pheS</i> gene sequences.....	64
Fig. 3.1	Change in pH, as well as PCR-DGGE banding patterns representing the 16S rRNA gene fragments of the bacteria involved in the spontaneous fermentation of sorghum at 25°C.....	82
Fig. 3.2	Maximum likelihood (ML) phylogeny of species in the genera <i>Lactobacillus</i> and <i>Weissella</i> , based on 16S rRNA and <i>pheS</i> gene sequences.....	84
Fig. 3.3	Maximum likelihood (ML) phylogeny of species in the genera <i>Lactococcus</i> and <i>Enterococcus</i> based on 16S rRNA gene sequences.....	86
Fig. 4.1	Changes in pH and titratable acidity in sorghum powder slurries during natural fermentation for 54 h and inoculated fermentations with different LAB starter cultures for 24 h.....	123

LIST OF TABLES

Table 1.1	Examples of sorghum-based fermented foods, natural microflora and the methods used for characterization.....	12
Table 1.2	Families and genera in the order <i>Lactobacillales</i>	20
Table 2.1	Differential characteristics of nine LAB isolates from fermented sorghum, based on API 50 CHL analysis.....	62
Table 3.1	Microbial counts (cfu/g) at the start and end of spontaneous sorghum fermentations obtained on different selective media.....	80
Table 3.2	Likely species identities of bacteria isolated at the beginning and end of sorghum fermentations.....	81
Table 3.3	Comparison of the bacterial diversity in <i>ting</i> using culture-independent and culture-dependent approaches.....	88
Table 4.1	Experimental protocol for the six sorghum fermentations.....	109
Table 4.2	Culture conditions used for microbiological analysis of sorghum samples.....	111
Table 4.3	Microbial counts (log cfu/g) of sorghum powder, fermented sorghum slurries and corresponding cooked fermented porridge (<i>ting</i>).....	118
Table 4.4	Occurrence of bacterial food-borne pathogens in sorghum powder, fermented sorghum slurries and cooked <i>ting</i>	120
Table 4.5	Antimicrobial interactions among lactic acid bacteria used as starter cultures for sorghum fermentation.....	121
Table 4.6	Acceptability of the sensory characteristics of <i>ting</i> produced using different starter cultures based on average hedonic ratings.....	125
Table 4.7	Pearson's product-moment correlation coefficients (<i>r</i>) for pH, titratable acidity and acceptability of sensory attributes associated with <i>ting</i> produced using different starter cultures.....	126

LIST OF ABBREVIATIONS

%	percentage
°C	degrees Celsius
bp	base pair
cfu	colony forming units
ClustalX	cluster analysis version X
cm	centimetre
DMSO	dimethyl sulphoxide
DNA	deoxyribonucleic acid
dNTP	deoxyribonucleotide triphosphate
EDTA	ethylenediaminetetraacetic acid
ERIC	enterobacterial repetitive intergenic consensus
Fig.	figure
g	gram
h	hour
LAB	lactic acid bacteria
<i>Lb.</i>	<i>Lactobacillus</i>
<i>L.</i>	<i>Lactococcus</i>
<i>Le.</i>	<i>Leuconostoc</i>
LMG	Laboratorium voor Microbiologie Gent
M	molar
mg	milligram
min	minute
ml	millilitre
mm	millimetre
mM	millimolar
ng	nanogram
nm	nanometre
PCR	polymerase chain reaction
RAPD	randomly amplified polymorphic DNA
rep	repetitive extragenic palindromic
RFLP	restriction fragment length polymorphism
RNA	ribonucleic acid
rpm	revolutions per minute
rRNA	ribosomal ribonucleic acid



TE	Tris-EDTA
TEMED	N, N, N', N'-tetramethylenediamine
U	unit
UPGMA	unweighted pair group method of arithmetic averages
v/v	volume per volume
w/v	weight per volume
w/w	weight per weight
μg	microgram
μl	microlitre

RESEARCH COMMUNICATIONS

Published papers:

Evelyn Madoroba, Emma T. Steenkamp, Jacques Theron, Geert Huys, Ilse Scheirlinck and T. Eugene Cloete (2008). Polyphasic taxonomic characterization of lactic acid bacteria used to produce *ting*, a South African fermented food. *African Journal of Biotechnology*, **8** (3), 458-463.

Manuscripts in preparation:

Evelyn Madoroba, Emma T. Steenkamp, Jacques Theron, Geert Huys, Ilse Sheirlinck and T. Eugene Cloete. Diversity and dynamics of bacterial populations during spontaneous sorghum fermentations used to produce *ting*, a South African food.

Evelyn Madoroba, Jacques Theron, Emma T. Steenkamp, Henriëtte L. De Kock and T. Eugene Cloete. Use of starter cultures of lactic acid bacteria for the production of *ting*, a South African fermented food.

Conference contributions:

Evelyn Madoroba, Jacques Theron, Henriëtte De Kock, Emma Steenkamp and Eugene Cloete. Influence of starter cultures of lactic acid bacteria to the microbiological safety and sensory acceptability of *ting*, a fermented food. Food Micro Symposium, Aberdeen, Scotland, 1-4 September 2008.

Evelyn Madoroba, Emma T. Steenkamp, Jacques Theron, Geert Huys, Ilse Scheirlinck and T. Eugene Cloete. Polyphasic taxonomic characterization of lactic acid bacteria used to produce *ting*, a South African fermented food. Science with Africa Conference, Addis Ababa, Ethiopia, 1-3 March 2008.



CHAPTER ONE

LITERATURE REVIEW

TABLE OF CONTENTS

1.1 INTRODUCTION	3
1.2 SORGHUM-BASED FERMENTATION	3
1.2.1 Anti-nutritional compounds in sorghum	4
1.2.2 Nutritional and health advantages of lactic acid fermentation.....	5
1.2.3 Sensory advantages of lactic acid fermentation	6
1.2.4 Safety benefits of lactic acid fermentation.....	9
1.3 SORGHUM-BASED FERMENTED FOODS	11
1.4 STARTER CULTURES	14
1.5 TAXONOMY OF LACTIC ACID BACTERIA (LAB)	14
1.5.1 General characteristics of LAB	16
1.5.2 Horizontal gene transfer (HGT) and LAB	16
1.5.3 The order <i>Lactobacillales</i>	19
1.6 IDENTIFICATION OF LAB ASSOCIATED WITH FERMENTED FOOD .21	
1.6.1 Phenotypic methods	21
1.6.2 DNA-based methods	22
1.7 DNA-BASED METHODS FOR STUDYING LAB COMMUNITIES	26
1.8 CONCLUSIONS AND AIMS OF THIS INVESTIGATION	31
1.9 REFERENCES	32

1.1 INTRODUCTION

It is generally accepted that sorghum (*Sorghum bicolor* (L.) Moench) originated in Ethiopia, between 5 000 and 7 000 years ago (ICRISAT, 2005). From there, sorghum was distributed along the trade and shipping routes around Africa and the Middle East to India, China and the Far East (Doggett, 1970; Kimber, 2000). Sorghum was introduced in America from West Africa through the slave trade. Currently, sorghum is mostly cultivated in the arid and the semi-arid regions of the world and is the fifth most important cereal after wheat, maize, rice and barley in terms of production (Food and Agricultural Organization, 2003; 2006).

In sub-Saharan Africa, sorghum is especially significant for food security and comes second after maize in being a staple food (Dendy, 1995). Sorghum is drought resistant by nature (Jordan and Sullivan, 1982; National Research Council, 1996; Dicko *et al.*, 2006), which makes it particularly relevant to sub-Saharan Africa, as an estimated half a million families are susceptible to hunger due to drought (FAO, 2003). In addition, the ability to tolerate periods of water-logging and high temperatures (Doggett, 1988) makes sorghum an attractive crop for providing dietary energy and protein for people living in these regions.

The use of sorghum as a main constituent of the diet is usually associated with under-nourishment. This is mainly due to it lacking some essential amino acids (Neucere and Sumrell, 1979; Kazanas and Fields, 1981) and the presence of anti-nutritional factors (Hamacker *et al.*, 1986; Ibrahim *et al.*, 2005). However, fermentation, facilitated mainly by lactic acid bacteria (LAB), greatly improves the nutritive value and sensory properties of sorghum (Au and Fields, 1981; Chavan *et al.*, 1988; Hassan and El Tinay, 1995). In this Chapter, the fermentation process and its associated advantages with specific reference to sorghum is reviewed. Aspects such as the taxonomy of LAB and DNA-based methods to study them are also emphasized.

1.2 SORGHUM-BASED FERMENTATION

The term fermentation is generally used to describe the desirable biochemical changes brought about by microorganisms and/or enzymes on primary food products (Nout and Motarjemi, 1997; Blandino *et al.*, 2003). Enzymatic fermentations involve chemical reactions whereby the enzyme acts as a catalyst, as occurs when starch is converted to dextrin by α -amylase. When fermentation involves microorganisms, these are either naturally present on

the substrate or they may be deliberately added in the form of starter cultures (Nout and Motarjemi, 1997). In microbiological terms, fermentation is a type of energy-yielding microbial metabolism whereby organic substrates are incompletely oxidized and organic carbohydrates act as electron acceptors (Adams, 1990).

The four main fermentation types are alcoholic, lactic acid, acetic acid and alkali based. Yeasts usually predominate in alcoholic fermentations, resulting in the production of alcohol, whereas bacteria such as *Acetobacter* species are responsible for fermentations where alcohol is converted to acetic acid in the presence of oxygen (Blandino *et al.*, 2003). The production of foods such as Japanese *natto* from cooked soybeans, *dawadawa* from African locust beans and *ogiri* from melon seeds often involve some form of alkaline fermentation, with *Bacillus subtilis* being the dominant species (Wang and Fung, 1996). LAB mainly carry out lactic acid fermentation.

LAB are essential for fermentation of most plant-based products where they usually co-exist with yeasts. In this regard, LAB play various roles that include the production of safe products through biopreservation as a result of bacterial antagonism (Oyewole, 1997; Soomro *et al.*, 2002), enhancement of sensorial and nutritive value of the foods, and saving of energy through reduced cooking time (Simango, 1997). A plethora of LAB predominate in plant-based foods such as *pozol*, a Mexican maize dough (Escalante *et al.*, 2001), Colombian *chicha* (Steinkraus, 1996), and *idli*, a fermented Asian food made from rice blended with black gram (Soni and Sandhu, 1991). LAB also play a central role during the production process of most fermented sorghum-based foods such as Ethiopian *injera* (Vogel *et al.*, 1993), Sudanese *kisra* (Mohammed *et al.*, 1991), Ugandan *bushera* (Muyanja *et al.*, 2003) and Tanzanian *togwa* (Mugula *et al.*, 2003).

1.2.1 Anti-nutritional compounds in sorghum

Sorghum contains significant levels of anti-nutritional compounds that include protease inhibitors, phytic acid (myoinositol hexakis) and oligosaccharides (Serna-Saldivar and Rooney, 1995). The protease inhibitors include non-specific tannins and Kunitz trypsin inhibitor. Although tannins have been shown to reduce serum lipids, increase blood clotting and reduce blood pressure (Chung *et al.*, 1998), their anti-nutritive attributes in foods far outweigh their medical value. For example, impaired utilization of minerals and vitamins and

inhibition of digestive enzymes are associated with these compounds (Chung *et al.*, 1998). Together, tannins and phytic acid inhibit digestive enzymes and reduce protein availability via different modes of action. Tannins typically form complexes with proteins (Hagerman, 1989), whilst phytic acid, a highly charged molecule (Serna-Saldivar and Rooney, 1995), chelates cations such as potassium, calcium, magnesium, iron and zinc, resulting in the formation of phytate (Wodzinski and Ullah, 1996; Dvorakova, 1998; Stodolak *et al.*, 2007). Oligosaccharides such as raffinose, stachyose and verbascose are responsible for causing flatulence (FAO, 1998). This is due to their indigestion by enzymes of the human gastrointestinal tract in the small intestine. Consequently, these oligosaccharides are passed to the large intestine where fermentation by resident microorganisms takes place, resulting in the production of gas (FAO, 1998).

1.2.2 Nutritional and health advantages of lactic acid fermentation

Lactic acid fermentation of sorghum foods by LAB prior to cooking and/or consumption significantly alters their biochemical properties, rendering a food product with enhanced nutritional value and flavour (Towo *et al.*, 2006). For instance, lactic acid fermentation improves the *in vitro* digestibility of starch (Hassan and El Tinay, 1995) and sorghum protein (Kazanas and Fields, 1981; Mertz *et al.*, 1984; Taylor and Taylor, 2002; Ibrahim *et al.*, 2005). The increase in *in vitro* protein digestibility may be due to a rapid decrease in pH that affects the structure of insoluble sorghum proteins such as prolamines and glutelins in such a way that they become more accessible to pepsin digestion (Taylor and Taylor, 2002).

Lactic acid fermentation has been shown to reduce the level of anti-nutritive oligosaccharides, phytate and tannins. The amount of poly- and oligosaccharides in sorghum, e.g., raffinose, stachyose and verbascose, are significantly reduced due to the action of α -galactosidases produced by some LAB that include *Leuconostoc mesenteroides* subsp. *mesenteroides*, *Weissella paramesenteroides*, *Lactobacillus fermentum*, *Lb. brevis* and *Lb. buchneri* (Mital *et al.*, 1973; Milliere *et al.*, 1989). The α -galactosidases disrupt the α -D-galactosidic bonds of oligosaccharides, resulting in lowered abdominal distention and flatulence (Nout and Motarjemi, 1997). Tannin levels are also reduced by lactic acid fermentation (Lorri, 1993; Osman, 2004). The reduction may be due to metabolic processes of the microorganisms. These processes include oxidation, reduction or dissociation of the tannins as a defense mechanism against toxicity by endogenous sorghum microflora (Bvochora *et al.*, 2005). Reduced levels of phytate, as a result of lactic acid fermentation, may increase B vitamins

(Kazanas and Fields, 1981; Nout and Motarjemi, 1997) and improve the availability of minerals (Marfo *et al.*, 1990).

Some LAB are thought to have probiotic properties (Fuller, 1989). These include prevention of kidney stones (Campieri *et al.*, 2001), treatment of heart disease (Schaafsma *et al.*, 1998), control of cholesterol (Gilliland *et al.*, 1985), stimulation of anti-carcinogenic action (Goldin, 1990), prevention of antibiotic-induced diarrhea (Fooks *et al.*, 1999), prevention of lactose intolerance (Gilliland and Kim, 1984), prevention of food allergies (Sütas *et al.*, 1996), blockage of the formation of biogenic amines (Joosten *et al.*, 1996) and prevention of vaginosis in humans (Kontiokari *et al.*, 2001; Reid, 2001). Lactic acid fermentation is, however, linked to certain diseases, especially in immunocompromised hosts. For instance, the D (-) lactic acid produced during fermentation may potentially be toxic to malnourished and sick children due to acidosis (Nout and Motarjemi, 1997). LAB have also been associated with clinical infections such as urethritis, endocarditis, endometritis, pneumonia, arthritis, bacteremia and meningitis (Sussman *et al.*, 1986; Struve *et al.*, 1988; Isenberg *et al.*, 1988; Ruoff *et al.*, 1988; Aguirre and Collins, 1993; Salminen *et al.*, 2002; Flaherty *et al.*, 2003). In addition, other members of genera such as *Leuconostoc*, *Enterococcus*, *Weissella* and *Lactobacillus* may cause dental caries (Monchois *et al.*, 1999; Sbordone and Bortolaia, 2003). For these reasons, thorough characterization of LAB for use in food fermentations is of paramount importance.

1.2.3 Sensory advantages of lactic acid fermentation

During the production of many fermented foods, the development of appetizing flavour characteristics is crucial (Caplice and Fitzgerald, 1999). Particularly, during cereal fermentations, several volatile compounds are formed, which contribute to a complex blend of appetizing flavours (Chavan and Kadam, 1989). The specific flavour occurs due to secondary metabolites produced during fermentation of macromolecules such as sugars (Longo and Sanromán, 2006). The specific flavour compounds produced include diacetyl, which imparts a buttery aroma (Rankine *et al.*, 1969; Davis *et al.*, 1985), acetaldehyde (yoghurt flavour) (Gobbetti and Corsetti, 1997; Abd El-Salam and Alichanidis, 2004) and the amino acid alanine, a natural sweetener (Hols *et al.*, 1999). Generally, these flavour compounds are products of pyruvate, which acts as an electron or hydrogen acceptor during metabolism by some LAB and yeasts, thereby perpetuating fermentation through recycling of NAD⁺ (Axelsson, 2004; Fig. 1.1). Although both yeasts and bacteria may contribute to the

formation of diacetyl, the role of yeasts during lactic acid fermentation is negligible (Martineau and Henick-Kling, 1995).

Depending on the bacterial strain under investigation and the growth conditions, pyruvate may act as substrate for the following pathways: dehydrogenase pathway (aerobic metabolism), pyruvate oxidase pathway, pyruvate-formate lyase system (anaerobic) or the diacetyl/acetoin pathway (Caplice and Fitzgerald, 1999; Axelsson, 2004). When pyruvate is present in excess (compared to NAD^+ regeneration), as occurs in the presence of another carbon source such as citrate, or if an alternative electron acceptor such as oxygen is present, the diacetyl/acetoin pathway is followed (Hugenholtz, 1993; Fig. 1.1). Yeasts also convert pyruvate to diacetyl via hydroxyethyl-thiamine (acetylaldehyde-thiamine pyrophosphate complex/active acetaldehyde) (Romano and Suzzi, 1996). The subsequent reactions are divided into three pathways, depending on the substrate, which interacts with the active acetaldehyde. Ultimately, pleasant sensory compounds, which include acetoin and diacetyl, are formed. Diacetyl formation usually occurs at low pH and decreased sugar concentration (Axelsson, 2004). Despite the pleasant sensory attributes imparted by these flavour compounds, high concentrations of diacetyl (above 5-7 mg/L) are considered unpleasant (Rankine *et al.*, 1969; Davis *et al.*, 1985).

The pyruvate-formate lyase system is active under anaerobic conditions or when some substrate is limiting (Axelsson, 2004). This results in mixed acid fermentation wherein pyruvate and Coenzyme A (CoA) react to form formate and acetyl CoA with pyruvate-formate lyase acting as the catalyst (Fig. 1.1). The utilization of acetyl CoA results in the formation of ethanol (if it is used as an electron acceptor) or acetate (if it acts as a precursor for substrate-level phosphorylation) (Fig. 1.1), or a mixture of both. *Lb. casei* and some *Lactococcus lactis* strains mainly use this pathway (Thomas *et al.*, 1979). The pyruvate dehydrogenase pathway, which is mainly active in lactococci, involves pyruvate dehydrogenase that catalyses the formation of acetyl CoA under aerobic conditions in a manner similar to the pyruvate-formate lyase system, thus resulting in the formation of acetate (Fig. 1.1). In the pyruvate oxidase pathway, pyruvate oxidase catalyses the conversion of pyruvate to acetyl-phosphate (acetyl-P) and carbon dioxide with the simultaneous formation of hydrogen peroxide. Ultimately, acetate is formed from acetyl-P (Fig. 1.1). *Lb. plantarum* was reported to produce high concentrations of acetic acid, using pyruvate oxidase under aerobic conditions (Sedewitz *et al.*, 1984).




Figure 1.1 Generalized scheme for the formation of important metabolic products from pyruvate by lactic acid bacteria. Brackets indicate the four different pathways that are involved in pyruvate metabolism. Dotted arrows illustrate an external source of pyruvate. Black arrows show the direction of the reaction. Curved arrows indicate redox reactions involving NAD/NADH and energy-yielding reactions. (Adopted and modified from: Caplice and Fitzgerald, 1999; Axelsson, 2004).

Despite the sensory advantages that LAB impart to fermented foods, these bacteria may, however, also produce off-flavours in beer and wine (Caplice and Fitzgerald 1999), and in fish (Lyhs *et al.*, 2002). For instance, some strains of *Lb. plantarum* may cause off-flavours through the formation of aldehydes (Kandler and Weiss, 1986; Bottazi, 1988; Henick-Kling, 1995). Furthermore, the production of diacetyl by *Pediococcus damnosus* results in cloudy beer with an acid taste (Satokari *et al.*, 2000). Some species of LAB (e.g., *Lb. kunkei*) and acetic acid bacteria (e.g., *A. aceti*) may also suppress yeast growth during grape juice fermentation, resulting in sluggish or stuck fermentations (Drysdale and Fleet, 1989; Huang *et al.*, 1996). This highlights the importance of accurate characterization of the LAB for use during specific fermentations.

1.2.4 Safety benefits of lactic acid fermentation

LAB render fermented foods safe for consumption by controlling the growth of pathogenic and spoilage microorganisms (Caplice and Fitzgerald, 1999). In order to create an unfavourable environment for these pathogenic and spoilage microorganisms, LAB produce a range of compounds that act as natural antimicrobials in different ways. These compounds include, but are not limited to, bacteriocins, reutericyclin, diacetyl, hydrogen peroxide, carbon dioxide and organic acids such as lactic and acetic acid (Silva *et al.*, 1987; De Vuyst and Vandamme, 1994; Lavermicocca *et al.*, 2000; Messens and De Vuyst, 2002; Leroy and De Vuyst, 2004). Some of these antimicrobial compounds were shown to be bactericidal or bacteriostatic for a diverse range of microorganisms belonging to the following genera: *Streptococcus*, *Mycobacteria*, *Lactococcus*, *Listeria*, *Clostridium*, *Bacillus*, *Neisseria*, *Campylobacter*, and *Haemophilus* (van Kraaij *et al.*, 1999; Mota-Meira *et al.*, 2000; Soomro *et al.*, 2002; Savadago *et al.*, 2006).

Some of the antimicrobial agents, e.g., bacteriocins, organic acids and carbon dioxide, produced by LAB disrupt the cytoplasmic membranes of target organisms by virtue of different properties (Riley and Wertz, 2002). For example, lactococcin A depolarizes or makes pores in the cytoplasmic membrane, thus disrupting the integrity of this structure (Van Belkum *et al.*, 1991). Reutericyclin is a highly hydrophobic and charged molecule that is produced during anaerobic growth of *Lb. reuteri* (Gänzle and Vogel, 2003). This compound is bacteriostatic or bactericidal against a broad range of food-associated spoilage and pathogenic microbes, including *Staphylococcus aureus* and *Listeria innocua*, due to its

activity as a proton ionophore, resulting in translocation of protons across the cytoplasmic membrane and consequently, dissipating the transmembrane pH gradient (Gänzle, 2004).

Organic acids such as lactic and acetic acids disrupt the cytoplasmic membranes of target microorganisms via both the dissociated and undissociated acid (Cherrington *et al.*, 1991). When these organic acids are in an undissociated form, they become lipid soluble and enter the cytoplasm directly where they dissociate, releasing anions (De Vuyst and Vandamme, 1994). The growth of affected bacteria is thus inhibited, as they divert most of their energy to curbing this acidification and its effects (Adams and Nicolaidis, 1997).

Carbon dioxide contributes to food safety by creating anaerobic conditions, which inhibit the growth of oxygen-dependent microorganisms such as moulds and Gram-negative bacteria (De Vuyst and Vandamme, 1994; Adams and Nicolaidis, 1997). Carbon dioxide also curbs the growth of pathogenic and spoilage bacteria by disrupting enzyme-dependent reactions due to a decrease in pH (King and Nagel, 1975).

Hydrogen peroxide, whose mode of action is not clear (since LAB do not possess a true catalase system), is believed to activate the lactoperoxidase system, resulting in the formation of other antimicrobials (De Vuyst and Vandamme, 1994). Its bactericidal effect is due to oxidation of the bacterial cell, membrane lipids and sulfhydryl groups of cell proteins (Lindgren and Dobrogosz, 1990). Hydrogen peroxide may accumulate to sufficient concentrations capable of inhibiting some pathogenic and spoilage bacteria (Condon, 1987). However, peroxidases, flavoproteins and pseudocatalase are believed to decompose hydrogen peroxide, prohibiting its accumulation to effective amounts *in vivo* (Fontaine *et al.*, 1996). Moreover, diacetyl contributes to food safety by disrupting the arginine utilization system of Gram-negative bacteria (Jay, 1982; Caplice and Fitzgerald, 1999). However, the concentrations that are required for antimicrobial action are not acceptable in foods (De Vuyst and Vandamme, 1994).

In spite of the safety contribution of lactic acid fermentation, the inhibition of pathogenic and spoilage organisms by LAB is complex because the factors involved are difficult to quantify. These include the organism in question, temperature, level of hygiene, amount of undissociated acid and the buffering capacity of the food (Adams and Nicolaidis, 1997; Nout and Motarjemi, 1997). For instance, when both the pathogen (*Escherichia coli*) and the LAB

(*Lactococcus lactis*) were added to a model weaning food simultaneously, the pathogen still grew for 5 h despite the fact that the LAB outnumbered the pathogen by more than 5 log cycles (Yusof *et al.*, 1993). Consequently, the antimicrobial effect of fermentation must be an adjunct to good hygiene but not a substitute for it (Adams and Nicolaidis, 1997; Nout and Motarjemi, 1997).

1.3 SORGHUM-BASED FERMENTED FOODS

Throughout the world, a plethora of fermented foods are metabolized by a variety of microorganisms to yield products with unique and appealing characteristics (Caplice and Fitzgerald, 1999; Taylor, 2003). In Africa, sorghum-based fermented foods and beverages include *injera* (Chavan and Kadam, 1989), *kisra* (Mohammed *et al.*, 1991), *ogi* (Akingbala *et al.*, 1981), *mahewu* (Bvochora *et al.*, 1999), *uji* (Mbugua *et al.*, 1984), *muramba* (Mukuru, 1992), *bushera* (Muyanja, 2003), *togwa* (Lorri and Svanberg, 1995) and *ting* (Boling and Eisener, 1982). LAB are mainly associated with most of these fermented foods (Table 1.1).

Injera is a circular, spongy-textured Ethiopian sour bread (Stewart and Getachew, 1962; Chavan and Kadam, 1989) made from sorghum, tef, corn, finger millet or barley. Over 8% of the total sorghum production in Ethiopia is used for *injera* production (Gebrekidan and Gebrettiwat, 1982). The production process for *injera* involves mixing of the mechanically dehulled grains with water, followed by addition of the starter (*ersho*) from a previous batch and fermentation for 2-3 days. A small part of the fermented mixture is gelatinized in boiling water, added back to the fermented dough and allowed to stand for 2-3 h. The resulting dough is thinned down after fermentation and poured into a thinly oiled pan, covered with a lid for 2-3 min and cooked (Parker *et al.*, 1989). This food is valued for its sour flavour and high nutritional value, being rich in calcium and iron (Zegeye, 1997). Yeasts and some fungi are mainly involved in the fermentation of *injera* (Blandino *et al.*, 2003; Table 1.1).

Fermented sorghum is used to make *kisra* that is baked into thin sheets to make pancakes, which are consumed throughout the Arabian Gulf, Sudan and Iraq (Eggum *et al.*, 1983). *Kisra* pancakes are similar to *injera*. The fermented dough (*ajin*) is prepared by using a portion of a previous batch (back-slopping). Fermentation was found to increase the crude protein, thiamine and niacin content of *kisra* (El Tinay *et al.*, 1979; Axtell *et al.*, 1981; Eggum *et al.*, 1983). A range of LAB, yeasts and fungi are involved in the production of *kisra* (Table 1.1).

Table 1.1 Examples of sorghum-based fermented foods, natural microflora and the methods used for characterization

Product	Country	Microorganisms isolated	Method of characterizing organisms	Reference
<i>Togwa</i>	Tanzania	<i>Lb. plantarum</i> , <i>Lb. brevis</i> , <i>Lb. fermentum</i> , <i>Lb. cellobiosus</i> , <i>P. pentosaceus</i> , <i>Weissella confusa</i> , <i>Issatchenkia orientalis</i> , <i>Saccharomyces cerevisiae</i> , <i>Candida pelliculosa</i> , <i>Candida tropicalis</i>	API 50 CHL	Mugula <i>et al.</i> , 2003
<i>Bushera</i>	Uganda	<i>Lb. plantarum</i> , <i>Lb. paracasei</i> subsp. <i>paracasei</i> , <i>Lb. fermentum</i> , <i>Lb. brevis</i> , <i>Lb. delbrueckii</i> subsp. <i>delbrueckii</i> , <i>Streptococcus thermophilus</i> , <i>Lactococcus lactis</i> subsp. <i>lactis</i> , <i>Leuconostoc mesenteroides</i> subsp. <i>mesenteroides</i> , <i>Leuconostoc mesenteroides</i> subsp. <i>dextranicum</i> , <i>Weissella confusa</i>	API 50 CHL	Muyanja <i>et al.</i> , 2003
Sorghum-based fermented weaning cereal	South Africa	<i>Lb. plantarum</i> , <i>Leuconostoc mesenteroides</i> , <i>Lb. sake-Lb. curvatus</i> , <i>P. pentosaceus</i> , <i>P. acidilactici</i> , <i>L. lactis</i>	Analysis of soluble proteins, amplified fragment length polymorphism	Kunene <i>et al.</i> , 2000
<i>Injera</i>	Ethiopia	<i>Candida guilliermondii</i> , <i>Pullaria</i> sp., <i>Aspergillus</i> sp., <i>Penicillium</i> sp., <i>Rhodotorulla</i> sp., <i>Homodendrum</i> sp., <i>Candida</i> sp., unidentified bacteria	Morphological and biochemical tests	Ashenafi, 1994; Dirar, 1993; Gashe <i>et al.</i> , 1982; Steinkraus, 1983; Vogel <i>et al.</i> , 1977
<i>Kisra</i>	Sudan	<i>Pediococcus pentosaceus</i> , <i>Lb. confusus</i> , <i>Lb. brevis</i> , <i>Lactobacillus</i> sp., <i>Erwinia ananas</i> , <i>Klebsiella pneumoniae</i> , <i>Enterobacter cloacae</i> , <i>Candida intermedia</i> , <i>Debaryomyces hansenii</i> , <i>Aspergillus</i> sp., <i>Penicillium</i> sp., <i>Fusarium</i> sp., <i>Rhizopus</i> sp.	Morphological and biochemical tests	Mohammed <i>et al.</i> , 1991
<i>Ogi</i> *	Nigeria	<i>Lactococcus raffinolactis</i> , <i>Lactococcus garviae</i> , <i>P. pentosaceus</i> , <i>Lb. paracasei</i> subsp. <i>tolerans</i> , <i>Lb. plantarum</i> , <i>Lb. suebicus</i> , <i>Lb. brevis</i> , <i>Pediococcus</i> sp., <i>Cephalosporium</i> sp., <i>Fusarium</i> sp., <i>Aspergillus</i> sp., <i>Penicillium</i> spp., <i>Saccharomyces cerevisiae</i> , <i>Candida mycorderma</i> , <i>C. vini</i>	API 50 CHL	Teniola <i>et al.</i> , 2005 ; Caplice and Fitzgerald, 1999

**Ogi* was prepared using maize.

In South Africa and Botswana, *ting* is used as a weaning food and is also consumed by adults (Boling and Eisener, 1982). This food represents a fermented sour porridge made traditionally by soaking sorghum flour for 2-3 days in water, followed by cooking. *Ting* is highly prized by local people for its unique taste, texture and aroma. Despite its popularity, there is little information regarding the microorganisms associated with the production of *ting*.

Togwa, a Tanzanian fermented gruel or beverage, is produced from sorghum, millet, maize, cassava or combinations of these (Mugula *et al.*, 2001). It is useful as a weaning food or as a refreshment (Mugula *et al.*, 2001). Like other indigenous African foods, the predominant microflora are LAB and yeasts (Table 1.1). The consumption of *togwa* is declining, probably due to the unhygienic nature of the preparation method and its poor shelf-life (Mugula *et al.*, 2001).

Bushera, a non-alcoholic fermented beverage, is the most common indigenous beverage produced in the western highlands of Uganda (Muyanja *et al.*, 2003). Production of *bushera* involves mixing germinated sorghum flour with boiled water, followed by cooling at ambient temperature and addition of germinated sorghum flour for starting the fermentation (Muyanja *et al.*, 2003). The fermentation is dominated by LAB and normally takes place in clay pots for 1-4 days (Muyanja *et al.*, 2003).

Ogi, a fermented sorghum, maize or millet gruel, is considered the most significant weaning food in West Africa and has been produced on a semi-industrial scale (Achi, 2005). *Ogi* has a sour flavour akin to that of yoghurt and a characteristic aroma, which differentiates it from other cereal-based fermented foods (Chavan and Kadam, 1989). Briefly, *ogi* is prepared by steeping the cereal grains in pots for 1-3 days, followed by wet milling and sieving of the fermented grains to yield a slurry (Steinkraus, 1998) of which the colour depends on the cereal grain. Fermentation of the grains is steered by LAB, yeasts and moulds, whilst flavour development is imparted by members of the genera *Saccharomyces* and *Candida* (Caplice and Fitzgerald, 1999; Table 1.1).

The common feature of many of these fermented foods is spontaneous fermentation, usually at household level. Despite the simple traditional method of preparation, natural fermentation involves a complex uncontrolled microbial process (Daeschel *et al.*, 1987), resulting in

products with variable sensory characteristics. Such variability in product quality hampers commercial production of indigenous fermented sorghum-based foods. Nevertheless, starter cultures may be used to overcome the inconsistency in product quality associated with traditional fermented foods by improving process control.

1.4 STARTER CULTURES

Starter cultures are microbial preparations containing large numbers of cells with at least one type of microbe, which shorten and steer the fermentation process when added to the raw material (Holzapfel, 2002; Leroy and De Vuyst, 2004). These cultures are critical for the formation of the desired flavour compounds that contribute to specific sensory characteristics. In addition, starter cultures pose technological, nutritional and health advantages (De Vuyst, 2000; Hansen, 2002). A typical starter culture facilitates improved control, predictability, safety, stability against mutations and standardization of a fermentation process (Holzapfel, 1997; Holzapfel, 2002). A starter culture may consist of single or mixed strains. Although single strain cultures may improve both process control and predictability of metabolic activities, they are relatively sensitive to bacteriophage infection, spontaneous mutation and they easily lose key physiological features such as plasmid-mediated fermentation of lactose (Holzapfel, 2002). In contrast, mixed strains are relatively unaffected by fluctuating conditions of handling and storage (Holzapfel, 2002).

Starter cultures may include moulds, yeasts, lactic acid bacteria or combinations of these. Moulds such as *Penicillium nalgiovense*, *Aspergillus oryzae* or *P. chrysogenum* are important starter cultures in food products such as cheese, fermented sausages, soy sauce and *miso*. Yeasts such as *Saccharomyces*, *Candida* and *Hansenula* species are important starter cultures for plant-based material that contain relevant fermentable sugars (Holzapfel, 1997), and are especially used in spontaneous African alcoholic fermentations. However, LAB occupy a central role as starter cultures due to their long safe history of interaction with humans. Hence, they are generally regarded as safe (GRAS) (Mugula *et al.*, 2003; Muyanja *et al.*, 2003).

1.5 TAXONOMY OF LACTIC ACID BACTERIA (LAB)

Taxonomy, a synonym of biosystematics, involves the cataloguing of biodiversity with the aim of arranging and characterizing organisms in an organized way (Staley and Kreig, 1984).

It is divided into classification, nomenclature and identification (Vandamme *et al.*, 1996). Classification is the orderly arrangement or clustering of organisms into taxonomic groups based on similarity, whilst nomenclature is the labeling of units according to international rules, and identification of organisms is the process of determining whether a query isolate belongs to one of the defined units (Staley and Kreig, 1989; Vandamme *et al.*, 1996). The species is the most significant and central element of bacterial taxonomy (Vandamme *et al.*, 1996) and has been described as a cluster of separate organisms, which are monophyletic, genomically coherent, and have a high degree of overall similarity with respect to independent characteristics (Rosselló-Mora and Amann, 2001). Bacterial species are usually identified using phenotypic traits, phylogenetic data or combinations of these.

Phenotypic classification usually involves carbohydrate fermentation patterns, lactic acid configuration and other growth requirements (Sharpe, 1981). Based on carbohydrate fermentation patterns, lactobacilli are classified as: (i) obligately homofermentative, (ii) facultatively heterofermentative, and (iii) obligately heterofermentative. The differences between the three fermentation types are as follow: obligate heterofermenters use the Emden-Meyerhof-Panas (EMP) pathway to convert hexose sugar to almost solely lactic acid, but not pentoses and gluconate due to the lack of phosphoketolase; facultative heterofermenters possess both aldolase and phosphoketolase, and can thus metabolize pentoses and gluconate in addition to hexoses, which are degraded using the EMP pathway; obligate heterofermenters metabolize pentoses, and hexoses are degraded via the phosphogluconate pathway with the formation of lactate, carbon dioxide, ethanol or acetic acid (Hammes and Vogel, 1995). Other phenotypic data may be obtained from the following: (i) expressed characteristics such as morphology, physiology, enzymology and serology, (ii) chemotaxonomic markers, which include cellular fatty acids and exopolysaccharides, among others, and (iii) proteins, including functionality (Vandamme *et al.*, 1996).

The classification of LAB, which is based on phenotypic traits and true phylogenetic relationships inferred from rRNA gene sequences (see below), remains largely unresolved due to lack of correlation between these approaches (Vandamme *et al.*, 1996; Felis and Dellaglio, 2007). For instance, the genus *Lactobacillus* is phylogenetically intermixed with some *Pediococcus* and *Leuconostoc* strains, despite their traditional separation due to differences in morphology (Stackebrandt and Teuber, 1988; Axelsson, 2004). In addition, phenotypic tests mostly fail to distinguish the genus *Carnobacterium* from that of *Lactobacillus*, and

Vagococcus from *Lactococcus* members (Axelsson, 2004). Despite this lack of correlation in some groups between phenotypic and genotypic results, phenotyping generally provides important information for separating taxa (Botina *et al.*, 2006). In order to circumvent limitations inherent in either phenotyping or phylogeny, polyphasic taxonomy, involving an integration of phenotypic, genotypic and phylogenetic data, is important (Colwell, 1970; Vandamme *et al.*, 1996). However, if the main thrust is speed and reliability, DNA-based techniques are more advantageous (Ehrmann and Vogel, 2005).

1.5.1 General characteristics of LAB

LAB represent a diverse and paraphyletic group of bacteria (Fig. 1.2). They constitute a heterogeneous group of Gram-positive, acid-tolerant and strictly fermentative cocci, coccobacilli or rods that produce lactic acid as the main product during carbohydrate fermentation (Axelsson, 2004; Temmermann *et al.*, 2004; Ehrmann and Vogel, 2005). The heterogeneity is clearly evidenced in the genus *Weissella* that contains both coccoid and rod-shaped members (Collins *et al.*, 1993). Furthermore, the largest and most heterogeneous group within the LAB group comprise lactobacilli with a G + C content of 32-55%, which is wider than that expected for members of a single genus (Goodfellow *et al.*, 1997). Generally, the G + C content do not vary more than 3 and 10% for a well-defined species and genus level, respectively (Stackebrandt and Liesack, 1993).

LAB generally lack catalase, although pseudocatalase was detected in cultures grown at low sugar concentrations (Caplice and Fitzgerald, 1999). They are asporogenous and fastidious organisms and hence, occupy nutritionally rich habitats. LAB are mesophilic and their growth temperatures range from 5°C to 45°C (Caplice and Fitzgerald, 1999). This variability also applies to pH where LAB grow at pH 9.6 and pH 3.6, with the majority growing at pH 4.0-4.5 (Caplice and Fitzgerald, 1999).

1.5.2 Horizontal gene transfer (HGT) and LAB

The common ancestor of LAB was shaped by HGT (Koonin *et al.*, 2001), like all lineages of bacteria (Doolittle, 1999; Jordan *et al.*, 2001; Brown, 2003; Koonin, 2003; Lawrence and Hendrickson, 2003; Ochman, 2005; Choi and Kim, 2007). HGT is the transfer of genes among organisms of different species either by acquiring paralogous genes that already exist in a specific lineage, acquiring genes that do not pre-exist in the lineage or through




Figure 1.2 Neighbor-joining phylogenetic tree, based on 16S rRNA gene sequence analysis, showing the relationships among type species of the representative genera from the families of *Lactobacillales*. These genera represent the families *Lactobacillaceae*, *Aerococcaceae*, *Carnobacteriaceae*, *Enterococcaceae*, *Leuconostoccaceae* and *Streptococcaceae* and non-*Lactobacillales*, *Listeria monocytogenes*. *Bacillus subtilis* is the outgroup. The bar indicates the number of nucleotide substitutions per site.

xenologous gene displacement (Koonin *et al.*, 2001). In LAB, gene acquisition is thought to occur by conjugation or it is facilitated by bacteriophages (Wood and Warner, 2003). Consequently, duplication, acquisition and/or loss of genes occurred as these organisms underwent evolution (Makarova and Koonin, 2007). It was inferred that the ancestor of LAB might have acquired about 84 genes from various sources through HGT (Makarova and Koonin, 2007). This gene acquisition, especially of a group of peptidases, allows LAB to adapt and out-compete other microorganisms found in complex and nutritionally rich environments such as soil, silage, sewage, manure, water, plant materials, and mucous membranes such as the intestines, vagina and mouth where LAB normally reside (Axelsson, 2004; Makarova and Koonin, 2007).

HGT also influences the taxonomy of LAB. For instance, when a limited number of gene families are used for inferring phylogeny (see below), the topology of the gene trees are not concordant with species trees (Smith *et al.*, 1992; McGowan *et al.*, 1998; Friedrich, 2002; Ge *et al.*, 2005). However, when either small subunit (SSU) rRNA genes, whole genomes or numerous housekeeping genes are used for phylogenetic analysis of organisms, HGT will have very little influence on the phylogenetic tree (Kurland, 2000; Daubin *et al.*, 2003; Kurland *et al.*, 2003; Snel *et al.*, 2005; Choi and Kim, 2007). In other instances, HGT might result in unusual similarity of a gene originating from different genomes in comparison to orthologous genes (Koonin *et al.*, 2001). The continuous genome decay through gene loss in LAB such as *Streptococcus thermophilus* explains the ongoing evolving taxonomy of these bacteria (Bolotin *et al.*, 2004).

HGT plays an important part in LAB technologies, since genetic material, usually plasmids containing some characteristics, may be easily transferred among LAB, thus leading to the production of new genetic systems (Dalezios and Siebert, 2001). For instance, transferring a conjugative transposon with genes that encode for the production of nisin, an industrially important bacteriocin, to non-nisin producing strains has a huge industrial impact (Nga, 2005). Recombinant strains of *L. lactis* have been used for expression of bovine β -lactoglobulin (Chatel *et al.*, 2001) and for supplementation of lipase in pigs with pancreatic lipase deficiency (Drouault *et al.*, 2002). The production of the flavour compound diacetyl by some LAB from citrate is mediated by citrate permease. The gene that codes for this enzyme is linked to a small plasmid (Kempler and McKay, 1981). Hence, recombinant strains that are deficient for this gene may not be used for production of diacetyl.

1.5.3 The order *Lactobacillales*

The order *Lactobacillales* belongs to the phylum Firmicutes, class Bacilli and comprises the LAB. Together with the *Lactobacillaceae*, this order includes five other families, as well as a number of species not currently classified in any known family (Table 1.2) (Garrity *et al.*, 2001). The total number of genera within these families is 34 (Garrity *et al.* 2004; Table 1.2), with the following being important fermenters or spoilers of fermented products: *Lactobacillus*, *Aerococcus*, *Carnobacterium*, *Enterococcus*, *Lactococcus*, *Leuconostoc*, *Oenococcus*, *Pediococcus*, *Streptococcus*, *Tetragenococcus*, *Vagococcus* and *Weissella* (Axelsson, 2004). The valid species included in each of these genera are shown in Appendices A and B.

The family *Lactobacillaceae* consists of three genera, namely *Lactobacillus*, *Paralactobacillus* and *Pediococcus* (Garrity *et al.*, 2004). The genus *Lactobacillus* currently contains 106 species, making it the largest group in the order (Felis and Delaglio, 2007; Appendix A). The genus *Pediococcus* contains 11 members. The genus *Paralactobacillus* contains only one species, *Paralactobacillus selangolensis* (Leisner *et al.*, 2000), which was isolated from *chili bo*, a Malaysian food ingredient (Leisner *et al.*, 2000).

The family *Leuconostocaceae* consists of three genera, namely *Leuconostoc*, *Oenococcus* and *Weissella*. *Leuconostoc*, the largest of these, has 23 members and some of them are associated with food (Shaw and Harding, 1989; Antunes *et al.*, 2002; Kim *et al.*, 2003; Chambel *et al.*, 2006). *Weissella*, the second largest genus in this family, has 12 members. Although more members of this genus have been isolated from fermented food (Collins *et al.*, 1993; Choi *et al.*, 2002), some *Weissella* species have been detected in clinical samples (Björkroth *et al.*, 2002; Shin *et al.*, 2007). The genus *Oenococcus* consists of only two members, *Oenococcus oeni* and *O. kitaharae* (Appendix B).

The genera *Streptococcus* and *Lactococcus* make up the family *Streptococcaceae* (Appendix B). Of the 97 *Streptococcus* members, *Streptococcus thermophilus* is of industrial significance and it is used during the manufacture of a variety of fermented milk products. Most of the members in these genera are associated with a variety of diseases, both in humans and animals (Smith and Sherman, 1939; Bouvet *et al.*, 1989; Whiley *et al.*, 1999). The genus *Lactococcus* has six members with *Lactococcus lactis* playing a central role in the industrial production of various compounds such as nisin and diacetyl.

Table 1.2 Families and genera in the order *Lactobacillales**

Family	Genera
<i>Lactobacillaceae</i>	<ul style="list-style-type: none"> • <i>Lactobacillus</i> • <i>Paralactobacillus</i> • <i>Pediococcus</i>
<i>Aerococcaceae</i>	<ul style="list-style-type: none"> • <i>Aerococcus</i> • <i>Abiotrophia</i> • <i>Dolosicoccus</i> • <i>Eremococcus</i> • <i>Facklamia</i> • <i>Globicatella</i> • <i>Ignavigranum</i>
<i>Carnobacteriaceae</i>	<ul style="list-style-type: none"> • <i>Carnobacterium</i> • <i>Agitococcus</i> • <i>Alkalibacterium</i> • <i>Allofustis</i> • <i>Alloicoccus</i> • <i>Desemzia</i> • <i>Dolosigranulum</i> • <i>Isobaculum</i> • <i>Lactosphaera</i> • <i>Marinalactibacillus</i> • <i>Trichococcus</i> • <i>Granulicatella</i>
<i>Leuconostocaceae</i>	<ul style="list-style-type: none"> • <i>Leuconostoc</i> • <i>Oenococcus</i> • <i>Weissella</i>
<i>Streptococcaceae</i>	<ul style="list-style-type: none"> • <i>Streptococcus</i> • <i>Lactococcus</i>
<i>Enterococcaceae</i>	<ul style="list-style-type: none"> • <i>Enterococcus</i> • <i>Atopobacter</i> • <i>Mellisococcus</i> • <i>Tetragenococcus</i> • <i>Vagococcus</i>
<i>Incertae sedis</i>	<ul style="list-style-type: none"> • <i>Acetoanerobium</i> • <i>Oscillospira</i> • <i>Syntrophococcus</i>

*Adopted from: Garrity *et al.*, 2004

Enterococcus, *Atopobacter*, *Mellisococcus*, *Tetragenococcus* and *Vagococcus* genera belong to the family *Enterococcaceae* (Garrity *et al.*, 2004). The largest of these genera *Enterococcus* consists of 37 members that are associated with both foods and clinical samples (Fortina *et al.*, 2004; Koort *et al.*, 2004). The remaining genera consist of a total of eight members (Garrity *et al.*, 2004).

The family *Aerococcaceae* consists of seven genera (Garrity *et al.*, 2004; Table 1.2). The largest genus is *Facklamia*, which consists of six species (Appendix B). The genera *Abiotrophia* and *Aerococcus* have five members each. *Globicatella* genus has two members, whilst *Dolosicoccus*, *Eremococcus* and *Ignavigranum* consist of only one member per genus (Garrity *et al.*, 2004; Appendix B).

The family *Carnobacteriaceae* consists of 12 genera (Garrity *et al.*, 2004; Appendix B), with a total of 24 members. The largest genus *Carnobacterium* consists of eight members, whilst the remainder contains one or two members only (Appendix B). Some members of *Carnobacteria* are associated with food spoilage (Holley *et al.*, 2002).

1.6 IDENTIFICATION OF LAB ASSOCIATED WITH FERMENTED FOODS

1.6.1 Phenotypic methods

Phenotyping of bacteria involves all the identification methods that are not based on DNA or RNA (Caplice and Fitzgerald, 1999). These include morphology (e.g., shape, presence or absence of endospores or inclusion bodies), physiology and biochemical attributes such as conditions required for growth, activities of enzymes and metabolic activities. The significance of individual phenotypic characteristics may appear insignificant for determining genetic relatedness, yet, when taken together, taxa may be determined using these attributes (Vandamme *et al.*, 1996).

For routine phenotyping, miniaturized identification kits such as API 50 CHL or BIOLOG, which are based on carbohydrate fermentation profiles, are often popular due to affordability (Temmerman *et al.*, 2004). Although this approach is simple, affordable and provides evidence for functionality of strains, phenotyping is generally laborious, time-consuming and the efficiency and resolution may be poor, especially when complex samples with high diversity are studied (Meroth *et al.*, 2003). For instance, most LAB isolates from freshwater

fish could not be identified, even at genus level, due to the poor taxonomic resolution of phenotypic methods (Gonzalez *et al.*, 2000). In addition, only 38% of LAB from sourdough could be identified using the API 50 CHL system, as well as morphological and physiological characteristics (Corsetti *et al.* 2001). The fact that LAB strongly adapt to their environment and share many common attributes, makes differentiation of species using phenotypic methods challenging (e.g., Hayford *et al.*, 1999; Van Reenen and Dicks, 1996). Moreover, it is challenging to compare results obtained using different phenotypic methods and similar phenotypes may not be concordant with related genotypes (Temmerman *et al.*, 2004). Therefore, it is essential to use an approach that combines phenotyping and fast, reliable and reproducible DNA-based techniques with high resolution for identification of LAB associated with food fermentations (Vandamme *et al.*, 1996; Nigatu, 2000).

1.6.2 DNA-based methods

LAB are characterized using a wide variety of DNA-based methods (Fig. 1.3). One DNA-based method that will allow identification of LAB to the species level is species-specific PCR (Nomura *et al.*, 2002). Species-specific PCR involves the amplification of target genes, using primers that correspond to the oligonucleotide sequences present within parts of the whole genome of target organisms (Lupski and Weinstock, 1992), followed by subsequent analysis. Species-specific PCR targeting the 16S rRNA gene was used for identification of *Lb. brevis* and the method was found to be very sensitive, specific and efficient (Guarneri *et al.*, 2001).

DNA-based methods that allow identification of LAB to the species and strain level include restriction fragment length polymorphism (RFLP) (Giraffa *et al.*, 2002), pulsed-field gel electrophoresis (PFGE) (Ventura and Zink, 2002), randomly amplified polymorphic DNA-PCR (RAPD-PCR) (Williams *et al.*, 1990), repetitive extragenic palindromic-PCR (rep-PCR) (Versalovic *et al.*, 1991), amplified ribosomal DNA restriction analysis (ARDRA), amplified fragment length polymorphism (AFLP) (Vos *et al.* 1995) and phylogenetic analyses of housekeeping loci.




Figure 1.3 Main steps involved in DNA-based methods for characterization of LAB. R.E. represents restriction enzyme. (Adopted and modified from: Olive and Bean, 1999).

RFLP involves restriction enzyme digestion of the whole-genome, followed by size fractionation of the resultant DNA fragments by agarose gel electrophoresis. Southern blotting (Southern, 1975) may be used to transfer DNA fragments onto either a nitrocellulose or nylon membrane, followed by hybridization to a labeled homologous probe (e.g., rRNA gene probe), which allows visualization of the hybridized fragments (Vandamme *et al.*, 1996). The hallmark of RFLP is high resolution up to strain level.

In ARDRA, bacterial rRNA genes are initially amplified by PCR using conserved sequences of rRNA genes as primers and the amplicon is digested with restriction endonucleases, followed by electrophoresis to obtain a fingerprint (Massol-Deya *et al.*, 1995). ARDRA is fast to perform, but highly dependent on the degree of polymorphism in the region studied and the discriminatory power is often inferior (Massol-Deya *et al.*, 1995) because smaller areas of the rRNA operon are targeted. Roy *et al.* (2001) used ARDRA for molecular discrimination of *Lb. helveticus*, *Lb. delbrueckii* subspecies *delbrueckii*, *Lb. delbrueckii* subsp. *lactis*, *Lb. delbrueckii* subsp. *bulgaricus*, *Lb. acidophilus* and *Lb. casei*, making it a suitable technique for discrimination of these closely related bacteria at subspecies level.

PFGE involves separation of large DNA representing the whole genome in an oscillating electric field (Tenover *et al.*, 1995; McCartney, 2002). Excellent subspecies discrimination is the hallmark of PFGE. For instance, Ventura and Zink (2002) observed heterogeneity among isolates of *Lb. johnsonii* after using PFGE, despite the close relationship between these organisms. However, since the extraction of intact chromosomal DNA is critical to PFGE, the technique becomes labourious and expensive. Therefore, PFGE it is seldomly applied in studies where large numbers of isolates are to be characterized (Temmerman *et al.*, 2004).

In RAPD-PCR fingerprinting, one or two primers (usually 10-12 bp) are arbitrarily selected and allowed to anneal to the bacterial genomic DNA template at low stringency, resulting in several amplification products of varying sizes, which are resolved electrophoretically to yield a RAPD fingerprint (Welsh and McClelland, 1990). RAPD analyses do not require prior knowledge of the target sequences. It has been used to follow unmarked starter cultures in commercial fermentations (Plengvidhya *et al.*, 2004). Reguant and Bordons (2003) developed a multiplex RAPD-PCR, based on the combination of one random 10-mer and one specific 23-mer oligonucleotide in a single PCR. The method generated unique patterns and distinguished DNA profiles of strains of *Oenococcus oeni*. RAPD-PCR is simple and fast in

methodological terms, but its reproducibility is limited due to the low stringency of the PCR amplification and sensitivity to small variations in template concentration, purity, magnesium concentration and primer annealing temperature (Erlandson and Batt, 1997). Therefore, RAPD-PCR is best suited for studies where a specific bacterial strain is sought among a large number of isolates, and it is not well suited for inter-laboratory and taxonomic studies where the aim is to develop a fingerprint database.

Repetitive DNA elements allow the analysis of whole-genomes through DNA band profiles created using PCR-based techniques such as ERIC (enterobacterial repetitive intergenic consensus) sequences for typing enterobacteria (Hulton *et al.*, 1991), BOX for typing *Streptococcus pneumoniae* using invertedly repeated DNA elements (Martin *et al.*, 1992) and rep (repetitive extragenic palindromic) elements. In rep-PCR (Versalovic *et al.*, 1991), primers anneal to repetitive parts of the chromosome and amplification occurs when the distance between the primer-binding sites are short enough to enable amplification. The advantages are high resolution (Ventura and Zink, 2002), making it suitable for analyzing closely related LAB using a (GTG)₅ primer, low cost, reliability (Olive and Bean, 1999) and high throughput (Gevers *et al.*, 2001). In addition, standardization and inter-laboratory comparisons are feasible due to automation of rep-PCR with fluorescently labeled primers. However, very few publications are available. Bouton *et al.* (2002) used rep-PCR, RAPD-PCR and PFGE to identify selected starter strains during cheese ripening and showed that *Lb. helveticus* diminished fast, while *Lb. delbrueckii* subsp. *lactis* maintained viability during ripening.

Amplified fragment length polymorphism (AFLP) involves restriction enzyme digestion of DNA using two restriction enzymes, resulting in fragments with sticky ends. Prior to selective PCR-amplification of the fragments, adapters, which are short oligonucleotides, are ligated to the sticky ends (Zabeau and Vos, 1993). The subsequent PCR amplicons are then separated by polyacrylamide gel electrophoresis. The hallmark of AFLP is the ability to differentiate isolates at species level and below. Kunene *et al.* (2000) distinguished *Lb. plantarum* and *L. mesenteroides* isolated from different sorghum doughs using AFLP (Kunene *et al.*, 2000).

Phylogenetic analyses of conserved genomic regions are increasingly being used for the identification of bacteria. Phylogenetic relationships among species are usually inferred using

rRNA genes due to the following: (i) ubiquitous nature of rRNA genes in bacteria, (ii) presence of highly conserved regions due to the essential role played by ribosomes during protein synthesis, (iii) presence of variable regions within the sequence, (iv) availability of large public domain databases (e.g., the Ribosomal Database), which enable sequence comparisons, and (v) rRNA-encoding genes are not usually influenced by horizontal gene transfer (HGT) (Woese, 1987; Axelsson, 2004; Felis and Dellaglio, 2007). However, the limitation of rRNA genes for determining phylogenetic relationships is that generally a sequence identity of less than 97% indicates a poor relationship of organisms at the genomic level (Felis and Dellaglio, 2007). Furthermore, ribosomal gene sequences may not always allow clear-cut LAB identifications as these genes are highly conserved (Felis and Dellaglio, 2007; De Vuyst and Vancanneyt, 2007). As is the case for many other bacterial taxa, LAB are known to encode 16S rRNA gene sequences that are not sufficiently polymorphic to allow species separation (Torriani *et al.*, 2001; Gevers *et al.*, 2005; Felis and Dellaglio, 2007). Microorganisms are still described only as closely related even if they share identical rRNA sequences. Therefore, bacterial species are increasingly defined using multilocus sequence analysis (MLSA), involving housekeeping loci that evolve more rapidly than 16S rRNA genes (Gevers *et al.*, 2005; Kostantinidis and Tiedje, 2007).

1.7 DNA-BASED METHODS FOR STUDYING LAB COMMUNITIES

Microbial population dynamics is studied using either culture-dependent or culture-independent techniques. The limitations of culture-dependent methods for such studies are well-documented (Ampe *et al.*, 1999; Walter *et al.*, 2001; Randazzo *et al.*, 2002). Culture-dependent methods may underestimate microbial diversity due to failure of some organisms to grow. For instance, culture-dependent methods failed to detect *Lb. delbrueckii*, despite its dominance during cheese ripening (Randazzo *et al.*, 2002). Similarly, *Carnobacterium* sp. was not identified in cold-smoked salmon due to inhibition of its growth by sodium acetate in de Mann, Sharpe and Rogosa (MRS) medium (Giacomazzi *et al.*, 2004). In addition, culture-dependent methods may overestimate microbial diversity due to non-selectivity of the culture media (Randazzo *et al.*, 2002). Therefore, culture-independent genetic fingerprinting techniques are the methods of choice since they are easy to perform and are reproducible. These techniques include PCR-denaturing gradient gel electrophoresis (PCR-DGGE; Muyzer *et al.*, 1993), single-stranded conformation polymorphism (SSCP; Orita *et al.*, 1989), and terminal-restriction fragment length polymorphism (T-RFLP; Avannis-Aghajani *et al.*, 1994;

Fig. 1.4). Of these culture-independent techniques, PCR-DGGE is a frequently used method (Ercolini, 2004).

Despite their numerous advantages, culture-independent techniques with respect to food matrices are limited by DNA extraction and PCR amplification biases (Von Wintzingerode *et al.*, 1997; Ercolini, 2004). DNA extraction of bacteria in complex ecosystems may be complicated due to differences in cell structure among varying taxa, leading to bias in extraction in favour of cells that are easily disrupted (Niemi *et al.*, 2001; Ercolini, 2004). Niemi and co-workers (2001) concluded that PCR-DGGE profiles are a function of the method used for DNA isolation and purification. This bias in efficiency of DNA extraction may cause further bias in downstream applications such as PCR amplification (Ercolini, 2004). PCR amplification may be biased due to preferential amplification of target genes (Reysenbach *et al.*, 1992). Consequently, some microorganisms in a complex ecosystem are excluded (Ercolini, 2004). Therefore, in order to obtain a true reflection of the microbial population dynamics during food fermentations, a holistic approach that combines both culture-dependent and culture-independent methods is important.

In DGGE, total bacterial DNA from the habitat of interest is extracted and a variable region of the 16S rRNA gene is PCR-amplified (Fischer and Lerman, 1983; Muyzer *et al.*, 1993; Fig. 1.4). DNA amplicons of the same length but containing different DNA sequences are subjected to polyacrylamide gel electrophoresis by using a denaturing gradient established with urea and formamide. DGGE enables direct visualization of bacterial diversity and subsequent identification of community members by DNA fragment sequence analysis or hybridization with specific probes (Muyzer *et al.*, 1993). Therefore, it is a suitable tool for rapid and economical investigation of microbial community dynamics during whole-food fermentation processes (ben Omar and Ampe, 2000; Meroth *et al.*, 2003). Well-defined microflora produced by using starter cultures in controlled conditions are suitable for DGGE analysis, as the food substrates usually display a simple profile where each band corresponds to the expected species (Ercolini *et al.*, 2004).




Figure. 1.4 Main steps involved in DNA-based methods for studying microbial population dynamics of LAB.

Many studies involving microbial population dynamics of LAB during food fermentations have employed PCR-DGGE. Two LAB-specific PCR-DGGE systems were used to monitor the development of starter mixtures of three commercially available sourdough starters and a baker's yeast preparation, containing various species of LAB, and *Lb. sanfranciscensis*, *Lb. pontis*, *Lb. mindensis*, *Lb. crispatus*, *Lb. johnsonii*, *Lb. frumenti* and *Lb. reuteri* were shown to be the predominant species (Meroth *et al.*, 2003). Randazzo *et al.* (2002) examined microbial succession during the manufacture of artisanal Sicilian cheese using PCR-DGGE and cultivation. For total microflora, the V6-V8 variable region of the 16S rRNA gene was analyzed, whilst species-specific primers targeting the V1-V3 region were used to detect *Lactobacillus* spp. *Streptococcus thermophilus* prevailed during fermentation, whilst mesophilic bacteria were dominant in raw milk after cloning and sequencing of the 16S rRNA gene amplicons.

The major advantages of DGGE are reproducibility, reliability and speed (Muyzer, 1999). However, DGGE has particular limitations. For instance, PCR-DGGE failed to detect *Lb. acidophilus* and *Lb. casei*, declared in the probiotic yoghurt products Biospega and ABC, due to bacterial concentrations below the detection limit (Fasoli *et al.*, 2003). This may also be due to the choice of PCR primers, as these are crucial for determining the resolution of DGGE (Temmerman *et al.*, 2004). However, group-specific primers may improve the detection limit for less abundant groups in the microbial ecosystem. Furthermore, a single band may represent more than one strain in DGGE (Muyzer and Smalla, 1998; Sekiguchi *et al.*, 2001; Van Beek and Priest, 2001; Cocolin *et al.*, 2001; Meroth *et al.*, 2003), a phenomenon known as co-migration. For instance, *Lb. manihotivorans*, which represented up to 13% of the total LAB of sour cassava starch, was not detected by DGGE as the PCR product migrated at the same position as that of *L. lactis* (Ampe *et al.*, 2001). This may be circumvented by using a set of different primer pairs or varying the DGGE gradient (Ercolini, 2004). In addition, targeting the 16S rRNA gene to study community structure may overestimate species diversity in DGGE profiles due to the heterogeneity of this region (Farrelly *et al.*, 1995; Fogel *et al.*, 1999). However, targeting single copy genes such as *rpoB*, encoding the RNA polymerase beta subunit, may circumvent this limitation (Dahllöf *et al.*, 2000; Giacomazzi *et al.*, 2004). DGGE may also be biased by the failure of PCR to distinguish between viable and non-viable cells (Uyttendaele *et al.*, 1999). In some instances, PCR detected cells after autoclaving at 121°C for 15 min (Masters *et al.*, 1994).

In SSCP (Orita *et al.*, 1989), a single base modification can change the conformation of a single-strand DNA molecule, altering the migration distance of the molecules in a non-denaturing gel. Hence, sequence variations between DNA fragments of the same size (Hayashi, 1991; Fig. 1.4), which are usually PCR-amplified from variable regions of the 16S rRNA gene, are detected (Giraffa and Neviani, 2001). Duthoit *et al.* (2003) used SSCP analysis targeting the V2 or V3 regions of the 16S rRNA gene to study the dynamics of the microbial flora of Salers cheese without the addition of starter culture. The authors found *L. lactis*, *S. thermophilus*, *E. faecium*, *Le. mesenteroides*, *Le. pseudomesenteroides*, *Lb. plantarum* and *Lb. pentosus* to be predominant during manufacturing and ripening. In contrast to DGGE, SSCP gives at least two bands (one for each strand) and often a third one (the re-annealed dsDNA product) that represents one DNA sequence, making interpretation of the results complex, especially for studies that involve community analysis of microorganisms. The major advantage of SSCP is that it is rapid to perform and automatization of the technique by capillary electrophoresis allows microbial dynamics of many samples to be analyzed (Duthoit *et al.*, 2003). The discriminatory power and reproducibility of SSCP analysis is usually most effective for fragments up to 400 bp (Vaneechoutte, 1996). PCR-SSCP detects bacterial populations that make up 1% or more of a bacterial community (Lee *et al.*, 1996).

In T-RFLP, the target gene is amplified from the community DNA using standard PCR techniques with a 5' fluorescently tagged primer (Avaniss-Aghajani *et al.*, 1994; Liu *et al.*, 1997; Fig. 1.4), and the amplification products are digested with restriction enzymes and electrophoresed on an automated sequencer. Only the restriction fragment containing labeled primer is detected by the automated system. In general, each population of the community contributes a terminal fragment of one size (Moeseneder *et al.*, 1999; Osborn *et al.*, 2000; Dunbar *et al.*, 2001). Hence, direct comparisons can be made rapidly and easily using the large sequence database of the Ribosomal Database Project (Marsh *et al.*, 2000; Dunbar *et al.*, 2001). However, there is a possibility that two different species could have the same restriction site in their 16S rRNA gene sequence, resulting in an identical peak (Moeseneder *et al.*, 1999). Nevertheless, the use of different enzymes increases information for discriminating samples (Moeseneder *et al.*, 1999). T-RFLP of amplified 16S rRNA genes was successfully used to study the dynamics of metabolically active mixed dairy cultures consisting of *L. lactis* subsp. *lactis* and *Le. citreum* (Sánchez *et al.*, 2006).

1.8 CONCLUSIONS AND AIMS OF THIS INVESTIGATION

In summary, most studies involving characterization of LAB isolated from fermented sorghum foods have focussed on phenotypic methods. However, if traditional sorghum-based fermented foods are to be upgraded to commercial status, whilst maintaining their unique pleasant sensory attributes, characterization of wild-strain LAB as potential starter cultures is of paramount importance. The limitations of phenotyping are well documented. Various DNA-based methods offer a rapid and more reliable alternative. However, the inherent limitations in each technique necessitate the need for a polyphasic approach. To select the most suitable starter cultures, which can adapt rapidly to the sorghum matrix and dominate fermentations, it is essential to study the dynamics of the microflora involved. A combination of both culture-dependent and culture-independent approaches may give an overview of the diversity and microbial population dynamics occurring in sorghum fermentations.

Although PFGE has excellent resolution, routine use for characterization of LAB during food fermentations is limited by the labourious nature of the technique. The use of RAPD-PCR as a method for characterization of LAB in the food industry is limited by reproducibility challenges. However, rep-PCR, which is cost-effective and reliable, has a high resolution and can be used to analyze complex genomes (Ventura and Zink, 2002). This may be an important technique in future for characterization of LAB isolated from indigenous fermented sorghum-based foods since it was shown to have good correlation with PFGE results, albeit with slightly less resolution.

Predominant and dominant wild-strain LAB starter cultures used in most trial fermentations of sorghum-based foods have been selected based on methods that relied on culture-dependent approaches. However, it is imperative in future to include culture-independent methods for such studies in order to obtain a true reflection of LAB that dominate sorghum fermentations since some microorganisms may be excluded or under-represented by cultivation on particular media. PCR-DGGE is a powerful tool for microbial population studies involving food fermentations, and it has been used to monitor microbial composition and dynamics during food fermentations (Randazzo *et al.*, 2002; Meroth *et al.*, 2003). Its current limitations, however, necessitate the use of PCR-DGGE in combination with other techniques. Therefore, an approach that combines both culture-dependent and culture-

independent methods to give an overview of microbial population dynamics in indigenous sorghum-based foods is recommended. Consequently, the aims of this study were:

- To characterize bacteria isolated from spontaneous sorghum fermentations using DNA-based techniques.
- To perform PCR-DGGE on total bacterial genomic DNA obtained from two types of commercial sorghum flour spontaneously fermented at 20, 25 and 30°C, so as to determine the diversity and microbial dynamics that occur during sorghum fermentations.
- To clarify the role of selected starter cultures of LAB with regards to the microbiological safety and consumer acceptance of the sensory characteristics of *ting*.

1.9 REFERENCES

- Abd El-Salam, M.H. and Alichanidis, E.** (2004). Cheese varieties ripened in brine. In: *Cheese: Chemistry, physics and microbiology*. Fox *et al.* (Eds.). Elsevier, Amsterdam, The Netherlands, pp. 227-249.
- Achi, O.K.** (2005). The potential of upgrading traditional fermented foods through biotechnology. *African Journal of Biotechnology*, **4**, 375-380.
- Adams, M.R.** (1990). Topical aspects of fermented foods. *Trends in Food Science and Technology*, **1**, 141-144.
- Adams, M.R. and Nicolaidis, L.** (1997). Review of the sensitivity of different food-borne pathogens to fermentation. *Food Control*, **8**, 227-239.
- Aguirre, M. and Collins, M.D.** (1993). Lactic acid bacteria and human clinical infection. *Journal of Applied Bacteriology*, **75**, 95-107.
- Akingbala, J.O., Rooney, L.W. and Faubion, J.M.** (1981). A laboratory procedure for the preparation of *ogi*, a Nigerian fermented food. *Journal of Food Science*, **46**, 1523-1526.
- Ampe, F., ben Omar, N., Moizan, C., Wach, C. and Guyot, J.P.** (1999). Polyphasic study of the partial distribution of microorganisms in Mexican *pozol*, fermented maize dough demonstrates the need for cultivation-independent methods to investigate traditional fermentations. *Applied and Environmental Microbiology*, **65**, 5464-5473.

- Ampe, F., Sirvent, A. and Zakhia, N.** (2001). Dynamics of the microbial community responsible for the traditional sour cassava starch fermentation studied by denaturing gradient gel electrophoresis and quantitative rRNA hybridization. *International Journal of Food Microbiology*, **65**, 45-54.
- Antunes, A., Rainey, F.A., Nobre, M.F., Schumann, P., Ferreira, A.M., Ramos, A., Santos, H. and Da Costa, M.S.** (2002). *Leuconostoc ficulneum* sp. nov., a novel lactic acid bacterium isolated from a ripe fig, and reclassification of *Lactobacillus fructosus* as *Leuconostoc fructosum* comb. nov. *International Journal of Systematic and Evolutionary Microbiology*, **52**, 647-655.
- Ashenafi, M.** (1994). Microbial flora and some chemical properties of *ersho*, a starter for *teff* (*Eragrostis tef*) fermentation. *World Journal of Microbiology and Biotechnology*, **10**, 69-73.
- Au, P.M. and Fields, M.L.** (1981). Nutritive quality of fermented sorghum. *Journal of Food Science*, **46**, 652-654.
- Avaniss-Aghajani, E., Jones, K., Chapman, D. and Brunk, C.** (1994). A molecular technique for the identification of bacteria using small subunit ribosomal RNA sequences. *BioTechniques*, **17**, 144-149.
- Axelsson, L.** (2004). Lactic acid bacteria: Classification and physiology. In: *Lactic acid bacteria. Microbiological and functional aspects*. Salimen, S., von Wright, A. and Ouwehand, A. (Eds.). Marcel Dekker, Inc., New York, USA, pp. 1-66.
- Axtell, J.D., Kirleis, A., Hassen, M.M., D’Croz-Mason, N., Mertz, E.T. and Munck, L.** (1981). Digestibility of sorghum proteins. *Proceedings of the National Academy of Sciences of the United States of America*, **78**, 1333-1335.
- ben Omar, N. and Ampe, F.** (2000). Microbial community dynamics during production of Mexican fermented dough *pozol*. *Applied and Environmental Microbiology*, **66**, 3664-3673.
- Björkroth, K.J., Schillinger, U., Geisen, R., Weiss, N., Hoste, B., Holzapfel, W.H., Korkeala, H.J. and Vandamme, P.** (2002). Taxonomic study of *Weissella confusa* and description of *Weissella cibaria* sp. nov., detected in food and clinical samples. *International Journal of Systematic and Evolutionary Microbiology*, **52**, 141-148.
- Blandino, A., Al-Aseeri, M.E., Pandiella, S.S., Cantero, D. and Webb, C.** (2003). Cereal-based fermented foods and beverages. *Food Research International*, **36**, 527-543.

Boling, M.B. and Eisener, N. (1982). *Bogobe*: Sorghum porridge of Botswana. In: Proceedings of the International Symposium on Sorghum Grain Quality. International Crops Research Institute for the Semi-Arid Tropics. Mertin, J.V. (Ed.). Patancheru, India, pp. 32-35.

Bolotin, A., Quinquis, B., Renault, P., Sorokin, A., Ehrlich, S.D., Kulakauskas, S., Lapidus, A., Goltsman, E., Mazur, M., Pusch, G.D., Fonstein, M., Overbeek, R., Kyprides, N., Purnelle, B., Prozzi, D., Ngui, K., Masuy, D., Hancy, F., Burteau, S., Boutry, M., Delcour, J., Goffeau, A. and Hols, P. (2004). Complete sequence and comparative genome analysis of the dairy bacterium *Streptococcus thermophilus*. *Nature Biotechnology*, **22**, 1554-1558.

Bouvet, A., Grimont, F. and Grimont, P.A.D. (1989). *Streptococcus defectivus* sp. nov. and *Streptococcus adjacens* sp. nov., nutritionally variant streptococci from human clinical specimens. *International Journal of Systematic Bacteriology*, **39**, 290-294.

Botina, S.G., Tsygankov, Y.D. and Sukhodolets, V.V. (2006). Identification of industrial strains of lactic acid bacteria by methods of molecular genetic typing. *Russian Journal of Genetics*, **42**, 1367-1379.

Bottazi, V. (1988). An introduction to rod-shaped lactic acid bacteria. *Biochimie*, **70**, 303-315.

Bouton, Y., Guyot, P., Beuquier, E., Tailliez, P. and Grappin, R. (2002). Use of PCR-based methods and PFGE for typing and monitoring homofermentative lactobacilli during Comte cheese ripening. *International Journal of Food Microbiology*, **76**, 27-38.

Brown, J.R. (2003). Ancient horizontal gene transfer. *Nature Reviews Genetics*, **4**, 121-132.

Bvochora, J.M., Reed, J.D., Read, J.S. and Zvauya, R. (1999). Effect of fermentation processes on proanthocyanidins in sorghum during preparation of *mahewu*, a non-alcoholic beverage. *Process Biochemistry*, **35**, 21-25.

Bvochora, J.M., Danner, H., Miyafuji, H., Braun, R. and Zvauya, R. (2005). Variation of sorghum phenolic compounds during the preparation of opaque beer. *Process Biochemistry*, **40**, 1207-1213.

Campieri, C., Campieri, M., Bertuzzi, V., Swennen, E., Matteuzzi, D., Stefoni, S., Pirovano, F., Centi, C., Ulisse, S., Famularo, G. and De Simone, C. (2001). Reduction of oxaluria after an oral course of lactic acid bacteria at high concentration. *Kidney International*, **60**, 1097-1105.

Caplice, E. and Fitzgerald, G.F. (1999). Food fermentations: Role of microorganisms in food production and preservation. *International Journal of Food Microbiology*, **50**, 131-149.

Chambel, L., Chelo, I.M., Zé-zé, L., Pedro, L.G., Santos, M.A. and Tenreiro, R. (2006). *Leuconostoc pseudoficulneum* sp. nov., isolated from a ripe fig. *International Journal of Systematic and Evolutionary Microbiology*, **56**, 1375-1381.

Chatel, J.M., Langella, P., Adel-Patient, K., Commissaire, J., Wal, J.M. and Corthier, G. (2001). Induction of mucosal immune response after intranasal or oral inoculation of mice with *Lactococcus lactis* producing bovine beta-lactoglobulin. *Clinical and Diagnostic Laboratory Immunology*, **8**, 545-551.

Chavan, J.K. and Kadam, S.S. (1989). Nutritional improvement of cereals by fermentation. *Critical Reviews in Food Science and Nutrition*, **28**, 349-400.

Chavan, U.D., Chavan, J.K. and Kadam, S.S. (1988). Effect of fermentation on soluble proteins and *in vitro* digestibility of sorghum, green gram and sorghum-green gram blends. *Journal of Food Science*, **53**, 1574-1575.

Cherrington, C.A., Hinton, M., Mead, G.C. and Chopra, I. (1991). Organic acids: Chemistry, antibacterial activity and practical applications. *Advances in Microbiology and Physics*, **32**, 87-108.

Choi, H.J., Cheigh, C.I., Kim, S.B., Lee, D.W., Choi, S.W., Park, J.M. and Pyun, Y.R. (2002). *Weissella kimchii* sp. nov., a novel lactic acid bacterium from *kimchi*. *International Journal of Systematic and Evolutionary Microbiology*, **52**, 507-511.

Choi, I.G. and Kim, S.H. (2007). Global extent of horizontal gene transfer. *Proceedings of the National Academy of Sciences of the United States of America*, **104**, 4489-4494.

Chung, K.T., Wong, T.Y., Wei, C.I., Huang, Y.W. and Lin, Y. (1998). Tannins and human health: A review. *Critical Reviews in Food Science and Nutrition*, **38**, 421-464.

Cocolin, L., Manzano, M., Cantoni, C. and Comi, G. (2001). Denaturing gradient gel electrophoresis analysis of the 16S rRNA gene V1 region to monitor dynamic changes in the bacterial population during fermentation of Italian sausages. *Applied and Environmental Microbiology*, **67**, 5113-5121.

Collins, M.D., Samelis, J., Metaxopoulos, J. and Wallbanks, S. (1993). Taxonomic studies on some leuconostoc-like organisms from fermented sausages: Description of a new genus *Weissella* for the *Leuconostoc paramesenteroides* group of species. *Journal of Applied Bacteriology*, **75**, 595-603.

Colwell, R.R. (1970). Polyphasic taxonomy of the genus *Vibrio*: Numerical taxonomy of *Vibrio cholerae*, *Vibrio parahaemolyticus*, and related *Vibrio* species. *Journal of Bacteriology*, **104**, 410-433.

Condon, S. (1987). Responses of lactic acid bacteria to oxygen. *FEMS Microbiology Reviews*, **46**, 269-280.

Corsetti, A., Lavermicocca, P., Morea, M., Baruzzi, F., Tosti, N. and Gobbetti, M. (2001). Phenotypic and molecular identification and clustering of lactic acid bacteria and yeasts from wheat (species *Triticum durum* and *Triticum aestivum*) sourdoughs of Southern Italy. *International Journal of Food Microbiology*, **65**, 95-104.

Daeschel, M.A., Andersson, R.E. and Fleming, H.P. (1987). Microbial ecology of fermenting plant materials. *FEMS Microbiology Reviews*, **46**, 357-367.

Dahllöf, I., Baillie, H. and Kjelleberg, S. (2000). *rpoB*-based microbial community analysis avoids limitations inherent in 16S rRNA gene intraspecies heterogeneity. *Applied and Environmental Microbiology*, **66**, 3376-3380.

Dalezios, I. and Siebert, K.J. (2001). Comparison of pattern recognition techniques for the identification of lactic acid bacteria. *Journal of Applied Microbiology*, **91**, 225-236.

Daubin, V., Moran, N.A. and Ochman, H. (2003). Phylogenetics and the cohesion of bacterial genomes. *Science*, **301**, 829-832.

Davis, C.R., Wibowo, D., Eschenbruch, R., Lee, T.H. and Fleet, G.H. (1985). Practical implications of malolactic fermentation: A review. *American Journal of Enology and Viticulture*, **36**, 290-301.

Devriese, L.A. and Pot, B. (1995). The genus *Enterococcus*. In: The genera of lactic acid bacteria. Wood, B.J.B and Holzapfel, W.H. (Eds.). Blackie Academic, London, UK, pp. 327-367.

De Vuyst, L. (2000). Technology aspects related to the application of functional starter cultures. *Food Technology and Biotechnology*, **38**, 105-112.

De Vuyst, L. and Vandamme, E.J. (1994). Antimicrobial potential of lactic acid bacteria. In: Bacteriocins of lactic acid bacteria. De Vuyst, L. and Vandamme, E.J. (Eds.). Blackie Academic, London, UK, pp. 91-142.

Dendy, D.A.V. (1995). Sorghum and millets: Production and importance. In: Sorghum and millets: Chemistry and technology. Dendy, D.A.V. (Ed.). American Association of Cereal Chemists, St. Paul, USA, pp. 11-26.

- Dicko, M.H., Gruppen, H., Traoré, A.S., Voragen, A.G.J. and van Berkel, W.J.H.** (2006). Sorghum grain as human food in Africa: Relevance of content of starch and amylase activities. *African Journal of Biotechnology*, **5**, 384-395.
- Dirar, H.A.** (1993). The indigenous fermented foods of the Sudan. CAB International.
- Doggett, H.** (1970). Sorghum. John Wiley and Sons, New York, USA.
- Doggett, H.** (1988). Sorghum. John Wiley and Sons, New York, USA.
- Doolittle, W.F.** (1999). Phylogenetic classification and the universal tree. *Science*, **284**, 2124-2129.
- Drouault, S., Juste, C., Marteau, P., Renault, P. and Corthier, G.** (2002). Oral treatment with *Lactococcus lactis* expressing *Staphylococcus hyicus* lipase enhances lipid digestion in pigs with induced pancreatic insufficiency. *Applied and Environmental Microbiology*, **68**, 3166-3168.
- Drysdale, G.S. and Fleet, G.H.** (1989). The effect of acetic acid bacteria upon the growth and metabolism of yeast during the fermentation of grape juice. *Journal of Applied Bacteriology*, **67**, 471-481.
- Dunbar, J., Ticknor, L.O. and Kuske, C.R.** (2001). Phylogenetic specificity and reproducibility and new methods for analysis of terminal-restriction fragment profiles of 16S rRNA genes from bacterial communities. *Applied and Environmental Microbiology*, **67**, 190-197.
- Duthoit, F., Godon, J.J. and Montel, M.C.** (2003). Bacterial community dynamics during production of registered designation of origin Salers cheese as evaluated by 16S rRNA gene single-strand conformation polymorphism analysis. *Applied and Environmental Microbiology*, **69**, 3840-3848.
- Dvorakova, J.** (1998). Phytase: Sources, preparation and exploitation. *Folia Microbiology*, **43**, 323-338.
- Eggum, B.O., Monowar, L., Boch Knusden, K.E., Munck, L. and Axtell, J.** (1983). Nutritional quality of sorghum foods from Sudan. *Journal of Cereal Science*, **1**, 127-137.
- Ehrmann, M.A. and Vogel, R.F.** (2005). Molecular taxonomy and genetics of sourdough lactic acid bacteria. *Trends in Food Science and Technology*, **16**, 31-42.
- El Tinay, A.H., Abdel Gadir, A.M. and El Hidai, M.** (1979). Sorghum fermented *kisra* bread. Nutritive value of *kisra*. *Journal of the Science of Food and Agriculture*, **30**, 859-863.

Ercolini, D. (2004). PCR-DGGE fingerprinting: Novel strategies for detection of microbes in food. *Journal of Microbiological Methods*, **56**, 297-314.

Ercolini, D., Mauriello, G., Blaiotta, G., Moschetti, G. and Coppola, S. (2004). PCR-DGGE fingerprints of microbial succession during a manufacture of traditional water buffalo mozzarella cheese. *Journal of Applied Microbiology*, **96**, 263-270.

Erlandson, K. and Batt, C.A. (1997). Strain-specific differentiation of lactococci in mixed starter culture populations using randomly amplified polymorphic DNA-derived probes. *Applied and Environmental Microbiology*, **63**, 2702-2707.

Escalante, A., Wachter, C. and Farres, A. (2001). Lactic acid bacteria diversity in the traditional Mexican fermented dough *pozol* as determined by 16S rDNA sequence analysis. *International Journal of Food Microbiology*, **64**, 21-31.

Farrelly, V., Rainey, F.A. and Stackebrandt, E. (1995). Effect of genome size and *rrn* copy number on PCR amplification of 16S rRNA genes from a mixture of bacterial species. *Applied and Environmental Microbiology*, **61**, 2798-2801.

Fasoli, S., Marzotto, M., Rizzotti, L., Rossi, F., Dellaglio, F. and Torriani, S. (2003). Bacterial composition of commercial probiotic products as evaluated by PCR-DGGE analyses. *International Journal of Food Microbiology*, **82**, 59-70.

Felis, G.E. and Dellaglio, F. (2007). Taxonomy of *Lactobacilli* and *Bifidobacteria*. *Current Issues in Intestinal Microbiology*, **8**, 44-61.

Fischer, S.G. and Lerman, L.S. (1983). DNA fragments differing by single base pair substitutions are separated in denaturing gradient gels: Correspondence with melting theory. *Proceedings of the National Academy of Sciences of the United States of America*, **80**, 1579-1583.

Flaherty, J.D., Levett, P.N., Dewhirst, F.E., Troe, T.E., Warren, J.R. and Johnson, S. (2003). Fatal case of endocarditis due to *Weissella confusa*. *Journal of Clinical Microbiology*, **41**, 2237-2239.

Fogel, G.B., Collins, C.R., Li, J. and Brunk, C.F. (1999). Prokaryotic genome size and SSU rDNA copy number: Estimation of microbial relative abundance from a mixed population. *Microbial Ecology*, **38**, 93-113.

Fontaine, E.A., Claydon, E. and Taylor-Robinson, D. (1996). *Lactobacilli* from women with or without bacterial vaginosis and observations on the significance of hydrogen peroxide. *Microbial Ecology in Health and Disease*, **9**, 135-141.

- Food and Agriculture Organization of the United Nations.** (1998). Carbohydrates in human nutrition. (<http://www.fao.org/docrep/W8079E/w8079e0h.htm>).
- Food and Agriculture Organization of the United Nations.** (2003). The state of food insecurity in the world. (www.fao.org).
- Food and Agriculture Organization of the United Nations.** (2006). FAOSTAT – FAO Statistical Databases. (<http://faostat.fao.org/>).
- Fooks, L.J., Fuller, R. and Gibson, G.R.** (1999). Prebiotics, probiotics and human gut microbiology. *International Dairy Journal*, **9**, 53-61.
- Fortina, M.G., Ricci, G., Mora, D. and Manachini, P.L.** (2004). Molecular analysis of artisanal Italian cheeses reveals *Enterococcus italicus* sp. nov. *International Journal of Systematic and Evolutionary Microbiology*, **54**, 1717-1721.
- Friedrich, M.W.** (2002). Phylogenetic analysis reveals multiple lateral transfers of adenosine-5'-phosphosulfate reductase genes among sulfate-reducing microorganisms. *Journal of Bacteriology*, **184**, 278-289.
- Fuller, R.** (1989). Probiotics in man and animals. *Journal of Applied Bacteriology*, **66**, 365-378.
- Gänzle, M.G.** (2004). Reutericyclin: Biological activity, mode of action, and potential applications. *Applied Microbiology and Biotechnology*, **64**, 326-332.
- Gänzle, M.G. and Vogel, R.F.** (2003). Studies on the mode of action of reutericyclin. *Applied and Environmental Microbiology*, **69**, 1305-1307.
- Garrity, G.M., Winters, M. and Searles, D.B.** (2001). Taxonomic outline of the prokaryotic genera. *Bergey's Manual of Systematic Bacteriology*.
- Garrity, G.M., Bell, J.A. and Lilburn, T.** (2004). Taxonomic outline of the prokaryotes. *Bergey's Manual of Systematic Bacteriology*.
- Gashe, B.A., Girma, M. and Bisrat, A.** (1982). *Tef* fermentation. The role of microorganisms in fermentation and their effect on the nitrogen content of *tef*. *Ethiopian Journal of Science*, **5**, 69-76.
- Ge, F., Wang, L.S. and Kim, J.** (2005). The cobweb of life revealed by genome-scale estimates of horizontal gene transfer. *PLoS Biol*, **3**, e316.

- Gebrekidan, B. and Gebrettiwat, B.** (1982). Sorghum *injera*: Preparation and quality parameters. In: Proceedings of the International Symposium on Sorghum Grain Quality. International Crops Research Institute for the Semi-Arid Tropics. Mertin, J.V. (Ed.). Patancheru, India, pp. 55-56.
- Gevers, D., Huys, G. and Swings, J.** (2001). Applicability of rep-PCR fingerprinting for identification of *Lactobacillus* species. FEMS Microbiology Letters, **205**, 31-36.
- Giacomazzi, S., Leroi, F., L'Henaff, C. and Joffraud, J.J.** (2004). *rpoB*-PCR amplified gene and temporal temperature gradient gel electrophoresis: A rapid tool to analyse bacterial strains representative of cold-smoked salmon microflora. Letters in Applied Microbiology, **38**, 130-134.
- Gilliland, S.E. and Kim, H.S.** (1984). Effect of viable starter culture bacteria in yogurt on lactose utilization in humans. Journal of Dairy Science, **67**, 1-6.
- Gilliland, S.E., Nelson, C.R. and Maxwell, C.** (1985). Assimilation of cholesterol by *Lactobacillus acidophilus*. Applied and Environmental Microbiology, **49**, 377-381.
- Giraffa, G. and Neviani, E.** (2000). Molecular identification and characterization of food-associated lactobacilli. Italian Journal of Food Science, **4**, 403-423.
- Giraffa, G. and Neviani, E.** (2001). DNA-based, culture-independent strategies for evaluating microbial communities in food-associated ecosystems. International Journal of Food Microbiology, **67**, 19-34.
- Giraffa, G., Lazzi, C., Gatti, M., Rossetti, L., Mora, D. and Neviani, E.** (2002). Molecular typing of *Lactobacillus delbrueckii* of dairy origin by PCR-RFLP of protein-coding genes. International Journal of Food Microbiology, **82**, 163-172.
- Gobbetti, M. and Corsetti, A.** (1997). *Lactobacillus sanfrancisco*, a key sourdough lactic acid bacterium: A review. Food Microbiology, **14**, 175-188.
- Goldin, B.R.** (1990). Intestinal microflora: Metabolism of drugs and carcinogens. Annals of Medicine, **22**, 43-48.
- Gonzalez, C.J., Encinas, J.P., Garcia-Lopez, M.L. and Otero, A.** (2000). Characterisation and identification of lactic acid bacteria from fresh water fishes. Food Microbiology, **17**, 383-391.
- Goodfellow, M., Manfio, G.P. and Chun, J.** (1997). Towards a practical species concept for cultivable bacteria. In: Species: The units of biodiversity. Clarridge, M.F. and Dawah, H.A. (Eds.). Chapman and Hall, London, UK, pp. 25-29.

Guarneri, T., Rossetti, L. and Giraffa, G. (2001). Rapid identification of *Lactobacillus brevis* using polymerase chain reaction. *Letters in Applied Microbiology*, **33**, 377-381.

Hagerman, A.E. (1989). Chemistry of tannin-protein complexation. In: Chemistry and significance of condensed tannin. Hemingway, R.W. and Karchesy, J.J. (Eds.). Plenum Press, New York, USA, pp. 323-333.

Hamacker, B.R., Kirleis, A.W., Mertz, E.T. and Axtell, J.D. (1986). Effect of cooking on the protein profiles and *in vitro* protein digestibility of sorghum and maize. *Journal of Agricultural and Food Chemistry*, **34**, 647-649.

Hammes, W.P. and Vogel, R.F. (1995). The genus *Lactobacillus*. In: The genera of lactic acid bacteria. Wood, B.J.B. and Holzapfel, W.H. (Eds.). Blackie Academic, London, UK, pp. 19-54.

Hansen, E.B. (2002). Commercial bacterial starter cultures for fermented foods of the future. *International Journal of Food Microbiology*, **78**, 119-131.

Hassan, I.A.G. and El Tinay, A.H. (1995). Effect of fermentation on tannin content and *in vitro* protein and starch digestibilities of two sorghum cultivars. *Food Chemistry*, **53**, 149-151.

Hayashi, K. (1991). PCR-SSCP: A simple and sensitive method for detection of mutations in a genomic DNA. *PCR Methods and Applications*, **1**, 34-38.

Hayford, A.E., Petersen, A., Vogensen, F.K. and Jakobsen, M. (1999). Use of conserved randomly amplified polymorphic DNA (RAPD) fragments and RAPD pattern for characterization of *Lactobacillus fermentum* in Ghanaian fermented maize dough. *Applied and Environmental Microbiology*, **65**, 3213-3221.

Henick-Kling, T. (1993). Malolactic fermentation. In: Wine microbiology and biotechnology. Fleet, G.H. (Ed.). Harwood Academic Publishers, Zurich, Switzerland, pp. 286-326.

Holley, R.A., Guan, T.Y., Peirson, M. and Yost, C.K. (2002). *Carnobacterium viridans* sp. nov., an alkaliphilic, facultative anaerobe isolated from refrigerated, vacuum-packed bologna sausage. *International Journal of Systematic and Evolutionary Microbiology*, **52**, 1881-1885.

Hols, P., Kleerebezem, M., Schanck, A.N., Ferain, T., Hugenholtz, J., Delcour, J. and de Vos, W.M. (1999). Conversion of *Lactococcus lactis* from homolactic to homoalanine fermentation through metabolic engineering. *Nature Biotechnology*, **17**, 588-592.

- Holzappel, W.H.** (1997). Use of starter cultures in fermentation on a household scale. *Food Control*, **8**, 241-258.
- Holzappel, W.H.** (2002). Appropriate starter culture technologies for small-scale fermentation in developing countries. *International Journal of Food Microbiology*, **75**, 197-212.
- Huang, Y.C., Edwards, C.G., Peterson, J.C. and Haag, K.M.** (1996). Relationship between sluggish fermentations and antagonism of yeast by lactic acid bacteria. *American Journal of Enology and Viticulture*, **47**, 1-9.
- Hughenoltz, J.** (1993). Citrate metabolism in lactic acid bacteria. *FEMS Microbiology Reviews*, **12**, 165-178.
- Hulton, C.S.J., Higgins, C.F. and Sharp, P.M.** (1991). ERIC sequences: A novel family of repetitive elements in the genomes of *Escherichia coli*, *Salmonella typhimurium*, and other enterobacteria. *Molecular Microbiology*, **5**, 825-834.
- Ibrahim, F.S., Babiker, E.E., Yousif, N.E. and El Tinay, A.H.** (2005). Effect of fermentation on biochemical and sensory characteristics of sorghum flour supplemented with whey protein. *Food Chemistry*, **92**, 285-292.
- ICRISAT.** (2005). Sorghum report. Patancheru, India.
(<http://www.icrisat.org/text/research/grep/homepage/sorghum/sorghumhomepage.htm>).
- Isenberg, H.D., Vellozzi, E.M., Shapiro, J. and Rubin, L.G.** (1988). Clinical laboratory challenges in the recognition of *Leuconostoc* spp. *Journal of Clinical Microbiology*, **26**, 479-483.
- Jay, J.M.** (1982). Antimicrobial properties of diacetyl. *Applied and Environmental Microbiology*, **44**, 525-532.
- Joosten, H.M.L.J. and Nuñez, M.** (1996). Prevention of histamine formation in cheese by bacteriocin-producing lactic acid bacteria. *Applied and Environmental Microbiology*, **62**, 1178-1181.
- Jordan, K., Makarova, K.S., Spouge, J.L., Wolf, Y.I. and Koonin, E.V.** (2001). Lineage-specific gene expansions in bacterial and archaeal genomes. *Genome Research*, **11**, 555-565.
- Jordan, W.R. and Sullivan, C.Y.** (1982). Reaction and resistance of grain sorghum to heat and drought. In: *Sorghum in the eighties*. Mertin, J.V. (Ed.). International Crops Research Institute for the Semi-Arid Tropics. Patancheru, India, pp. 131-142.

- Kandler, O. and Weiss, N.** (1986). Regular, nonsporing Gram-positive rods. In: Bergey's Manual of Systematic Bacteriology. Sneath *et al.* (Eds.). Williams and Wilkins, Baltimore, USA, pp. 1208-1234.
- Kazanas, N. and Fields, M.L.** (1981). Nutritional improvement of sorghum by fermentation. Journal of Food Science, **46**, 819-821.
- Kempler, G.M. and McKay, L.L.** (1981). Biochemistry and genetics of citrate utilization in *Streptococcus lactis* subsp. *diacetylactis*. Journal of Dairy Science, **64**, 1527-1539.
- Kim, B., Lee, J., Jang, J. and Han, H.** (2003). *Leuconostoc inhae* sp. nov., a lactic acid bacterium isolated from *kimchi*. International Journal of Systematic and Evolutionary Microbiology, **53**, 1123-1126.
- Kimber, C.T.** (2000). Origins of domesticated sorghum and its early diffusion into India and China. In: Sorghum: Origin, history, technology, and production. Smith, C.W. and Frederiksen, R.A. (Eds.). John Wiley and Sons, New York, USA, pp. 3-98.
- King, A.D.G. and Nagel, C.W.** (1975). Influence of carbon dioxide upon the metabolism of *Pseudomonas aeruginosa*. Journal of Food Science, **40**, 362-366.
- Kontiokari, T., Sundqvist, K. and Nuutinen, M.** (2001). Randomised trial of cranberry lingonberry juice and *Lactobacillus* GG drink for the prevention of urinary tract infections in women. British Medical Journal, **322**, 1571-1576.
- Koonin, E.V.** (2003). Horizontal gene transfer: The path to maturity. Molecular Microbiology, **50**, 725-727.
- Koonin, E.V., Makarova, K.S. and Aravind, L.** (2001). Horizontal gene transfer in prokaryotes: Quantification and classification. Annual Reviews in Microbiology, **55**, 709-742.
- Koort, J., Coenye, T., Vandamme, P., Sukura, A. and Björkroth, J.** (2004). *Enterococcus hermanniensis* sp. nov., from modified atmosphere-packaged broiler and canine tonsils. International Journal of Systematic and Evolutionary Microbiology, **54**, 1823-1827.
- Kunene, N.F., Geornaras, I., von Holy, A. and Hastings, J.W.** (2000). Characterization and determination of origin of lactic acid bacteria from a sorghum-based fermented weaning food by analysis of soluble proteins and amplified fragment length polymorphism fingerprinting. Applied and Environmental Microbiology, **66**, 1084-1092.

- Kurland, C.G.** (2000). Something for everyone. Horizontal gene transfer in evolution. *EMBO Reports*, **1**, 92-95.
- Kurland, C.G., Canback, B. and Berg, O.G.** (2003). Horizontal gene transfer: A critical view. *Proceedings of the National Academy of Sciences of the United States of America*, **100**, 9658-9662.
- Lavermicocca, P., Valeria, F., Evidente, A., Lazzaroni, S., Corsetti, A. and Gobbetti, M.** (2000). Purification and characterization of novel antifungal compounds by sourdough *Lactobacillus plantarum* 21B. *Applied and Environmental Microbiology*, **66**, 4084-4090.
- Lawrence, J.G. and Hendrickson, H.** (2003). Lateral gene transfer: When will adolescence end? *Molecular Microbiology*, **50**, 739-749.
- Lee, D.H., Zo, Y.G. and Kim, S.J.** (1996). Non-radioactive method to study genetic profiles of natural bacterial communities by PCR-SSCP. *Applied and Environmental Microbiology*, **62**, 3112-3120.
- Leisner, J.J., Vancanneyt, M., Goris, J., Christensen, H. and Rusul, G.** (2000). Description of *Paralactobacillus selangolensis* gen. nov., sp. nov., a new lactic acid bacterium isolated from *chili bo*, a Malaysian food ingredient. *International Journal of Systematic and Evolutionary Microbiology*, **50**, 19-24.
- Leroy, F. and De Vuyst, L.** (2004). Lactic acid bacteria as functional starter cultures for the food fermentation industry. *Trends in Food Science and Technology*, **15**, 67-78.
- Lindgren, S.E. and Dobrogosz, W.J.** (1990). Antagonistic activities of lactic acid bacteria in food and feed fermentations. *FEMS Microbiology Letters*, **87**, 149-163.
- Liu, W.T., Marsh, T.L., Cheng, H. and Forney, L.J.** (1997). Characterization of microbial diversity by determining terminal-restriction fragment length polymorphisms of genes encoding 16S rRNA. *Applied and Environmental Microbiology*, **63**, 4516-4522.
- Longo, M.A. and Sanromán, M.A.** (2006). Production of food aroma compounds: Microbial and enzymatic methodologies. *Food Technology and Biotechnology*, **44**, 335-353.
- Lorri, W.** (1993). Nutritional and microbiological evaluation of fermented cereal weaning foods. Ph.D thesis, Chambers University of Technology, Gotenborg, Sweden.
- Lorri, W. and Svanberg, U.** (1995). An overview of the use of fermented foods for child feeding in Tanzania. *Ecology of Food and Nutrition*, **34**, 65-81.

- Lupski, J.R. and Weinstock, G.M.** (1992). Short, interspersed repetitive DNA sequences in prokaryotic genomes. *Journal of Bacteriology*, **174**, 4525-4529.
- Lyhs, U., Korkeala, H. and Björkroth, J.** (2002). Characterization of lactic acid bacteria from spoiled, vacuum-packaged 'gravad' rainbow trout using ribotyping. *International Journal of Food Microbiology*, **72**, 147-153.
- Makarova, K.S. and Koonin, E.V.** (2007). Evolutionary genomics of lactic acid bacteria. *Journal of Bacteriology*, **189**, 1199-1208.
- Marfo, E.K., Simpson, B.K., Idowu, J.S. and Oke, O.L.** (1990). Effect of local food processing on phytate levels in cassava, cocoyam, yam, maize, sorghum, rice, cowpea and soybean. *Journal of Agricultural and Food Chemistry*, **38**, 1580-1585.
- Marsh, T.L., Saxman, P., Cole, J. and Tiedje, J.M.** (2000). Terminal-restriction fragment length polymorphism analysis program, a web-based research tool for microbial community analysis. *Applied and Environmental Microbiology*, **66**, 3616-3620.
- Martin, B., Humbert, O., Camara, M., Guenzi, E., Walker, J., Mitchell, T., Andrew, P., Prudhomme, M., Alloing, G. and Hakenbeck, R.** (1992). A highly conserved repeated DNA element located in the chromosome of *Streptococcus pneumoniae*. *Nucleic Acids Research*, **20**, 3479-3483.
- Martineau, B. and Henick-Kling, T.** (1995). Formation and degradation of diacetyl in wine during alcoholic fermentation with *Saccharomyces cerevisiae* strain EC1118 and malolactic fermentation with *Leuconostoc oenos* strain MCW. *American Journal of Enology and Viticulture*, **46**, 442-448.
- Massol-Deya, A.A., Odelson, D.A., Hickey, R.F. and Tiedje, J.M.** (1995). Bacterial community fingerprinting of amplified 16S and 16S-23S ribosomal DNA genes sequences and restriction endonuclease analysis (ARDRA). *Molecular Microbial Ecology Manual*, **3**, 1-18.
- Masters, C.I., Shallcross, J.A. and Mackey, B.M.** (1994). Effect of stress treatments on the detection of *Listeria monocytogenes* and enterotoxigenic *Escherichia coli* by the polymerase chain reaction. *Journal of Applied Bacteriology*, **77**, 73-79.
- Mbugua, S.K.** (1984). Isolation and characterization of lactic acid bacteria during the traditional fermentation of *uji*. *East African Agriculture and Forestry Journal*, **50**, 36-43.
- McCartney, A.L.** (2002). Application of molecular biological methods for studying probiotics and the gut flora. *British Journal of Nutrition*, **88**, S29-S37.

- McGowan, C., Fulthorpe, R., Wright, A. and Tiedje, J.M.** (1998). Evidence for interspecies gene transfer in the evolution of 2,4-dichlorophenoxyacetic acid degraders. *Applied and Environmental Microbiology*, **64**, 4089-4092.
- Meroth, C.B., Walter, J., Hertel, C., Brandt, M.J. and Hammes, W.P.** (2003). Monitoring the bacterial population dynamics in sourdough fermentation processes by using PCR-denaturing gradient gel electrophoresis. *Applied and Environmental Microbiology*, **69**, 475-482.
- Mertz, E.T., Hassen, M.M., Cairns-Whittern, C., Kirleis, A.W., Tu, L. and Axtell, J.D.** (1984). Pepsin digestibility of sorghum and other major cereals. *Proceedings of the National Academy of Sciences of the United States of America*, **81**, 1-2.
- Messens, W. and De Vuyst, L.** (2002). Inhibitory substances produced by lactobacilli isolated from sourdoughs-A review. *International Journal of Food Microbiology*, **72**, 31-43.
- Milliere, J.B., Mathot, A.G., Schmitt, P. and Divies, C.** (1989). Phenotypic characterisation of *Leuconostoc* species. *Journal of Applied Bacteriology*, **67**, 529-542.
- Mital, B.K., Schallenberger, R.S. and Steinkraus, K.H.** (1973). Alpha-galactosidase activity of lactobacilli. *Applied Microbiology*, **26**, 783-788.
- Moeseneder, M.M., Arrieta, J.M., Muyzer, G., Winter, C. and Herndl, G.J.** (1999). Optimisation of terminal-restriction fragment polymorphism analysis for complex marine bacterioplankton communities and comparison with denaturing gradient gel electrophoresis. *Applied and Environmental Microbiology*, **65**, 3518-3525.
- Mohammed, S.I., Steenson, L.R. and Kirleis, A.W.** (1991). Isolation and characterization of microorganisms associated with the traditional sorghum fermentation for production of Sudanese *kisra*. *Applied and Environmental Microbiology*, **57**, 2529-2533.
- Monchois, V., Willemot, R.M. and Monsan, P.** (1999). Glucansucrases: Mechanism of action and structure-function relationships. *FEMS Microbiology Reviews*, **23**, 131-151.
- Mota-Meira, M., Lapointe, G., Lacroix, C. and Lavoie, M.C.** (2000). MICs of mutacin B-Ny266, nisin A, vancomycin, and oxacillin against bacterial pathogens. *Antimicrobial Agents Chemotherapy*, **44**, 24-29.
- Mugula, J.K., Nnko, S.A.M. and Sørhaug, T.** (2001). Changes in quality attributes during storage of *togwa*, a lactic acid fermented gruel. *Journal of Food Safety*, **21**, 181-194.

Mugula, J.K., Nnko, S.A.M., Narvhus, J.A. and Sørhaug, T. (2003). Microbiological and fermentation characteristics of *togwa*, a Tanzanian fermented food. *International Journal of Food Microbiology*, **80**, 187-199.

Mukuru, S. (1992). Traditional technologies in small grain processing. In: Utilisation of sorghum and millets. Gomez *et al.* (Eds.). International Crops Research Institute for Semi-Arid Tropics, Patancheru, India, pp. 47-56.

Muyanja, C.M.B.K., Narvhus, J.A., Treimo, J. and Langsrud, T. (2003). Isolation, characterization and identification of lactic acid bacteria from *bushera*: A Ugandan traditional fermented beverage. *International Journal of Food Microbiology*, **80**, 201-210.

Muyzer, G. (1999). DGGE/TGGE: A method for identifying genes from natural ecosystems. *Current Opinion in Microbiology*, **2**, 317-322.

Muyzer, G. and Smalla, K. (1998). Application of denaturing gradient gel electrophoresis (DGGE) and temperature gradient gel electrophoresis (TGGE) in microbial ecology. *Antonie Van Leeuwenhoek*, **73**, 127-141.

Muyzer, G., de Waal, E.C. and Uitterlinden, A.G. (1993). Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction amplified genes coding for 16S rRNA. *Applied and Environmental Microbiology*, **59**, 695-700.

National Research Council. (1996). *Lost crops of Africa, 1, Grains*. National Academy Press, Washington DC, USA.

Neucere, N.J. and Sumrell, G. (1979). Protein fractions from varieties of grain sorghum: Amino acid composition and solubility properties. *Journal of Agricultural and Food Chemistry*, **27**, 809-812.

Nga, B.H. (2005). Genome analysis of lactic acid bacteria in food fermentations and biotechnological applications. *Current Opinion in Microbiology*, **8**, 1-6.

Niemi, R.M., Heiskanen, I., Wallenius, K. and Lindström, K. (2001). Extraction and purification of DNA in rhizosphere samples for PCR-DGGE analysis of bacterial consortia. *Journal of Microbiological Methods*, **45**, 155-165.

Nigatu, A. (2000). Evaluation of numerical analyses of RAPD and API 50 CH patterns to differentiate *Lactobacillus plantarum*, *Lact. fermentum*, *Lact. rhamnosus*, *Lact. sake*, *Lact. parabuchneri*, *Lact. gallinarum*, *Lact. casei*, *Weissella minor* and related taxa isolated from *kocho* and *tef*. *Journal of Applied Microbiology*, **89**, 969-978.

- Nomura, M., Kobayashi, M. and Okamoto, T.** (2002). Rapid PCR-based method which can determine both phenotype and genotype of *Lactococcus lactis* subspecies. *Applied and Environmental Microbiology*, **68**, 2209-2213.
- Nout, M.J.R. and Motarjemi, Y.** (1997). Assessment of fermentation as a household technology for improving food safety: A joint FAO/WHO workshop. *Food Control*, **8**, 221-226.
- Ochman, H.** (2005). Genomes on the shrink. *Proceedings of the National Academy of the Sciences of the United States of America*, **102**, 11959-11960.
- Olive, D.M. and Bean, P.** (1999). Principles and applications of methods for DNA-based typing of microbial organisms. *Journal of Clinical Microbiology*, **37**, 1661-1669.
- Orita, M., Iwahana, H., Knazawa, H., Hayashi, K. and Sekiya, T.** (1989). Detection of polymorphisms of human DNA by gel electrophoresis as single strand conformation polymorphisms. *Proceedings of the National Academy of Sciences of the United States of America*, **86**, 2766-2770.
- Osborn, A.M., Moore, R.B. and Timmis, K.N.** (2000). An evaluation of terminal-restriction fragment length polymorphism (T-RFLP) analysis for the study of microbial community structure and dynamics. *Environmental Microbiology*, **2**, 39-50.
- Osman, M.A.** (2004). Changes in sorghum enzyme inhibitors, phytic acid, tannins and *in vitro* protein digestibility occurring during *khamir* (local bread) fermentation. *Journal of Food Chemistry*, **88**, 129-134.
- Oyewole, O.B.** (1997). Lactic fermented foods in Africa and their benefits. *Food Control*, **8**, 289-297.
- Parker, M.L., Melaku, U. and Faulks, R.M.** (1989). The contribution of flour components to the structure of *injera*, an Ethiopian fermented bread made from *tef* (*Eragrostis tef*). *Journal of Cereal Science*, **10**, 93-104.
- Plengvidhya, V., Breidt, F. and Fleming, H.P.** (2004). Use of RAPD-PCR as a method to follow the progress of starter cultures in sauerkraut fermentation. *International Journal of Food Microbiology*, **93**, 287-296.
- Randazzo, C.L., Torriani, S., Akkermans, A.D.L., de Vos, W.M. and Vaughan, E.E.** (2002). Diversity, dynamics, and activity of bacterial communities during production of an artisanal Sicilian cheese as evaluated by 16S rRNA analysis. *Applied and Environmental Microbiology*, **68**, 1882-1892.

Rankine, B.C., Fornachon, J.C.M. and Bridson, D.A. (1969). Diacetyl in Australian dry red wines and its significance in wine quality. *Vitis*, **8**, 129-134.

Reguant, C. and Bordons, A. (2003). Typification of *Oenococcus oeni* strains by multiplex RAPD-PCR and study of population dynamics during malolactic fermentation. *Journal of Applied Microbiology*, **95**, 344-353.

Reid, G. (2001). Probiotic agents to protect the urogenital tract against infection. *American Journal of Clinical Nutrition*, **73**, 437S-443S.

Reysenbach, A.L., Giver, L.J., Wickham, G.S. and Pace, N.R. (1992). Differential amplification of rRNA genes by polymerase chain reaction. *Applied and Environmental Microbiology*, **58**, 3417-3418.

Riley, M.A. and Wertz, J.E. (2002). Bacteriocins: Evolution, ecology and application. *Annual Reviews in Microbiology*, **56**, 117-137.

Romano, P. and Suzzi, G. (1996). Origin and production of acetoin during wine yeast fermentation. *Applied and Environmental Microbiology*, **62**, 309-315.

Rosselló-Mora, R. and Amann, R. (2001). The species concept for prokaryotes. *FEMS Microbiology Reviews*, **25**, 39-67.

Roy, D., Sirois, S. and Vincent, D. (2001). Molecular discrimination of lactobacilli used as starter and probiotic cultures by amplified ribosomal DNA restriction analysis. *Current Microbiology*, **42**, 282-289.

Ruoff, K.L., Kuritzkes, D.R., Wolfson, J.S. and Ferraro, M.J. (1988). Vancomycin-resistant Gram-positive bacteria isolated from human sources. *Journal of Clinical Microbiology*, **26**, 2064-2068.

Salminen, M.K., Tynkkynen, S., Rautelin, H., Saxelin, M., Vaara, M., Ruutu, P., Sarna, S., Valtonen, V. and Järvinen, A. (2002). *Lactobacillus* bacteremia during a rapid increase in probiotic use of *Lactobacillus rhamnosus* GG in Finland. *Clinical and Infectious Diseases*, **35**, 1155-1160.

Sánchez, J.I., Rossetti, L., Martinez, B., Rodriguez, A. and Giraffa, G. (2006). Application of reverse transcriptase PCR-based T-RFLP to perform semi-quantitative analysis of metabolically active bacteria in dairy fermentations. *Journal of Microbiological Methods*, **65**, 268-277.

Satokari, R., Mattila-Sandholm, T. and Suihko, M.L. (2000). Identification of *Pediococci* by ribotyping. *Journal of Applied Microbiology*, **88**, 260-265.

- Savadago, A., Quattara, C.A.T., Bassole, I.H.N. and Traore, S.A.** (2006). Bacteriocins and lactic acid bacteria - A minireview. *African Journal of Biotechnology*, **5**, 678-683.
- Sbordone, L. and Bortolaia, C.** (2003). Oral microbial biofilms and plaque-related diseases: Microbial communities and their role in the shift from oral health to disease. *Clinical Oral Investigations*, **7**, 181-188.
- Schaafsma, G., Meulig, W.J.A., van Dokkum, W. and Bouley, C.** (1998). Effects of a milk product, fermented by *Lactobacillus acidophilus* and with fructo-oligosaccharides added, on blood lipids in male volunteers. *European Journal Clinical Nutrition*, **52**, 436-440.
- Sedewitz, B., Schleifer, K.H. and Gotz, F.** (1984). Purification and biochemical characterization of pyruvate oxidase from *Lactobacillus plantarum*. *Journal of Bacteriology*, **160**, 273-278.
- Sekiguchi, H., Tomioka, N., Nakahara, T. and Uchiyama, H.** (2001). A single band does not always represent single bacterial strains in denaturing gradient gel electrophoresis analysis. *Biotechnology Letters*, **23**, 1205-1208.
- Serna-Saldivar, S. and Rooney, L.W.** (1995). Structure and chemistry of sorghum and millets. In: *Sorghum and millets: Chemistry and technology*. Dendy, D.A.V. (Ed.). American Association of Cereal Chemists, St Paul, USA, pp. 69-124.
- Sharpe, M.E.** (1981). The genus *Lactobacillus*. In: *The Prokaryotes*. Starr *et al.* (Eds.). Springer-Verlag, Berlin, Germany, pp. 1653-1679.
- Shaw, B.G. and Harding, C.D.** (1989). *Leuconostoc gelidum* sp. nov. and *Leuconostoc carnosum* sp. nov. from chill-stored meats. *International Journal of Systematic Bacteriology*, **39**, 217-223.
- Shin, J.H., Kim, D.I., Kim, H.R., Dong Soo Kim, D.S., Kook, J.K. and Lee, J.N.** (2007). Severe infective endocarditis of native valves caused by *Weissella confusa* detected incidentally on echocardiography. *Journal of Infection*, **54**, e149-e151.
- Silva, M., Jacobus, N.V., Deneke, C. and Gorbach, S.L.** (1987). Antimicrobial substance from a human *Lactobacillus* strain. *Antimicrobial Agents and Chemotherapy*, **31**, 1231-1233.
- Simango, C.** (1997). Potential use of traditional fermented foods for weaning in Zimbabwe. *Social Science and Medicine*, **44**, 1065-1068.
- Smith, F.R. and Sherman, J.M.** (1939). The hemolytic streptococci of human feces. *Journal of Infectious Diseases*, **62**, 186-189.

- Smith, M.W., Feng, D.F. and Doolittle, R.F.** (1992). Evolution by acquisition: The case for horizontal gene transfers. *Trends in Biochemical Sciences*, **17**, 489-493.
- Snel, B., Huynen, M.A. and Dutilh, B.E.** (2005). Genome trees and the nature of genome evolution. *Annual Reviews in Microbiology*, **59**, 191-209.
- Soni, S.K. and Sandhu, D.K.** (1991). Role of yeast domination in Indian *idli* batter fermentation. *World Journal of Microbiology and Biotechnology*, **7**, 505-507.
- Soomro, A.H., Masud, T. and Anwaar, K.** (2002). Role of lactic acid bacteria (LAB) in food preservation and human health - A review. *Pakistan Journal of Nutrition*, **1**, 20-24.
- Southern, E.M.** (1975). Detection of specific sequences among DNA fragments separated by gel electrophoresis. *Journal of Molecular Biology*, **98**, 503-517.
- Stackebrandt, E. and Teuber, M.** (1988). Molecular taxonomy and phylogenetic position of lactic acid bacteria. *Biochimie*, **70**, 317-324.
- Stackebrandt, E. and Liesack, W.** (1993). Nucleic acids and classification. In: *Handbook of new bacterial systematics*. Goodfellow, M. and O'Donnell, A.G. (Eds.). Academic Press, London, UK, pp. 151-194.
- Staley, J.T. and Krieg, N.R.** (1989). Classification of prokaryotic organisms: An overview. In: *Bergey's Manual of Systematic Bacteriology*. Staley *et al.* (Eds.). Williams and Wilkins, Baltimore, USA, pp. 1601-1603.
- Steinkraus, K.H.** (1983). Fermented foods, feeds and beverages. *Biotechnology Advances*, **10**, 31-46.
- Steinkraus, K.H.** (1996). *Handbook of indigenous fermented foods*. Marcel Dekker, Inc., New York, USA.
- Steinkraus, K.H.** (1998). Bio-enrichment: Production of vitamins in fermented foods. In: *Microbiology of fermented foods*. Wood, J.B. (Ed.). Blackie Academic, London, UK, pp. 603-619.
- Stewart, R.B. and Getachew, A.** (1962). Investigations on the nature of *injera*. *Economic Botany*, **16**, 127-130.
- Stodolak, B., Starzyńska, A., Czystoń, M. and Żyła, K.** (2007). The effect of phytic acid on oxidative stability of raw and cooked meat. *Food Chemistry*, **101**, 1041-1045.

- Struve, J., Weiland, O. and Nord, C.E.** (1988). *Lactobacillus plantarum* endocarditis in a patient with benign monoclonal gammopathy. *Journal of Infection*, **17**, 127-130.
- Sussman, J.I., Baron, E.J., Goldberg, S.M., Kaplan, M.H. and Pizzarello, R.A.** (1986). Clinical manifestations and therapy of *Lactobacillus* endocarditis: Report of a case and review of the literature. *Reviews of Infectious Diseases*, **8**, 771-776.
- Sütas, Y., Soppi, E. and Korhonen, H.** (1996). Suppression of lymphocyte proliferation *in vitro* by bovine caseins hydrolysed with *Lactobacillus* GG-derived enzymes. *Journal of Allergy and Clinical Immunology*, **98**, 86-92.
- Taylor, J. and Taylor, J.R.N.** (2002). Alleviation of the adverse effects of cooking on protein digestibility in sorghum through fermentation in traditional African porridges. *International Journal of Food Science and Technology*, **37**, 129-138.
- Taylor, J.R.N.** (2003). Overview: Importance of sorghum in Africa. (<http://www.Afripro.org.uk/papers/Paper01Taylor.pdf> cited 20/05/03).
- Temmerman, R., Huys, G. and Swings, J.** (2004). Identification of lactic acid bacteria: Culture-dependent and culture-independent methods. *Trends in Food Science and Technology*, **15**, 348-359.
- Teniola, O.D., Odunfa, S.A. and Holzapfel, W.H.** (2005). Selection, use and the influence of starter cultures in the nutrition and processing improvement of *ogi*. In: *Proceedings of the Second International Workshop: Food-based approaches for a healthy nutrition*. Brouwer *et al.* (Eds.). Presses Universitaires de Ouagadougou, Burkina Faso, pp. 697-708.
- Tenover, F.C., Arbeit, R.D., Goering, R.V., Mickelsen, P.A., Murray, B.E., Persing, D.H. and Swaminathan, B.** (1995). Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: Criteria for bacterial strain typing. *Journal of Clinical Microbiology*, **33**, 2233-2239.
- Thomas, T.D., Ellwood, D.C. and Longyear, V.M.C.** (1979). Change from homo- to heterolactic fermentation by *Streptococcus lactis* resulting from glucose limitation in anaerobic chemostat cultures. *Journal of Bacteriology*, **138**, 109-117.
- Towo, E., Matuschek, E. and Svanberg, U.** (2006). Fermentation and enzyme treatment of tannin sorghum gruels: Effects of phenolic compounds, phytate and *in vitro* accessible iron. *Food Chemistry*, **94**, 369-376.

- Uyttendaele, M., van Boxstael, S. and Debevere, J.** (1999). PCR assay for detection of the *E. coli* 0157:H7 *eae*-gene and effect of the sample preparation method on PCR detection of heat-killed *E. coli* 0157:H7 in ground beef. *International Journal of Food Microbiology*, **52**, 85-95.
- Van Kraaij, C., De Vos, W., Siezen, R.J. and Kuipers, O.P.** (1999). Lantibiotics: Biosynthesis, mode of action and applications. *Natural Product Reports*, **16**, 575-587.
- Van Beek, S. and Priest, F.G.** (2001). Evolution of the lactic acid bacterial community during malt whisky fermentation: A polyphasic study. *Applied and Environmental Microbiology*, **68**, 297-305.
- Van Belkum, M.J., Kok, J., Venema, G., Holo, H., Nes, I.F., Konings, W.N. and Abee, T.** (1991). The bacteriocin lactococcin A specifically increases permeability of lactococcal cytoplasmic membranes in a voltage-independent, protein mediated manner. *Journal of Bacteriology*, **173**, 7934-7941.
- Van Reenen, C.A. and Dicks, L.M.T.** (1996). Evaluation of numerical analysis of Random Amplified Polymorphic DNA (RAPD)-PCR as a method to differentiate *Lactobacillus plantarum* and *Lactobacillus pentosus*. *Current Microbiology*, **32**, 183-187.
- Vandamme, P., Pot, B., Gillis, M., de Vos, P., Kersters, K. and Swings, J.** (1996). Polyphasic taxonomy, a consensus approach to bacterial systematics. *Microbiological Reviews*, **60**, 407-438.
- Vanechoutte, M.** (1996). DNA fingerprinting techniques for microorganisms. *Molecular Biotechnology*, **6**, 115-142.
- Ventura, M. and Zink, R.** (2002). Specific identification and molecular typing analysis of *Lactobacillus johnsonii* by using PCR-based methods and pulsed-field gel electrophoresis. *FEMS Microbiology Letters*, **217**, 141-154.
- Versalovic, J., Koeuth, T. and Lupsi, J.R.** (1991). Distribution of repetitive DNA sequences in eubacteria and application to fingerprinting of bacterial genomes. *Nucleic Acids Research*, **19**, 6823-6831.
- Vogel, S., Gobeze, A. and Gifawesen, C.** (1977). *Enjera*-A fermented pancake-like bread of Ethiopia. Symposium on Indigenous Fermented Foods, Bangkok, Thailand.
- Von Wintzingerode, F., Gobel, U.B. and Stackebrandt, E.** (1997). Determination of microbial diversity in environmental samples: Pitfalls of PCR-based rRNA analysis. *FEMS Microbiology Reviews*, **21**, 213-229.

Vos, P., Hogers, R., Bleeker, M., Reijans, M., van de Lee, T., Hornes, M., Frijters, A., Pot, J., Peleman, J., Muiper, M. And Zabeau, M. (1995). AFLP: A new concept for DNA fingerprinting. *Nucleic Acids Research*, **21**, 4407-4414.

Walter, J., Hertel, C., Tannock, G.W., Lis, C.M., Munro, K. and Hammes, W.P. (2001). Detection of *Lactobacillus*, *Pediococcus*, *Leuconostoc*, and *Weissella* species in human feces by using group-specific PCR primers and denaturing gradient gel electrophoresis. *Applied and Environmental Microbiology*, **67**, 2578-2585.

Wang, J. and Fung, D.Y. (1996). Alkaline-fermented foods: A review with emphasis on pidan fermentation. *Critical Reviews in Microbiology*, **22**, 101-138.

Welsh, J. and McClelland, M. (1990). Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Research*, **25**, 7212-7218.

Whiley, R.A., Hall, L.M.C., Hardie, J.M. and Beighton, D. (1999). A study of small-colony, β -haemolytic, Lancefield group C streptococci within the anginosus group: Description of *Streptococcus constellatus* subsp. *pharyngis* subsp. nov., associated with the human throat and pharyngitis. *International Journal of Systematic Bacteriology*, **49**, 1443-1449.

Williams, J.G.K., Kubelik, A.R., Livak, K.J., Rafalsky, J.A. and Tingey, S.V. (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Research*, **18**, 6531-6535.

Wodzinski, R.J. and Ullah, A.H.J. (1996). Phytase. *Advances in Applied Microbiology*, **42**, 263-302.

Woese, C.R. (1987). Bacterial evolution. *Microbiology Reviews*, **51**, 221-271.

Wood, B. and Warner, P. (2003). Genetics of lactic acid bacteria. Kluwer Academic Press, New York, USA.

Yusof, R.M., Morgan, J.B. and Adams, M.R. (1993). Bacteriological safety of a fermented weaning food containing L-lactate and nisin. *Journal of Food Protection*, **56**, 414-417.

Zabeau, M. and Vos, P. (1993). Selective restriction fragment amplification: A general method for DNA fingerprinting. European Patent Office, publication 0534858 A1.

Zegeye, A. (1997). Acceptability of *injera* with stewed chicken. *Food Quality and Preference*, **8**, 293-295.



CHAPTER TWO

**POLYPHASIC TAXONOMIC CHARACTERIZATION
OF LACTIC ACID BACTERIA ISOLATED FROM
SPONTANEOUS SORGHUM FERMENTATIONS USED
TO PRODUCE *TING*, A TRADITIONAL SOUTH
AFRICAN FOOD**

TABLE OF CONTENTS

2.1 INTRODUCTION	58
2.2 MATERIALS AND METHODS	59
2.2.1 Sorghum fermentation and LAB isolation	59
2.2.2 Phenotypic identification of LAB	59
2.2.3 DNA-based identification of LAB	60
2.2.3.1 Polymerase chain reaction (PCR)	60
2.2.3.2 Nucleotide sequencing and sequence analyses	60
2.2.3.3 Phylogenetic analyses	61
2.3 RESULTS	61
2.3.1 Phenotypic identification of LAB	61
2.3.2 DNA-based identification of LAB	61
2.4 DISCUSSION	63
2.5 REFERENCES	66

ABSTRACT

Ting, an indigenous cooked fermented food made from sorghum flour, is consumed extensively in South Africa. Due to the spontaneous nature of the sorghum fermentation considerable variations in the sensory and microbiological quality of the end-product may occur, thus hampering large-scale production of this food. The use of starter cultures purified from the fermented sorghum may be an alternative approach to obtain *ting* of consistent quality. The aim of this study was therefore to identify the lactic acid bacteria (LAB) associated with *ting* fermentation using a polyphasic approach. Phenotypic characterization and sequence analysis of the genes encoding the 16S subunit of the ribosomal RNA (*rrs*) and phenylalanyl tRNA synthase (*pheS*) were used. The results of these analyses showed that *ting* fermentation involved at least three different species of *Lactobacillus*, i.e. *Lb. fermentum*, *Lb. plantarum* and *Lb. rhamnosus*. This represents the first report of polyphasic taxonomic characterization of LAB from this food. This research forms an essential first step towards the development of relevant starter cultures to produce *ting* of consistent quality.

Key words: lactic acid bacteria; polyphasic taxonomy; sorghum fermentation; *ting*; *pheS*

2.1 INTRODUCTION

Ting is a fermented sour porridge made in rural South Africa (Boling and Eisener, 1982). Adults typically consume a thick form of the porridge at major ceremonies such as weddings and funerals, while a diluted form may be used as weaning food because it is relatively inexpensive to prepare and does not require refrigeration or re-heating prior to consumption (Kunene *et al.*, 1999). *Ting* is prepared by soaking sorghum, millet or maize meal for two to three days in excess water at ambient temperature, followed by cooking to make the porridge (Boling and Eisener, 1982). During the soaking step, lactic acid bacteria (LAB) associated with the ingredients and utensils are responsible for fermentation and the resulting flavour of *ting* (Tamang, 1998; M'hir *et al.*, 2007). As these fermentations are uncontrolled, the quality and sensory properties of different *ting* preparations may vary. However, the use of starter cultures could circumvent this problem and also facilitate the large-scale production and marketing of this food.

The development of starter cultures for a specific fermentation process usually involves isolation, selection and purification of microorganisms obtained from the fermented material (Mugula *et al.*, 2003a). The selected microorganisms are then propagated as a mixture of pure cultures or as single-strain starters (Mugula *et al.*, 2003a). Starter cultures that were developed from wild-type strains are generally better adapted to the specific food matrix (Holzapfel, 2002) and ensure that the end-product will have the expected flavour and aroma (Ayad *et al.*, 1999). However, fermentation processes usually require constant monitoring to ensure that the microorganisms involved remain competitive and retain their desired properties (Holzapfel, 2002). Mutations or the loss of important plasmids and contamination by other microorganisms may lead to a product of inferior quality. Accurate characterization of strains used as starter cultures is therefore of paramount importance.

Identification of LAB may be achieved using classical microbiological methods that are relatively simple to perform, but they often lack discriminatory power and reproducibility at species level (Ehrmann and Vogel, 2005). Genomic approaches using DNA fingerprint and/or sequence analyses, on the other hand, usually offer a higher level of accuracy, depending on the DNA regions targeted (Gevers *et al.*, 2001; Ehrmann and Vogel, 2005). For reliable species identification, however, a polyphasic approach is preferred in which both phenotypic and genotypic methods are used to obtain unambiguous identifications

(Vandamme *et al.*, 1996). The objectives of this study were to isolate the LAB responsible for the *ting* fermentation process, and to identify them using classical microbiological and DNA-based identification methods. For the latter, sequence analyses of the genes encoding 16S ribosomal RNA (*rrs*) and phenylalanyl tRNA synthase (*pheS*) were used.

2.2 MATERIALS AND METHODS

2.2.1 Sorghum fermentation and LAB isolation

Commercial packets of pure grain sorghum powder ('King Korn Mabele', King Food Corporation, Potchefstroom, South Africa) were obtained from a local supermarket. The sorghum powder was mixed (1:1 [w/v]) with luke-warm (ca. 40°C) water in glass containers to obtain a slurry. The containers were covered and incubated at 25°C for 54 h after which the pH of the mixture was measured with a Beckman model Ø 34 pH meter (Beckman Coulter, Fullerton, CA, USA). Using this approach, the experiments were carried out in triplicate.

For each fermentation, 10 g of the fermented slurry was aseptically removed at 54 h (when the pH of the slurries was around 3.8) and homogenized in 90 ml of quarter strength Ringer's solution (DAB 7, Braun, Melsungen, Germany). This 10⁻¹ dilution was then serially diluted to 10⁻⁸ using the same diluent, after which 1 ml of each of the diluted samples was inoculated in duplicate onto MRS agar (Oxoid, Basingstoke, UK). For the isolation of thermophilic lactobacilli, inoculated plates were anaerobically incubated using Oxoid's Anaerobic Gas Generating kit at 42°C for 48 h. For isolation of the mesophilic lactobacilli, the plates were anaerobically incubated at 30°C for 48 h. Colonies were purified and stored at -20°C in MRS broth supplemented with 20% glycerol.

2.2.2 Phenotypic identification of LAB

Gram-staining, motility, oxidase and catalase tests (Ehrlich, 1956) were performed for nine representative isolates. The ability of isolates to produce gas from glucose was also determined. To determine the carbohydrate fermentation profiles of the bacteria, API 50 CHL test kits (bioMérieux, France) were used according to the manufacturer's instructions. Isolates were identified to species level using the API database and accompanying software (Johansson *et al.*, 1995). Based on these identifications, six isolates were selected for sequence analysis.

2.2.3 DNA-based identification of LAB

2.2.3.1 Polymerase chain reactions (PCR)

LAB isolates were inoculated into MRS broth and incubated at 30°C overnight. DNA was extracted from these overnight cultures using the Wizard[®] Genomic DNA extraction kit (Promega, Madison, USA) or the phenol-chloroform method described by Gevers *et al.* (2001). Fragments of the 16S rRNA and *pheS* genes were amplified using the eubacterial universal primers 27F (5'-agagtttgatcctggctcag-3'; Lane, 1991) and 1507R (5'-tacctgttacgacttcaccca-3'; Heyndrickx *et al.*, 1996), and the primers PheS-21-F (5'-cayccngchcgycgayagatgc-3') and PheS-23-R (5'-ggtrgraccatvccngchcc-3') described by Naser *et al.* (2005), respectively. The *pheS* PCR was performed as described by Naser *et al.* (2005). For 16S rRNA genes, PCR reaction mixtures (25 µl) contained 50 ng of the extracted DNA, 1.0 mM MgCl₂, 0.2 mM of each dNTP, 0.4 µM of each of primers 27F and 1507R, 8% (v/v) dimethyl sulphoxide (DMSO), 1.25 U of *Taq* DNA polymerase (Fermentas, St. Leon-Rot, Germany) and PCR buffer containing NH₂SO₄ (Fermentas). The temperature programme consisted of an initial denaturation step at 95°C for 10 min, followed by 35 cycles of denaturation at 94°C for 1 min, primer annealing at 51°C for 30 s and primer extension at 72°C for 1 min, with a final extension at 72°C for 10 min.

2.2.3.2 Nucleotide sequencing and sequence analyses

The 16S rRNA gene amplicons were purified with a High Pure PCR Product Purification kit (Roche Applied Science, Penzberg, Germany) and sequenced in both directions with the primers used in PCR. For this purpose, the BigDye[®] Terminator v. 3.1 Cycle Sequencing kit (Applied Biosystems) and an ABI PRISM[™] 3100 DNA sequencer (Applied Biosystems) were used. The *pheS* amplicons were purified using the Nucleofast 96 PCR clean-up membrane system (Machery-Nagel, Germany) and sequenced, as described above, with the *pheS* PCR primers. The resulting sequences were analyzed using Chromas Lite 2.0 (Technelysium) and BioEdit v. 5.0.9 (Hall, 1999). The sequences were then compared to those in GenBank (National Centre for Biotechnology Information; www.ncbi.nih.gov) using the Basic Local Alignment Search Tool (Altschul *et al.*, 1990) for nucleotide sequences (*blastn*).

2.2.3.3 Phylogenetic analyses

For phylogenetic analyses, multiple alignments were generated using Multiple Sequence Alignment, which is based on Fast Fourier Transform (MAFFT v. 6) (Kato *et al.*, 2005). These alignments included the *ting* LAB sequences, as well as 16S rRNA and *pheS* gene sequences for the *Lactobacillus* type strains for the *Lb. reuteri*, *Lb. plantarum* and *Lb. casei* Groups (Naser *et al.* 2007) obtained from GenBank. Phylogenetic relationships based on maximum likelihood (ML) were inferred with PhyML v. 2.4.3 (Guindon and Gascuel, 2003). As indicated by Modeltest v. 3.7 (Posada and Crandall, 1998), ML analysis of the 16S rRNA gene sequence dataset utilized Tamura and Nei's (1993) substitution model with a proportion of invariable sites (I) and gamma correction for among site variation (G), while the *pheS* dataset utilized the general time reversible (GTR) model (Tavaré, 1986) with I and G. The same best-fit parameters were also used for bootstrap analyses based on 1000 replicates.

2.3 RESULTS

2.3.1 Phenotypic identification of LAB

All the selected isolates were catalase-negative, non-motile and Gram-positive, and were thus tentatively assigned to LAB. Among the nine representative LAB isolates, isolates 3.42, 4.35, 3.30 and 5.30 produced gas from glucose, while isolates 1.42, 5.42, 5.35, 4.30 and 2.35 did not produce gas from glucose. All nine isolates constituted short rods. Using the API 50 CHL system, the nine isolates were shown to utilize some of the sugars tested (Table 2.1). *L*-arabinose, melibiose and sucrose were fermented by all isolates, except isolate 5.30. *N*-acetyl glucosamine was fermented by all isolates, except isolates 3.30 and 3.42. Gluconate was fermented by all isolates, except isolates 2.35 and 5.30. The API 50 CHL identification system assigned the isolates to *Lb. plantarum* (isolates 1.42, 5.42, 5.35 and 4.30), *Lb. fermentum* (isolates 3.42, 4.35, 3.30 and 5.30) and *Lb. rhamnosus* (isolate 2.35). From these, six isolates (1.42, 3.42, 2.35, 4.35, 3.30 and 5.30) were selected for further analysis.

2.3.2 DNA-based identification of LAB

Comparison of the 16S rRNA gene sequences for the six selected isolates (GenBank accession numbers EU825657 through EU825662) from the fermented sorghum slurries to those in GenBank indicated that they all represent *Lactobacillus* species. The *blastn* results showed that the 16S rRNA gene sequence for isolate 1.42 was most similar to those of the

Table 2.1 Differential characteristics of nine LAB isolates from fermented sorghum, based on API 50 CHL analysis

Isolate	API 50 CHL Results*								
	1.42	3.42	5.42	2.35	4.35	5.35	3.30	4.30	5.30
Substrate tested									
<i>L</i> -arabinose	+	+	+	+	+	+	+	+	-
Mannose	+	+ ¹	+	+	+ ¹	+	+ ¹	+	+ ¹
<i>N</i> -acetyl glucosamine	+	-	+	+	+	+	-	+	+
Lactose	+ ²	+	+ ²	+	+	+ ²	+	+ ²	+
Melibiose	+	+	+	+	+	+	+	+	-
Sucrose	+	+	+	+	+	+	+	+	-
Raffinose	+	+	+	+	+	+	+	+	+
Gluconate	+	+	+	-	+	+	+	+	-
API 50 CHL identification	<i>Lb. plantarum</i>	<i>Lb. fermentum</i>	<i>Lb. plantarum</i>	<i>Lb. rhamnosus</i>	<i>Lb. fermentum</i>	<i>Lb. plantarum</i>	<i>Lb. fermentum</i>	<i>Lb. plantarum</i>	<i>Lb. fermentum</i>

* +, positive reaction; -, no reaction.

^{1,2} Weak reaction after 48 h.

so-called *Lb. plantarum* Group (Naser *et al.*, 2007) that includes *Lb. plantarum*, *Lb. paraplantarum* and *Lb. pentosus*. The 16S rRNA gene sequence for isolate 2.35 was most similar to those of the *Lb. casei* Group (Naser *et al.*, 2007) that includes *Lb. casei*, *Lb. paracasei*, *Lb. zaeae* and *Lb. rhamnosus*. The *blastn* results for isolates 3.42, 4.35, 3.30 and 5.30 indicated that their 16S rRNA gene sequences were most similar to that of *Lb. fermentum* in the *Lb. reuteri* Group (Naser *et al.*, 2007). ML analysis of the 16S rRNA gene sequence data indicated that isolates 3.42, 4.35, 3.30 and 5.30 were indeed most closely related to *Lb. fermentum* (Fig. 2.1A). The 16S rRNA gene sequence-based ML analysis did, however, not allow discrimination of isolates 2.35 and 1.42 from other *Lb. plantarum* Group and *Lb. casei* Group species, respectively. ML analysis of the *pheS* dataset containing sequences for the six selected LAB isolates (GenBank accession numbers EU825663 through EU825668) recovered similar groupings, but provided better resolution among the taxa included (Fig. 2.1B). Within the *Lb. plantarum* Group, isolate 1.42 grouped with known *Lb. plantarum* strains, and within the *Lb. casei* Group, isolate 2.35 grouped with *Lb. rhamnosus*.

2.4 DISCUSSION

This represents the first study investigating the identity of LAB isolated from *ting* fermentation using a polyphasic approach that combines API 50 CHL analysis and sequence-based identification. The polyphasic approach used in this study aimed to obtain an unequivocal identification of a selection of LAB isolates recovered from the fermentation of sorghum to produce *ting*. The results suggested that three *Lactobacillus* species, i.e. *Lb. plantarum*, *Lb. fermentum* and *Lb. rhamnosus*, are associated with this fermentation process. It has previously been reported that the API 50 CHL system does not allow accurate identification of the majority of LAB isolates. These findings have been attributed to the possible loss or acquisition of plasmids that encode many carbohydrate fermentation traits, resulting in isolates exhibiting atypical metabolic characteristics (Ahrné *et al.*, 1989). Moreover, carbohydrate fermentation patterns across the genus *Lactobacillus* have also been shown to be incongruent with the results of DNA-based studies (Boyd *et al.*, 2005). In addition, the subjective interpretation of similar carbohydrate fermentation patterns obtained using the API 50 CHL system may lead to atypical results (Randazzo *et al.*, 2004). However, in this study, use of the API 50 CHL system allowed identification of the LAB isolates to species level, the results of which correlated well with the sequence-based identification data.




Figure 2.1 Maximum likelihood (ML) phylogeny for the *Lb. plantarum*, *Lb. casei* and *Lb. reuteri* Groups (Naser *et al.*, 2007), based on 16S rRNA (A) and *pheS* (B) gene sequences. Isolates from fermented sorghum slurries are indicated in bold and bootstrap values >60%, based on 1000 replicates, are indicated at the internodes. For the various *Lactobacillus* species, type strain numbers and GenBank accession numbers for the 16S rRNA and *pheS* sequences, respectively, are as follows: *Lb. antri* LMG22111T, AY253659, AM263502; *Lb. casei* LMG6904T, M58815, AM087682; *Lb. coleohominis* LMG21591T, AJ292530, AM087683; *Lb. durianis* LMG19193T, AJ315640, AM087739; *Lb. fermentum* LMG6902T, M58819, AM087693; *Lb. frumenti* LMG19473T, AJ250074, AM087741; *Lb. gastricus* LMG22113T, AY253658, AM087696; *Lb. ingluvei* LMG20380T, AF333975, AM087731; *Lb. mucosae* LMG19534T, AF126738, AM087707; *Lb. oris* LMG9848T, X94229, AM087709; *Lb. panis* LMG21658T, X94230, AM087725; *Lb. paracasei* LMG13087T, D79212, AM087710; *Lb. paraplanctarium* LMG16673T, AJ306297, AM087727; *Lb. pentosus* LMG10755T, AB289240, AM087713; *Lb. plantarum* LMG6907T, X52653, AM087714; *Lb. plantarum** LMG19807T, AJ965482, AM087736; *Lb. plantarum* subsp. *argenteratensis* LMG9205T, AJ640078, AM694185; *Lb. pontis* LMG14187T, AJ422032, AM087715; *Lb. rhamnosus* LMG6400T, AB008211, AM087716; *Lb. reuteri* LMG9213T, X76328, AM087728; *Lb. rossiae* LMG22972T, AB370880, AM087768; *Lb. suebicus* LMG11408T, AM113785, AM087772; *Lb. vaccinostercus* LMG9215T, AJ621556, AM087750; *Lb. vaginalis* LMG12891T, AF243177, AM087751; *Lb. zaeae* LMG17315T, D86516, AM087761 (LMG = Laboratory of Microbiology, Ghent University, Belgium). *Lb. arizonensis* (type strain LMG19807) that was shown to represent a later heterotypic synonym of *Lb. plantarum* (Kostinek *et al.*, 2005) is indicated with an asterisk. The GenBank accession numbers for *pheS* and 16S rRNA gene sequences for the outgroup taxa *Weissella viridescens* (LMG3507T) and *Bacillus subtilis* (LMG7135T) are AM711182 and X60646, respectively.

Despite forming an integral part of all bacterial classifications (Vandamme *et al.*, 1996), ribosomal gene sequences may not always allow clear-cut LAB identifications as these genes are highly conserved (Felis and Dellaglio, 2007; De Vuyst and Vancanneyt, 2007). Among the six *ting* LAB isolates examined, only four could be identified with some level of certainty using 16S rRNA gene sequence data (Fig. 2.1A). The other two isolates formed part of the so-called *Lb. plantarum* and *Lb. casei* Groups (Naser *et al.*, 2007). As is the case for many other bacterial taxa, the members of these groups are known to encode 16S rRNA gene sequences that are not sufficiently polymorphic to allow species separation (Torriani *et al.*, 2001; Gevers *et al.*, 2005; Felis and Dellaglio, 2007). Therefore, bacterial species are increasingly being defined using housekeeping loci that evolve more rapidly than 16S rRNA genes (Gevers *et al.*, 2005). For LAB taxonomy, a number of alternative loci have been evaluated for taxonomic purposes and include protein-coding genes such as *tuf* (elongation factor Tu; Chavagnat *et al.*, 2002), *mal* (malolactic enzyme; Groisillier and Lonvaud-Funel, 1999), *pepC* (aminopeptidase C; Fortina *et al.*, 2001), *pepN* (aminopeptidase N; Fortina *et al.*, 2001), *htrA* (stress-inducible trypsin-like serine protease; Fortina *et al.*, 2001), *recA* (recombinase A; Felis *et al.*, 2001), *rpoB* (RNA polymerase beta subunit; Naser *et al.*, 2005), *hsp60* (60-kDa heat shock protein; Blaiotta *et al.*, 2008), and *pheS* (phenylalanyl tRNA synthase; Naser *et al.*, 2005). Most recently, Naser *et al.* (2007) examined the *pheS* and *rpoB* gene sequences of 201 strains representing 98 *Lactobacillus* species and demonstrated that these regions represent highly informative taxonomic markers for the identification of *Lactobacillus* species. In the current study, it was also found that *pheS* has greater discriminatory power, because it displayed sufficient interspecific variation that allowed the unequivocal identification of isolates 2.35 and 1.42 as *Lb. rhamnosus* and *Lb. plantarum*, respectively (Fig. 2.1B).

Of the three *ting* LAB species identified, *Lb. fermentum* and *Lb. plantarum* are commonly associated with a wide range of African traditional food and beverage fermentations (Steinkraus, 1996), including *fufu* (fermented cassava), *iru* (fermented African locust bean), *kenkey* and *ogi* (fermented maize), *kukun-zaki* (fermented millet), *ugba* (fermented African oil bean), and *wara* (fermented skimmed cow's milk). In fact, *Lb. plantarum* is the species most commonly isolated (Olasupo *et al.*, 1997) and has been identified as the dominant organism at the end of several natural lactic acid fermentations (Nout, 1980; Brauman *et al.*, 1996; Olasupo *et al.*, 1997; Mugula *et al.*, 2003b). This is probably due to its acid tolerance and superior ability to utilize the substrates involved (Akinrele, 1970; Oyewole and Odunfa,

1990). It should be possible to include the characterized isolates in starter cultures that would result in products with consistent microbiological and sensory properties. Ultimately, these starter cultures may also be used for upgrading this subsistence bioprocess technology to large-scale industrial production and marketing of *ting*. The entire process is therefore important, not only from an academic viewpoint, but also for the conservation of indigenous knowledge and technologies through the characterization and preservation of the microflora associated with this traditional fermented food.

2.5 REFERENCES

Ahrné, S., Molin, G. and Ståhl, S. (1989). Plasmids in *Lactobacillus* strains isolated from meat and meat products. *Systematic and Applied Microbiology*, **11**, 320-325.

Altschul, S.F., Gish, W., Myers, E.W. and Lipman, D.J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, **215**, 403-410.

Akinrele, A. (1970). Fermentation studies on maize during the preparation of a traditional African starch-cake food. *Journal of the Science of Food Agriculture*, **21**, 619-625.

Ayad, E.H.E., Verheul, A., de Jong, C., Wouters, J.T.M. and Smit, G. (1999). Flavour-forming abilities and amino acid requirements of *Lactococcus lactis* strains isolated from artisanal and non-dairy origin. *International Dairy Journal*, **9**, 725-735.

Blaiotta, G., Fusco, V., Ercolini, D. and Aponte, M. (2008). *Lactobacillus* strain diversity based on partial *hsp60* gene sequences and design of PCR-restriction fragment length polymorphism assays for species identification and differentiation. *Applied and Environmental Microbiology*, **74**, 208-215.

Boling, M.B. and Eisener, N. (1982). *Bogobe*: Sorghum porridge of Botswana. In: Proceedings of the International Symposium on Sorghum Grain Quality. International Crops Research Institute for the Semi-Arid Tropics. Mertin, J.V. (Ed.). Patancheru, India, pp. 32-35.

Boyd, M.A., Antonio, M.A.D. and Hillier, S.L. (2005). Comparison of API 50 CH strips to whole-chromosomal DNA probes for identification of *Lactobacillus* species. *Journal of Clinical Microbiology*, **43**, 5309-5311.

Brauman, A., Keleke, S., Malonga, M., Miambi, E. and Ampe, F. (1996). Microbiological and biochemical characterization of cassava retting, a traditional lactic acid fermentation for *foo-foo* (cassava powder) production. *Applied and Environmental Microbiology*, **62**, 2854-2858.

- Chavagnat, F., Haueter, M., Jimeno, J. and Casey, M.G.** (2002). Comparison of partial *tuf* gene sequences for the identification of lactobacilli. *FEMS Microbiology Letters*, **217**, 177-183.
- De Vuyst, L. and Vancanneyt, M.** (2007). Biodiversity and identification of sourdough lactic acid bacteria. *Food Microbiology*, **24**, 120-127.
- Ehrlich, P.** (1956). The requirement of the organism for oxygen. An analytical study with the aid of dyes. In: *Collected papers of Paul Ehrlich*. Himmelweit, F., Marquardt, M. and Dale, H. (Eds.). Pergamon Press, London, UK, pp. 433-496.
- Ehrmann, M.A. and Vogel, R.F.** (2005). Molecular taxonomy and genetics of sourdough lactic acid bacteria. *Trends in Food Science and Technology*, **16**, 31-42.
- Felis, G.E., Dellaglio, F., Mizzi, L. and Torriani, S.** (2001). Comparative sequence analysis of *recA* gene fragment brings new evidence for a change in taxonomy of the *Lactobacillus casei* group. *International Journal of Systematic and Evolutionary Microbiology*, **51**, 2113-2117.
- Felis, G.E. and Dellaglio, F.** (2007). Taxonomy of *Lactobacilli* and *Bifidobacteria*. *Current Issues in Intestinal Microbiology*, **8**, 44-61.
- Fortina, M.G., Ricci, G., Mora, D., Parini, C. and Manachini, P.L.** (2001). Specific identification of *Lactobacillus helveticus* by PCR with *pepC*, *pepN* and *htrA* targetted primers. *FEMS Microbiology Letters*, **198**, 85-89.
- Gevers, D., Huys, G. and Swings, J.** (2001). Applicability of rep-PCR fingerprinting for identification of *Lactobacillus* species. *FEMS Microbiology Letters*, **205**, 31-36.
- Gevers, D., Cohan, F.M., Lawrence, J.G., Spratt, B.G., Coenye, T., Feil, E.J., Stackebrandt, E., Van de Peer, Y., Vandamme, P., Thompson, F.L. and Swings, J.** (2005). Re-evaluating prokaryotic species. *Nature Reviews Microbiology*, **3**, 733-739.
- Groisillier, A. and Lonvaud-Funel, A.** (1999). Comparison of partial malolactic enzyme gene sequencing for phylogenetic analysis of some lactic acid bacteria species and relationships with the malic enzyme. *International Journal of Systematic Bacteriology*, **49**, 1417-1428.
- Guindon, S. and Gascuel, O.** (2003). A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology*, **52**, 696-704.
- Hall, T.A.** (1999). BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, **41**, 95-98.

- Heyndrickx, M., Vaterin, L., Vandamme, P., Kersters, K. and De Vos, P.** (1996). Applicability of combined amplified ribosomal DNA restriction analysis (ARDRA) patterns in bacterial phylogeny and taxonomy. *Journal of Microbiological Methods*, **26**, 247-259.
- Holzappel, W.H.** (2002). Appropriate starter culture technologies for small-scale fermentation in developing countries. *International Journal of Food Microbiology*, **75**, 197-212.
- Johansson, M.L., Sanni, A., Lonner, C. and Molin, G.** (1995). Phenotypically-based taxonomy using API 50 CH of lactobacilli from Nigerian *ogi*. *International Journal of Food Microbiology*, **25**, 159-168.
- Katoh, K., Kuma, T.H. and Miyata, T.** (2005). MAFFT version 5: Improvement in accuracy of multiple sequence alignment. *Nucleic Acids Research*, **33**, 511-518.
- Kostinek, M., Pukall, R., Rooney, A.P., Schillinger, U., Hertel, C., Holzappel, W.H. and Franz, C.M.A.P.** (2005). *Lactobacillus arizonensis* is a later heterotypic synonym of *Lactobacillus plantarum*. *International Journal of Systematic and Evolutionary Microbiology*, **55**, 2485-2489.
- Kunene, N.F., Hastings, J.E. and von Holy, A.** (1999). Bacterial populations associated with a sorghum-based fermented weaning cereal. *International Journal of Food Microbiology*, **49**, 75-83.
- Lane, D.L.** (1991). 16S/23S rRNA sequencing. In: *Nucleic acid techniques in bacterial systematics*. Stackebrandt, E. and Goodfellow, M. (Eds.). John Wiley and Sons, New York, USA, pp. 115-175.
- Mugula, J.K., Narvhus, J.A. and Sørhaug, T.** (2003a). Use of starter cultures of LAB and yeasts in the preparation of *togwa*, a Tanzanian fermented food. *International Journal of Food Microbiology*, **83**, 307-318.
- Mugula, J.K., Nnko, S.A.M., Narvhus, J.A. and Sørhaug, T.** (2003b). Microbiological and fermentation characteristics of *togwa*, a Tanzanian fermented food. *International Journal of Food Microbiology*, **80**, 187-199.
- M'hir, S., Mejri, M. and Hamdi, M.** (2007). Microflora distribution and species ratio of Tunisian fermented doughs for bakery industry. *African Journal of Biotechnology*, **6**, 2122-2129.
- Naser, S.M., Thompson, F.L., Hoste, B., Gevers, D., Dawyndt, P., Vancanneyt, M. and Swings, J.** (2005). Application of multilocus sequence analysis (MLSA) for rapid identification of *Enterococcus* species based on *rpoA* and *pheS* genes. *Microbiology*, **151**, 2141-2150.

- Naser, S.M., Dawyndt, P., Hoste, B., Gevers, D., Vandemeulebroecke, K., Cleenwerk, I., Vancanneyt, M. and Swings, J.** (2007). Identification of lactobacilli by *pheS* and *rpoA* gene sequence analyses. *International Journal of Systematic and Evolutionary Microbiology*, **57**, 2777-2789.
- Nout, M.J.R.** (1980). Microbiological aspects of the traditional manufacture of *busaa*, a Kenyan opaque maize beer. *Chemie Mikrobiologie und Technologie Lebensmitteln*, **6**, 137-142.
- Olasupo, N.A., Olukoya, D.K. and Odunfa, S.A.** (1997). Identification of *Lactobacillus* species associated with selected African fermented foods. *Zeitschrift fur Naturforschung*, **52**, 105-108.
- Oyewole, O.B. and Odunfa, S.A.** (1990). Characterization and distribution of lactic acid bacteria in cassava fermentation during *fufu* production. *Journal of Applied Bacteriology*, **68**, 145-152.
- Posada, D. and Crandall, K.A.** (1998). Modeltest: Testing the model of DNA substitution. *Bioinformatics*, **14**, 817-818.
- Randazzo, C.L., Restuccia, C., Romano, A.D. and Caggia, C.** (2004). *Lactobacillus casei*, dominant in naturally fermented Sicilian green olives. *International Journal of Food Microbiology*, **90**, 9-14.
- Steinkraus, K.H.** (1996). *Handbook of indigenous fermented foods*. Marcel Dekker, Inc., New York, USA.
- Tamang, J.P.** (1998). Role of microorganisms in traditional fermented foods. *Indian Food Industry*, **17**, 162-167.
- Tamura, K. and Nei, M.** (1993). Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution*, **10**, 512-526.
- Tavaré, S.** (1986). Some probabilistic and statistical problems in the analysis of DNA sequences. In: *Some mathematical questions in biology – DNA sequence analysis*. Miura, R.M. (Ed.). American Mathematical Society, Providence, Rhode Island, USA, pp. 57-86.
- Torriani, S., Felis, G.E. and Dellaglio, F.** (2001). Differentiation of *Lactobacillus plantarum*, *L. pentosus*, and *L. paraplantarum* by *recA* gene sequence analysis and multiplex PCR assay with *recA* gene-derived primers. *Applied and Environmental Microbiology*, **67**, 3450-3454.
- Vandamme, P., Pot, B., Gillis, M., de Vos, P., Kersters, K. and Swings, J.** (1996). Polyphasic taxonomy, a consensus approach to bacterial systematics. *Microbiology Reviews*, **60**, 407-438.



CHAPTER THREE

DIVERSITY AND DYNAMICS OF BACTERIAL POPULATIONS DURING SPONTANEOUS SORGHUM FERMENTATIONS USED TO PRODUCE *TING*, A SOUTH AFRICAN FOOD

TABLE OF CONTENTS

3.1 INTRODUCTION	73
3.2 MATERIALS AND METHODS	74
3.2.1 Sorghum fermentations	74
3.2.2 PCR-DGGE analyses	75
3.2.3 Enumeration, isolation and characterization of bacteria	76
3.2.4 PCR and sequencing of the 16S rRNA and <i>pheS</i> genes from pure cultures ...	77
3.2.5 Sequence and phylogenetic analyses.....	78
3.3 RESULTS	78
3.3.1 Enumeration and presumptive identification of bacteria	78
3.3.2 PCR-DGGE analyses	79
3.3.3 Phylogenetic analyses using 16S rRNA and <i>pheS</i> gene sequences	83
3.3.4 Detection of bacterial diversity using culture-independent and culture- dependent approaches	87
3.4 DISCUSSION	87
3.5 REFERENCES	92

ABSTRACT

Ting is a spontaneously fermented cooked South African food that is popular for its sour taste and unique flavour. Insight of microbial diversity and population dynamics during sorghum fermentations is an initial but imperative step in the development of starter cultures for commercial production of *ting*. In this study, microorganisms associated with spontaneous fermentations of two sorghum types were examined using culture-independent and culture-dependent methods. Culture-independent polymerase chain reaction (PCR) of the V3 region of the 16S rRNA gene and analysis of the resulting amplicons using denaturing gradient gel electrophoresis (DGGE), followed by sequence analyses of the most intense bands, revealed that lactic acid bacteria (LAB), including close relatives of *Lactococcus lactis*, *Lactobacillus curvatus*, *Weissella cibaria* and some members of the *Enterobacteriaceae* family, were predominant at the end of fermentation in both sorghum types. *L. lactis*, however, showed the most intense band. Culture-dependent methods, involving isolation of the various bacteria, followed by 16S rRNA gene sequence analyses indicated that *Enterococcus mundtii* was present at the start of fermentation, whilst *Lb. fermentum*, *Lb. plantarum*, *Lb. rhamnosus*, *E. faecalis*, *W. cibaria* and *L. lactis* were predominant at the end of fermentation. *Lb. curvatus* was not detected in culture media, but it was predominant in DGGE gels. Likewise, *Lb. fermentum*, *Lb. plantarum*, *Lb. rhamnosus*, *E. faecalis* and *E. mundtii* were isolated in culture media, but they were absent in DGGE gels. Therefore, the combined approach was effective in revealing the microbial diversity and dynamics during spontaneous sorghum fermentations.

Key words: culture-independent; culture-dependent; microbial diversity; population dynamics; PCR-DGGE; fermentation

3.1 INTRODUCTION

Sorghum is widely cultivated in arid and semi-arid regions of the world, and is considered to be the fifth most important cereal after wheat, maize, rice and barley (Food and Agricultural Organisation, 2006). However, sorghum as a main dietary constituent is usually associated with under-nourishment due to the lack of certain essential amino acids (Neucere and Sumrell, 1979; Kazanas and Fields, 1981) and the presence of anti-nutritional factors (Hamacker *et al.*, 1986; Ibrahim *et al.*, 2005). These shortcomings may be overcome by fermentation that is primarily facilitated by lactic acid bacteria (LAB) (Au and Fields, 1981; Chavan *et al.*, 1988; Hassan and El Tinay, 1995). In addition to improving the nutritive value of this commodity, fermentation also enhances its sensory properties (Au and Fields, 1981; Chavan *et al.*, 1988; Hassan and El Tinay, 1995). As a result, various sorghum-based fermented foods with unique and appealing characteristics are produced wherever sorghum is cultivated (Caplice and Fitzgerald, 1999; Taylor, 2003). These include *injera* (Gebrekidan and Gebrettiwat, 1982), *kisra* (Mohammed *et al.*, 1991), *ogi* (Akingbala *et al.*, 1981), *mahewu* (Bvochora *et al.*, 1999), *uji* (Mbugua, 1984), *muramba* (Mukuru, 1992), *bushera* (Muyanja, 2003), *togwa* (Lorri and Svanberg, 1995), and *ting* (Boling and Eisener, 1982).

Ting is a sour porridge made by cooking fermented sorghum (Boling and Eisener, 1982). It is frequently used as a weaning food for infants in rural South Africa because it is relatively inexpensive to prepare and does not require refrigeration or re-heating prior to consumption (Kunene *et al.*, 1999). Due to its appetizing taste, adults also consume *ting* at major ceremonies such as weddings and funerals. In such traditional preparations, sorghum undergoes spontaneous and uncontrolled fermentation steered by microflora endogenous to the sorghum, as well as those associated with the preparation equipment and local environments. Consequently, conventional *ting* preparations vary greatly with respect to product quality, taste and acceptability (Sanni, 1993). Also, little is known about the microorganisms that participate in this fermentation. Although bacteria were previously isolated from fermented sorghum and identified as *Lactobacillus plantarum*, *Lb. fermentum* and *Lb. rhamnosus* (Chapter 2), their involvement in the fermentation process and the possible role of other microbes remains to be determined. Such information is crucial for developing starter cultures that result in reduced fermentation time and *ting* with consistent microbiological quality.

To study the diversity and dynamics of microbial populations associated with specific fermented foods, a combination of culture-dependent and -independent approaches are typically applied. Culture-dependent approaches represents the only means of recovering microorganisms from the fermented substrate (Miambi *et al.*, 2003), although it is widely recognized that they do not allow analysis of true diversity and/or population dynamics (Ampe *et al.*, 1999; Meroth *et al.*, 2003; Nielsen *et al.*, 2007). On the other hand, various studies have shown that culture-independent approaches provide more reliable and reproducible (Ercolini, 2004) means for studying microbial populations in complex ecosystems such as food matrices (Ercolini, 2004; Rantsiou and Cocolin, 2006; Renouf *et al.*, 2006; Camu *et al.*, 2007). These approaches are based on methods such as PCR-DGGE (denaturing gradient gel electrophoresis; Muyzer *et al.*, 1993), SSCP (single-stranded conformation polymorphism; Orita *et al.*, 1989), T-RFLP (terminal-restriction fragment length polymorphism; Liu *et al.*, 1997) and LH-PCR (length heterogeneity-PCR; Ritchie *et al.*, 2000) analyses of a specific gene fragment (usually the 16S ribosomal RNA gene, *rrs*).

The aim of this study was to systematically and effectively describe the microbial populations associated with fermented sorghum, using a combination of culture-dependent and culture-independent methods. For the culture-independent analysis, PCR-DGGE was used to monitor the succession of dominant microbial populations from the onset to the end of spontaneous fermentations of sorghum. In parallel with the PCR-DGGE experiments, culture-dependent methods were used to enumerate, isolate and identify the bacteria involved in the fermentations by making use of phylogenetic analyses of the genes encoding 16S rRNA and phenylalanyl-tRNA synthase (*pheS*).

3.2 MATERIALS AND METHODS

3.2.1 Sorghum fermentations

In this study, fermentations were conducted at three different temperatures using sorghum flour obtained from two commercial sources in South Africa (King Food Corporation, Potchefstroom and Nola Pvt Ltd, Randfontein). For this purpose, individual sorghum slurries were prepared by mixing the respective sorghum flours thoroughly with sterile luke-warm (ca. 40°C) water (1:1 [w/v]) in sterile glass containers. The glass containers were then covered and incubated at 20, 25 and 30°C for 54 h. All fermentations were done in duplicate. During the incubation period, 50-ml samples of the fermenting sorghum slurries were aseptically

collected at the start of fermentation ($t = 0$ h) and after 6, 12, 18, 24, 30, 36, 42, 48 and 54 h of fermentation. The pH of a 15-ml aliquot of each sample was determined using a Beckman model Ø 34 pH meter (Beckman Coulter, Fullerton, CA, USA) by recording the averages of measurements taken in triplicate. For each of the collected samples, 15-ml aliquots were immediately frozen and stored at -20°C for later DNA extractions, while the remainder was used for culture-dependent analyses.

3.2.2 PCR-DGGE analyses

To prepare good quality DNA for the PCR-DGGE analyses, most of the sorghum was first removed from each of the various collected slurry samples, as follows. The individual 15-ml samples were vortexed for 10 min in the presence of glass beads (2% [v/v], 150 μm -diameter beads; Sigma-Aldrich) and then centrifuged for 5 min at $1\,500 \times g$ to remove the beads and large sorghum particles. The bacterial cells in 1.5 ml of the individual supernatants were harvested by centrifugation at $5\,000 \times g$ for 15 min. The pelleted cells were frozen at -20°C for 2 h and washed with sterile distilled water, after which DNA was extracted using the Wizard[®] Genomic DNA Purification Kit (Promega, Madison, USA).

For PCR-DGGE analyses, the V3 region of the bacterial 16S rRNA gene was targeted using the primers 518R (5'-attaccgcggtgg-3'; Muyzer *et al.*, 1993) and F357-GC. Primer F357-GC is similar to primer F357 (5'-tacgggaggcagcag-3', Rainey *et al.*, 1996), except that it includes a 40-base GC-clamp (5'-cgcccgccgcgcgcggcgggcgggggcgggggcacggggg-3'; Muyzer *et al.*, 1993) at its 5'-end to avoid complete melting of PCR products during DGGE (see below). PCR was performed according to Van der Meulen *et al.* (2007) on a MyCycler[™] thermal cycler (BioRad, Hercules, USA) with the following cycling conditions: denaturation at 95°C for 5 min, followed by 30 cycles of denaturation at 94°C for 20 s, annealing at 55°C for 45 s, extension at 72°C for 1 min and final extension for 7 min at 72°C .

The 16S rRNA V3 region was also amplified from 13 different bacteria and combined to use as a DGGE reference ladder (Van der Meulen *et al.*, 2007). These amplicons were generated using the same primers and cycling conditions as above. The bacteria comprised *Bacteriodes fragilis* (DSM 1396), *Bacteroides thetaiomicron* (LMG 11262), *Weissella cibaria* (LMG 17699), *Enterococcus flavescens* (LMG 13518T), *Enterococcus solitarius* (LMG 12890T), *Leuconostoc fructosum* (LMG 9498T), *Bacillus subtilis* (LMG 7135T), *Clostridium*

butyricum (LMG 1212), *Lactobacillus rhamnosus* (LMG 6400T), *Bifidobacterium longum* (LMG 13197T), *Bifidobacterium bifidum* (LMG 11041T), *Bifidobacterium lactis* (LMG 18314T), and *Bifidobacterium dentium* (LMG 11045T).

DGGE was essentially performed according to the procedure described by Muyzer *et al.* (1993) with slight modifications, as described previously (Van der Meulen *et al.*, 2007). The incorporation of a reference ladder on each DGGE gel allowed digital normalization of the band profiles by using the BioNumerics v. 4.0 software package and comparing with a standard reference (Van der Meulen *et al.*, 2007). Consequently, migration distances between different gels could be compared. All PCR-DGGE experiments were performed in duplicate.

To determine the identity of the bacteria for which individual 16S rRNA V3 PCR-DGGE fragments were generated, the most intense bands at each of the different fermentation sampling time points (t = 0 h to t = 54 h) were excised from DGGE gels. The gel slices were then incubated overnight at 4°C in sterile distilled water, after which the eluted DNA was used as template to re-amplify the specific fragments using identical PCR conditions and primers as above. After confirming that the re-amplified fragments co-migrated with the expected 16S rRNA V3 PCR fragments of the original samples on DGGE gels, the re-amplified fragments were subjected to another round of PCR using primers 518R and 357F to remove the GC-clamp. The resulting products were then purified with the QIAquick PCR purification kit (QIAGEN, Germany) and sequenced in both directions using primers 518R and 357F, the BigDye[®] Terminator v. 3.1 Cycle Sequencing kit (Applied Biosystems, USA) and an ABI PRISM[™] 3100 DNA sequencer (Applied Biosystems).

3.2.3 Enumeration, isolation and characterization of bacteria

For the fermented sorghum flour slurries, triplicate 10-fold serial dilutions were carried out by transferring 10-g samples to 90 ml of buffered peptone water (0.1% [w/v] peptone, 0.85% [w/v] NaCl; pH 7.2), followed by vortexing for 3 min to obtain homogenous mixtures. One (1) ml of this 10⁻¹ diluted homogenate was then transferred into 9 ml of the same diluent. Aliquots (100 µl) of the dilutions were surface inoculated onto the following selective media: MRS-5 (Oxoid, Basingstoke, UK) for lactobacilli (Meroth *et al.*, 2003), M17 for lactococci (Terzaghi and Sandine, 1975), *Enterococcus* selective agar (ESA) for enterococci (Slanetz and Bartley, 1957), Violet red bile agar (VRBA, Oxoid) to obtain Gram-negative counts

(GNC) (Mossel *et al.*, 1986) and plate count agar (PCA, Oxoid) to obtain numbers of total aerobic bacteria (Reasoner and Geldreich, 1985). The plates were incubated at 30°C for 24-48 h for MRS-5, M17 and ESA, 37°C for 24-48 h for VRBA, and 30°C for 72 h for PCA. For total aerobic counts and LAB counts, plates with colonies between 30 and 300 were selected for enumeration, whilst colonies between 15 and 150, and greater than 0.5 mm in diameter, were selected for GNC on VRBA. The counts were recorded as averages of three determinations and they were expressed as colony forming units (cfu) per gram.

After incubation, colonies on the various growth media were grouped according to macroscopic and microscopic appearance. Representative colonies were then randomly picked and sub-cultured to obtain pure cultures, after which they were routinely grown in appropriate broth media. *Lactobacillus*, *Weissella* and *Lactococcus* species were grown in MRS broth, whilst *Enterobacteriaceae* were grown in nutrient broth using the incubation conditions described. All pure cultures were microscopically examined using light microscopy to score cell morphology, motility and Gram stain. Catalase and oxidase activity were evaluated using 3% hydrogen peroxide (Morgulis, 1921) and tetramethyl-*p*-phenylenediamine (TMPD), respectively (Ehrlich, 1956). All cultures were stored at -20°C in sterile Eppendorf tubes, containing the appropriate broth media supplemented with 20% (v/v) glycerol as cryoprotectant.

3.2.4 PCR and sequencing of the 16S rRNA and *pheS* genes from pure cultures

DNA was extracted from pure cultures using the Wizard[®] Genomic DNA Purification Kit (Promega, Madison, USA) or the method described by Pitcher *et al.* (1989). The *pheS* gene was amplified for each of the isolated pure cultures using the primers PheS-21-F (5'-cayccngchcgycgayatgc-3') and PheS-23-R (5'-gggtgraccatvccngchcc-3'), as described by Naser *et al.* (2005). Amplicons were purified using the Nucleofast 96 PCR clean-up membrane system (Machery-Nagel, Germany) and sequenced, as described above, using the original *pheS* PCR primers.

The gene encoding 16S rRNA was amplified for each of the isolated pure cultures using the eubacterial universal primers 27F (5'-agagtttgatcctggctcag-3'; Lane, 1991) and 1507R (5'-tacctgttacgacttcaccca-3'; Heyndrickx *et al.*, 1996). PCR mixtures contained 1.25 U of *Taq* DNA polymerase (Fermentas, St. Leon-Rot, Germany) and PCR buffer containing NH₂SO₄

(Fermentas), 1.0 mM MgCl₂, 0.2 mM of each dNTP, 0.4 μM of each primer, 50 ng of template DNA and 8% (v/v) dimethyl sulphoxide (DMSO). The PCR cycling conditions consisted of an initial denaturation step at 95°C for 10 min, and 35 cycles with denaturation at 94°C for 1 min, primer annealing at 51°C for 30 s and primer extension at 72°C for 1 min, followed by final extension at 72°C for 10 min and cooling to 4°C. The resulting 16S rRNA PCR products were sequenced with the original primers by Inqaba Biotech (Pretoria, South Africa).

3.2.5 Sequence and phylogenetic analyses

All raw sequence files were inspected and corrected, where necessary, using Chromas Lite 2.0 (Technelysium) and BioEdit v. 5.0.9 (Hall, 1999). To match the sequences obtained for individual 16S rRNA V3 fragments excised from DGGE gels with those obtained for the complete 16S rRNA gene of the pure cultures, these sequences were compared in BioEdit. All sequences were also compared to those in the nucleotide database of the National Centre for Biotechnology Information (NCBI; <http://www.ncbi.nlm.nih.gov/>) using *blastn* (Altschul *et al.*, 1990) to obtain preliminary identifications for the isolated bacteria.

Alignments for the complete 16S rRNA gene and *pheS* data determined in this study were generated using Multiple sequence Alignment based on Fast Fourier Transform (MAFFT v. 6, Katoh *et al.*, 2002; 2005). These alignments also included the sequence information for the relevant type strains of species in the genera *Lactobacillus*, *Weissella*, *Lactococcus* and *Enterococcus*, which were obtained from GenBank. To determine the best-fit evolutionary models for the datasets, PAUP* v. 4.0b1 (Swofford, 2002), together with Modeltest v. 3.7 (Posada and Crandall, 1998), were used. The calculated parameters were then used to construct Neighbor-Joining (NJ) distance-based (Saitou and Nei, 1987) phylogenies with PAUP*, and maximum likelihood (ML) phylogenies with PhyML v. 2.4.3 (Guindon and Gascuel, 2003). Branch support for the various NJ and ML trees were determined using non-parametric bootstrap analysis based on 1000 replicates and the same parameters as before.

3.3 RESULTS

3.3.1 Enumeration and presumptive identification of bacteria

In order to select representative isolates that were dominant at the beginning and end of the fermentation of both sorghum samples at 25°C, enumeration of total bacteria and specific

groups of microorganisms were carried out by cultivation on five different types of media (PCA, MRS-5, M17, VRBA and ESA). Enumerations of total bacteria at 20°C and 30°C were not performed, as PCR-DGGE profiles were similar for fermentations performed at 20, 25 and 30°C. Total counts were initially low, but increased considerably toward the end of the fermentation (Table 3.1). Although LAB were predominant in most media (except VRBA), none of the media were sufficiently specific to allow growth of only particular LAB species. For example, both Gram-positive cocci and rods were isolated indiscriminately from MRS-5, ESA and M17 media.

A total of 192 isolates from both commercial sorghum sources and the various media were randomly selected. Of these, all Gram-positive, catalase-negative, non-motile and oxidase-negative isolates were considered presumptive LAB, whilst all Gram-negative, catalase-positive and oxidase-negative rods were presumptive *Enterobacteriaceae*. A total of 32 representative isolates were selected for further characterization (Table 3.2).

3.3.2 PCR-DGGE analyses

The diversity and dynamics of microflora during spontaneous fermentation of sorghum flour were studied using PCR-DGGE of the ca. 250-bp V3 hypervariable region of the 16S rRNA gene. Based on the PCR-DGGE profiles and pH, the overall fermentation process appeared to be divided into two stages (Fig. 3.1). The first stage (0-6 h) was associated with a relatively small change in pH (6.64-6.0) and simple DGGE profiles, mainly consisting of one intense band that was already present at $t = 0$ h. The second stage (12-54 h) was characterized by a more complex pattern, including 12 different fragments that remained almost the same until the end of fermentation. From 12-48 h of the second stage, the pH dropped from 5.69 to 3.79, after which it remained constant. Similar results were obtained for both of the commercial sorghum flour brands included in this study. All duplicate samples also generated identical PCR-DGGE profiles during the course of spontaneous fermentation, emphasizing the reproducibility of PCR-DGGE. The PCR-DGGE profiles for fermentations at 20, 25 and 30°C were also similar.

Table 3.1 Microbial counts (cfu/g) at the start and end of spontaneous sorghum fermentations (25°C) obtained on different selective media

Media	t = 0 h (K.F.C [*])	t =54 h (K.F.C)	t = 0 h (Nola [*])	t = 54 h (Nola)
VRBA	8.45×10^3	9.0×10^6	8.3×10^3	8.8×10^6
ESA	2×10^3	3.62×10^8	1.8×10^3	3.5×10^8
MRS-5	9×10^3	3.91×10^9	9.1×10^3	3.89×10^9
M17	3×10^3	2.8×10^9	2.8×10^3	2.92×10^9
PCA	1.9×10^5	3.9×10^9	1.5×10^5	2.3×10^9

^{*} K.F.C and Nola refer to commercial sorghum flour from King Food Corporation and Nola Pvt Ltd.

Table 3.2 Likely species identities of bacteria isolated at the beginning and end of sorghum fermentations

Sample identification ¹	Sorghum source ²	Sampling time (h)	DNA-based identification ³
M027M	K.F.C	0	<i>E. mundtii</i>
M033N	Nola	0	<i>E. mundtii</i>
M0331M	K.F.C	0	<i>E. mundtii</i>
107N	Nola	0	<i>E. mundtii</i>
M027N	Nola	0	<i>E. mundtii</i>
M030M	K.F.C	0	Novel, related to <i>E. mundtii</i>
106	K.F.C	0	<i>E. mundtii</i>
1052	Nola	0	<i>E. mundtii</i>
1053	K.F.C	0	<i>E. mundtii</i>
P11	K.F.C	54	<i>Lb. plantarum</i>
P3	K.F.C	54	<i>Lb. rhamnosus</i>
P31	Nola	54	<i>Lb. rhamnosus</i>
L235N	Nola	54	<i>Lb. rhamnosus</i>
P2M	K.F.C	54	<i>Lb. fermentum</i>
M5444M	K.F.C	54	<i>W. cibaria</i>
M5444N	Nola	54	<i>W. cibaria</i>
E548M	K.F.C	54	<i>E. faecalis</i>
E548N	Nola	54	<i>E. faecalis</i>
E5412M	K.F.C	54	<i>E. faecalis</i>
27291	K.F.C	54	<i>Lb. fermentum</i>
27292	Nola	54	<i>Lb. fermentum</i>
27293	Nola	54	<i>Lb. fermentum</i>
27294	Nola	54	<i>Lb. fermentum</i>
P1N	Nola	54	<i>Lb. plantarum</i>
P123r1	K.F.C	54	<i>L. lactis</i>
P123r2	Nola	54	<i>L. lactis</i>
P123r3	Nola	54	<i>L. lactis</i>
105	K.F.C	54	<i>W. cibaria</i>
V5422M	K.F.C	54	<i>Pantoea</i> species
V5423N	Nola	54	<i>Enterobacteriaceae</i>
V5430	Nola	54	<i>Enterobacteriaceae</i>
V5431	K.F.C	54	<i>Enterobacteriaceae</i>

¹ Sample identification refers to the arbitrary identification of bacterial isolates.

² Sorghum source refers to the commercial company that produces the sorghum flour (K.F.C for King Food Corporation and Nola for Nola Pvt Ltd).

³ DNA-based identification refers to the final identification of the isolate obtained from the results of sequence analysis of 16S rRNA and *pheS* genes.





Figure 3.1 Change in pH (A), as well as PCR-DGGE banding patterns (B) representing the 16S rRNA gene fragments of the bacteria involved in the spontaneous fermentation of sorghum at 25°C. Lane M: 16S rRNA V3 marker generated for this study with bands corresponding to *Bacteriodes fragilis* (DSM 1396), *Bacteroides thetaiomicron* (LMG 11262), *Weissella cibaria* (LMG 17699), *Enterococcus flavescens* (LMG 13518T), *Enterococcus solitarius* (LMG 12890T), *Leuconostoc fructosum* (LMG 9498T), *Bacillus subtilis* (LMG 7135T), *Clostridium butyricum* (LMG 1212), *Lactobacillus rhamnosus* (LMG 6400T), *Bifidobacterium longum* (LMG 13197T), *Bifidobacterium bifidum* (LMG 11041T), *Bifidobacterium lactis* (LMG 18314T), and *Bifidobacterium dentium* (LMG 11045T). Sampling time points for PCR-DGGE analysis and pH measurements are indicated at times 0, 6, 12, 18, 24, 30, 36, 42, 48 and 54 h. The identity of the different bacterial species, as suggested by sequence and phylogenetic analyses (Figs. 3.2 and 3.3), are indicated by arrows to the right-hand side of the gel image. Similar PCR-DGGE profiles were generated for fermentations at 25 and 30°C and for both sorghum sources.

To determine the identity of the bacteria represented by the various PCR-DGGE fragments, comparisons of V3 sequences to the full-length 16S rRNA gene sequences obtained from pure cultures and phylogenetic analysis (see below) were used. These analyses also showed that two PCR-DGGE fragments associated with the second fermentation stage represented members of the LAB (i.e. *L. lactis* and *W. cibaria*) and eight fragments represented *Enterobacteriaceae* (i.e. *Pantoea* sp. and *Enterobacter* sp.). The sequence of only one of the PCR-DGGE fragments evident at the end of the fermentation did not match the full-length 16S rRNA gene sequences obtained from any of the pure cultures, and based on the *blastn* results it most likely represents *Lb. curvatus*. The single band associated with the first fermentation stage (0-6 h) most likely represents the sorghum chloroplast 16S rRNA gene, because its sequence is most similar to that of the model monocotyledonous plant *Zea mays*. The *L. lactis* amplicon appeared at t = 18 h and remained until the end of fermentation in all experiments. Once it appeared the intensity of the *L. lactis* band remained constant or increased throughout the fermentation, while those representing the other bacteria, specifically the *Enterobacteriaceae*, decreased during the overall fermentation process.

3.3.3 Phylogenetic analyses using 16S rRNA and *pheS* gene sequences

Two separate data sets were created for the bacteria examined in this study. One included 16S rRNA gene sequences for putative *Lactobacillus* and *Weissella* species (i.e. the Gram-positive rod-shaped bacteria), while the other included *Lactococcus* and *Enterococcus* species (i.e. the Gram-positive coccoid bacteria). These alignments also included the full-length or near full-length sequences for the type strains of the species in these genera that were obtained from GenBank. For each of the datasets, similar trees were inferred using NJ and ML methods (Figs. 3.2 and 3.3). Within the *Lactobacillus* and *Weissella* phylogeny (Fig. 3.2), isolates P11 and P1N formed part of a clade containing *Lb. plantarum*, *Lb. paraplantarum*, *Lb. pentosus* and *Lb. arizonensis*, while P31, L235N and P3 were most closely related to *Lb. rhamnosus* and *Lb. casei*. However, phylogenetic analysis of *pheS* sequences clearly showed that isolates P11 and P1N represented *Lb. plantarum*, while P31, L235N and P3 represented *Lb. rhamnosus*. Isolates 27291, 27292, 27293, 27294 and P2 grouped together and represented *Lb. fermentum*, and isolates M5444N, M5444N and 105 were most closely related to *W. cibaria* (Fig. 3.2A). Within the *Lactococcus* and *Enterococcus* phylogeny (Fig. 3.3), isolates P123r1, P123r2, P123r3 represented *L. lactis*, and isolates M027N, M027M, M0331M, M0331N, 107N, 106 and 1052 represented *E. mundtii*. The isolates E548N, E5412M and E548M clustered with *E. faecalis* (Fig. 3.3).

(A)



(B)




Figure 3.2 Maximum likelihood (ML) phylogeny of species in the genera *Lactobacillus* and *Weissella*, based on 16S rRNA (A) and *pheS* (B) gene sequences. Trees with similar topologies were inferred using Neighbor-Joining (NJ) distance analysis. Isolates obtained from fermented sorghum are indicated in bold. All species names are preceded by strain numbers and GenBank accession numbers in brackets. *Bacillus subtilis* (A) and *Enterococcus faecalis* (B) were used as the outgroups and bootstrap values (> 60%), based on 1000 replications, are shown at internodes, as follows: ML/NJ. *Weissella kimchii* is a later heterotypic synonym of *Weissella cibaria* (Ennahar and Cai, 2004). *Lb. arizonensis* is a later heterotypic synonym of *Lb. plantarum* (Kostinek *et al.*, 2005).




Figure 3.3 Maximum likelihood (ML) phylogeny of species in the genera *Lactococcus* and *Enterococcus*, based on 16S rRNA gene sequences. Trees with similar topologies were inferred using Neighbor-Joining (NJ) distance analysis. Isolates obtained from fermented sorghum are indicated in bold. All species names are preceded by strain numbers and GenBank accession numbers in brackets. *Bacillus subtilis* was used as the outgroup and bootstrap values (> 60%), based on 1000 replications, are shown at internodes, as follows: ML/NJ.

3.3.4 Detection of bacterial diversity using culture-independent and culture-dependent approaches

Table 3.3 shows a comparison of the diversity of bacteria identified during spontaneous sorghum fermentations using culture-independent and culture-dependent approaches. Overall, PCR-DGGE appeared to be unable to detect most of the bacteria in the *ting* samples. For example, at time $t = 0$ h, *E. mundtii* was found to be present using culture-dependent methods (Table 3.2), although PCR-DGGE did not allow its detection, as all profiles at time $t = 0$ h for both sorghum brands at 20, 25 and 30°C included only the sorghum chloroplast 16S rRNA gene fragment (Fig. 3.1). Also, culture-dependent approaches allowed isolation and identification of *Lb. plantarum*, *Lb. rhamnosus* and *Lb. fermentum* at time $t = 54$ h, although their 16S rRNA gene fragments were absent from the PCR-DGGE profiles at comparable time points in both sorghum types at 20, 25 and 30°C (Table 3.3). However, an isolate presumably representing *Lb. curvatus* and that was predominant in PCR-DGGE profiles at time points $t = 42-54$ h was not isolated from the fermented samples, despite numerous attempts.

3.4 DISCUSSION

The use of culture-independent PCR-DGGE, in combination with culture-dependent microbiological methods, allowed for the effective description of the diversity and dynamics of the bacterial populations involved in the spontaneous fermentation process to produce *ting*. The results illustrated that various LAB and members of the *Enterobacteriaceae* were predominant at the end of fermentation, with *L. lactis* appearing to play a more important role compared to other bacteria. Consequently, *L. lactis* may be used to control sorghum fermentations, resulting in standardized products. The research presented here represents the first extensive study investigating bacterial diversity and dynamics in this ecosystem, and is relevant for development of appropriate and effective starter cultures.

In this study, species-level identification was primarily based on 16S rRNA gene sequences. This approach allowed positive identification of most of the bacteria involved in sorghum fermentations as *E. mundtii*, *E. faecalis*, *L. lactis*, and *Lb. fermentum* or *W. cibaria*. It did, however, not allow unambiguous identification of isolates that are related to *Lb. plantarum*, *Lb. paraplantarum*, *Lb. pentosus* and *Lb. arizonensis*, nor those related to *Lb. rhamnosus* and *Lb. casei*. The bacteria in these two groups are each also characterized by very similar

Table 3.3 Comparison of the bacterial diversity in *ting* using culture-independent and culture-dependent approaches

Time of detection	Culture-independent approach ¹	Culture-dependent approach ²
t = 0 h	None	<i>E. mundtii</i>
t = 54 h	<i>L. lactis</i> <i>W. cibaria</i> <i>Enterobacteriaceae</i> members (4) <i>Lb. curvatus</i>	<i>L. lactis</i> <i>W. cibaria</i> <i>Enterobacteriaceae</i> members (4) <i>E. faecalis</i> <i>Lb. plantarum</i> <i>Lb. fermentum</i> <i>Lb. rhamnosus</i>

¹ Identification was based on sequencing of the V3 region of the 16S rRNA gene.

² Identification was based on results of sequence analysis of the 16S rRNA and *pheS* genes.

phenotypes, further complicating their differentiation (Torriani *et al.*, 2001; Ennahar *et al.*, 2003). The fact that 16S rRNA gene sequence analysis did not allow their definitive identification is, however, not surprising as it has been shown on numerous occasions that this gene region is unable to separate closely related species (Fox *et al.*, 1992; Ward, 2002; Felis and Dellaglio, 2007). Recently, Naser *et al.* (2007) introduced the use of additional genes such as *pheS* and *rpoB* for separating species of *Lactobacillus*. In this study, it was similarly found that *pheS* has greater resolution, because it displayed sufficient interspecific variation that allowed for the unequivocal identification of isolates P3, P31 and L235N as *Lb. rhamnosus*, and isolates P1N and P11 as *Lb. plantarum* (Fig. 3.2B).

Spontaneous sorghum fermentations were characterized by two well-delineated stages that are linked to bacterial diversity and pH. There was an apparent shift in microbial diversity from simple PCR-DGGE profiles and relatively high pH to more complex PCR-DGGE profiles and increased acidity (Fig. 3.1). During the first phase of fermentation the bacterial counts were very low ($< 8.5 \times 10^3$ cfu/g, Table 3.1) and below the detection limit of PCR-DGGE. The single intense PCR-DGGE band dominating the first fermentation phase represented the V3 region of the 16S rRNA gene encoded on the sorghum chloroplast, which is also a eubacterial descendent (Krajčovič and Ebringer, 1990; Martin, 1999). During the second fermentation phase, PCR-DGGE detected *L. lactis*, which remained dominant until the end of the fermentation. *W. cibaria* and the putatively identified *Lb. curvatus* were also among the predominant species at the end of the fermentations. The dominance of *L. lactis*, a homofermenter, over other bacteria may be due to its tolerance of the low pH, which reduced the growth of heterofermentative LAB (Lin *et al.*, 1992). Therefore, *L. lactis* together with the other dominant species represent important bacteria that are well adapted to sorghum fermentation processes and may be suitable candidates to use as starter cultures for standardization of *ting* preparations. These results are also in agreement with the generally accepted concept that traditional fermentations are dominated by a few microbial species that are selected during the course of fermentation because of good adaptation to the food matrix (Hounhouigan *et al.*, 1993; Halm *et al.*, 1996).

All of the LAB species associated with the production of *ting* have previously been implicated in the production of fermented cereals with desirable sensory properties. *Lb. plantarum* was isolated from fermented cereals such as *ogi* (Odunfa and Adyele, 1985), *mageu* (Hesseltine, 1979) and *kunun zaki* (Gaffa and Gaffa, 2004), whilst the heterofermentative *Lb. fermentum*

was previously isolated from *mawe* (Hounhouigan *et al.*, 1993) and *kenkey* (Halm *et al.*, 1993). *Lb. curvatus-Lb. sake* was isolated from sorghum-based fermented weaning food (Kunene *et al.*, 2000). The production of the pleasant sensory characteristics of *ting* is likely due to some LAB such as *Lb. plantarum*, which has previously been shown to produce flavour-active compounds such as ethyl acetate and aldehydes (Rehman *et al.*, 2006).

In this study, a number of bacteria not normally associated with food were isolated from spontaneously fermented sorghum. The majority of these are naturally associated with plants and commonly found on plant-based material (Nout, 1991; Mohammed *et al.*, 1991). The yellow-pigmented species resembling *E. mundtii* that was isolated at the beginning of the process is typically associated with plants (Martin and Mundt, 1972) where it exhibits epiphytic relationships with them (Mundt *et al.*, 1962), and has also been reported as an environmental contaminant (Camu *et al.*, 2007). The same is also true for the *Pantoea* sp., which is a plant endophyte and commonly associated with rice and soybean (Kuklinsky-Sobral *et al.*, 2004). However, some *Pantoea* species are associated with diseases in humans such as septic monoarthritis (De Champs *et al.*, 2000). The presence of a bacterial species representing *E. faecalis* at the end of sorghum fermentations was expected, as this species is often prevalent in foods (Giraffa, 2002; Gomes *et al.*, 2008). Enterococci have been shown to form part of the microflora that is present throughout the cheese production process, and are responsible for sensory characteristics of the final product (Franz *et al.*, 2003). In addition, some enterococci strains, especially *E. faecalis* and *E. faecium* may produce bacteriocins that are active against a plethora of food borne pathogens (Franz *et al.*, 2007), making them suitable candidates for controlling emerging pathogens during food fermentation (Callewaert *et al.*, 2000). Despite the safety and pleasant sensory attributes imparted by *E. faecalis* in foods, some strains of *E. faecalis* and *E. faecium* are associated with infection that pose challenges to food safety (Carlos *et al.*, 2009).

One of the broadly recognized advantages of subjecting food materials to LAB-steered fermentation processes is the inhibitory effect this has on the growth of other microorganisms, especially food-borne pathogens (Nigatu and Gashe, 1994; Kingamkono *et al.*, 1995). Many previous studies have shown that the growth of members of the *Enterobacteriaceae* and other bacteria is inhibited by the lactic acid produced during fermentation (Nout, 1991; Adams and Nicolaides, 1997; Abegaz, 2007). The presence of bacteria representing *Pantoea* sp. and *E. faecalis* during and at the end of the sorghum fermentations might, however, be due to acid

resistance or the presence of microenvironments in the food matrix that support the growth of these bacteria (Wacher *et al.*, 1993; ben Omar and Ampe, 2000). Similar findings were also reported for other studies where significant numbers of coliforms and *Escherichia coli* were detected during fermentations involving LAB, especially when no starter cultures were added (Feresu and Nyathi, 1990; ben Omar and Ampe, 2000; Gran *et al.*, 2002; 2003). Overall, however, the intensity of PCR-DGGE bands corresponding to *Enterobacteriaceae* decreased as that of *L. lactis* increased. It is therefore likely that the unfavourable environment created by this and the other LAB slowed down the growth of *Pantoea* sp. and *E. faecalis* as fermentation progressed. This was also probably true for *E. mundtii* that was only isolated at the start of the fermentations and not thereafter.

The comparison of culture-independent and culture-dependent approaches used in this study highlights the limitations of each approach. Some species were not detected using culture-based methods, while others were not detected using culture-independent methods. PCR-DGGE showed a significant increase in the putatively identified *Lb. curvatus* numbers from time points $t = 42$ to 54 h of the fermentation process, but were not isolated from the food matrix. This may be due to it entering a viable but non-cultivable state, characterized by metabolically active cells that do not produce colonies on both selective and non-selective media (Giraffa and Neviani, 2001), which illustrates one of the main advantages of culture-independent approaches over culture-dependent methods (Muyzer and Smalla, 1998; Giraffa and Neviani, 2001). Overall, however, the use of culture-dependent methods allowed identification of more bacteria than culture-independent methods (Table 3.3). PCR-DGGE did not allow the detection of *E. faecalis*, *E. mundtii*, *Lb. rhamnosus*, *Lb. plantarum* and *Lb. fermentum* that were isolated on culture media at the end of fermentation. This was probably because the bacteria occurred in numbers below the detection limit of PCR-DGGE (Muyzer and Smalla, 1998). Biases that exist in terms of DNA extraction and PCR (Ercolini, 2004; De Vero *et al.*, 2006; Camu *et al.*, 2007) could also have played a role. The inability of PCR-DGGE to detect all the bacterial species associated with fermented sorghum was also observed for sourdough (Meroth *et al.*, 2003) and whey cultures of water buffalo mozzarella (Ercolini *et al.*, 2004). PCR-DGGE results are also strongly influenced by the intraspecific heterogeneous nature of the 16S rRNA gene region (Ueda *et al.*, 1999), which in the case of *Pantoea* species, yielded multiple bands associated with apparently a single species (Fogel *et al.*, 1999). Accurate and efficient description of bacterial populations during sorghum

fermentations was therefore strongly dependent on the combined application of culture-independent and culture-dependent approaches.

The use of a culture-dependent approach used in a previous study (Chapter 2) also highlights the advantages of using a combination of both culture-dependent and culture-independent approaches applied in this study. Some of the strains isolated from fermented sorghum in the previous study were similar to those obtained in this study (*Lb. rhamnosus*, *Lb. fermentum* and *Lb. plantarum*). However, *L. lactis* that was isolated in this study could not be detected in the previous study, probably because the addition of five nutritional supplements to molten MRS medium (Meroth *et al.*, 2003) improved the chances of isolating this LAB. Likewise, the use of ESA and M17 agar improved the probability of isolating *E. faecalis* that was isolated at the end of fermentation in this study, but could not be detected on MRS in the previous study.

In South Africa, *ting* is still produced from spontaneously fermented sorghum flour. Consequently, there is immense variation in the sensory characteristics and quality of *ting*, making it a daunting task to upgrade its status to commercial level. Detailed insight of the microbial processes occurring during spontaneous sorghum fermentations is a prerequisite for development of starter cultures that may result in the production of standardized *ting*. In this study, microbial population dynamics and diversity during sorghum fermentations were analyzed in detail, using a combination of culture-independent and culture-dependent methods, in order to get an overview of the important bacteria. Taken together, the use of these methods offered an excellent approach for increasing knowledge of microbial diversity and dynamics during spontaneous sorghum fermentations. Such information is useful, not only from an academic viewpoint, but for designing relevant starter cultures that may result in *ting* with standardized sensory profiles (appearance, aroma, sourness and taste) and fermentation time. Evaluation of the contribution of such starter cultures to the safety and acceptability of sensory characteristics of *ting* is important in future.

3.5 REFERENCES

Abegaz, K. (2007). Isolation, characterization and identification of lactic acid bacteria involved in traditional fermentation of *borde*, an Ethiopian cereal beverage. *African Journal of Biotechnology*, **6**, 1469-1478.

Adams, M.R. and Nicolaides, L. (1997). Review of the sensitivity of different pathogens to fermentation. *Food Control*, **8**, 227-239.

Akingbala, J.O., Rooney, L.W. and Faubion, J.M. (1981). A laboratory procedure for the preparation of *ogi*, a Nigerian fermented food. *Journal of Food Science*, **46**, 1523-1526.

Altschul, S.F., Gish, W., Myers, E.W. and Lipman, D.J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, **215**, 403-410.

Ampe, F., ben Omar, N. and Guyot, J.P. (1999). Culture-independent quantification of physiologically active microbial groups in fermented foods using rRNA-targeted oligonucleotide probes: Application to *pozol*, a Mexican lactic acid fermented maize dough. *Journal of Applied Microbiology*, **87**, 131-140.

Au, P.M. and Fields, M.L. (1981). Nutritive quality of fermented sorghum. *Journal of Food Science*, **46**, 652-654.

ben Omar, N. and Ampe, F. (2000). Microbial community dynamics during production of the Mexican fermented maize dough *pozol*. *Applied and Environmental Microbiology*, **66**, 3664-3673.

Boling, M.B. and Eisener, N. (1982). *Bogobe*: Sorghum porridge of Botswana. In: Proceedings of the International Symposium on Sorghum Grain Quality. International Crops Research Institute for the Semi-Arid Tropics. Mertin, J.V. (Ed.). Patancheru, India, pp. 32-35.

Bvochora, J.M., Reed, J.D., Read, J.S. and Zvauya, R. (1999). Effect of fermentation processes on proanthocyanidins in sorghum during preparation of *mahewu*, a non-alcoholic beverage. *Process Biochemistry*, **35**, 21-25.

Callewaert, R., Hugas, M. and De Vuyst, L. (2000). Competitiveness and bacteriocin production of enterococci in the production of Spanish-style dry fermented sausages. *International Journal of Food Microbiology*, **57**, 33-42.

Camu, N., De Winter, T., Verbrugghe, K., Cleenwerck, I., Vandamme, P., Takrama, J.S., Vancanneyt, M. and De Vuyst, L. (2007). Dynamics and biodiversity of populations of lactic acid bacteria involved in spontaneous heap fermentation of cocoa beans in Ghana. *Applied and Environmental Microbiology*, **73**, 1809-1824.

Carlos, A.R., Santos, J., Semedo-Lemsaddek, T., Barreto-Crespo, M.T. and Tenreiro, R. (2009). Enterococci from artisanal dairy products show high levels of adaptability. *International Journal of Food Microbiology*, **129**, 194-199.

Caplice, E. and Fitzgerald, G.F. (1999). Food fermentations: Role of microorganisms in food production and preservation. *International Journal of Food Microbiology*, **50**, 131-149.

Chavan, U.D., Chavan, J.K. and Kadam, S.S. (1988). Effect of fermentation on soluble proteins and *in vitro* protein digestibility of sorghum, green gram and sorghum-green gram blends. *Journal of Food Science*, **53**, 1574-1575.

De Champs, C., Le Seaux, S., Dubost, J.J., Boisgard, S., Sauvezie, B. and Sirot, J. (2000). Isolation of *Pantoea agglomerans* in two cases of septic monoarthritis after plant thorn and wood sliver injuries. *Journal of Clinical Microbiology*, **38**, 460-461.

De Vero, L., Gala, E., Gullo, M., Solieri, L., Landi, S. and Giudici, P. (2006). Application of denaturing gradient gel electrophoresis (DGGE) analysis to evaluate acetic acid bacteria in traditional balsamic vinegar. *Food Microbiology*, **23**, 809-813.

Devriese, L.A. and Pot, B. (1995). The genus *Enterococcus*. In: The genera of lactic acid bacteria. Wood, B.J.B. and Holzapel, W.H. (Eds.). Blackie Academic, London, UK, pp. 327-367.

Ehrlich, P. (1956). The requirement of the organism for oxygen. An analytical study with the aid of dyes. In: Collected papers of Paul Ehrlich. Himmelweit *et al.* (Eds.). Pergamon Press, London, UK, pp. 433-496.

Ennahar, S., Cai, Y. and Fujita, Y. (2003). Phylogenetic diversity of lactic acid bacteria associated with paddy rice silage as determined by 16S ribosomal DNA analysis. *Applied and Environmental Microbiology*, **69**, 444-451.

Ennahar, S. and Cai, Y. (2004). Genetic evidence that *Weissella kimchii* Choi *et al.* 2002 is a later heterotypic synonym of *Weissella cibaria* Björkroth *et al.* 2002. *International Journal of Systematic and Evolutionary Microbiology*, **54**, 463-465.

Ercolini, D. (2004). PCR-DGGE fingerprinting: Novel strategies for detection of microbes in food. *Journal of Microbiological Methods*, **56**, 297-314.

Ercolini, D., Mauriello, G., Blaiotta, G., Moschetti, G. and Coppola, S. (2004). PCR-DGGE fingerprints of microbial succession during a manufacture of traditional water buffalo mozzarella cheese. *Journal of Applied Microbiology*, **96**, 263-270.

Felis, G.E. and Dellaglio, F. (2007). Taxonomy of *Lactobacilli* and *Bifidobacteria*. *Current Issues in Intestinal Microbiology*, **8**, 44-61.

Feresu, S. and Nyathi, H. (1990). Fate of pathogenic and non-pathogenic *Escherichia coli* strains in two fermented milk products. *Journal of Applied Bacteriology*, **69**, 814-821.

Fogel, G.B., Collins, C.R., Li, J. and Brunk, C.F. (1999). Prokaryotic genome size and SSU rDNA copy number: Estimation of microbial relative abundance from a mixed population. *Microbial Ecology*, **38**, 93-113.

Food and Agriculture Organization of the United Nations. 2006. FAOSTAT – FAO Statistical Databases. (<http://faostat.fao.org/>).

Fox, G.E., Wisotzkey, J.D. and Jurtshuk, P. (1992). How close is close: 16S rRNA sequence identity may not be sufficient to guarantee species identity. *International Journal of Systematic Bacteriology*, **42**, 166-170.

Franz, C.M., Stiles, M.E., Schleifer, K.H. and Holzapfel, W.H. (2003). Enterococci in foods - a conundrum for food safety. *International Journal of Food Microbiology*, **88**, 105-122.

Franz, C.M.A.P., van Belkum, M.J., Holzapfel, W.H., Abriouel, H. and Gálvez, A. (2007). Diversity of enterococcal bacteriocins and their grouping into a new classification scheme. *FEMS Microbiology Reviews*, **31**, 293-310.

Gaffa, T. and Gaffa, A.T. (2004). Microbial succession during ‘*kunun zaki*’ production with sorghum (*Sorghum bicolor*) grains. *World Journal of Microbiology and Biotechnology*, **20**, 449-453.

Gebrekidan, B. and Gebrettiwat, B. (1982). Sorghum *injera*: Preparation and quality parameters. In: Proceedings of the International Symposium on Sorghum Grain Quality. International Crops Research Institute for the Semi-Arid Tropics. Mertin, J.V. (Ed.). Patancheru, India, pp. 55-56.

Giraffa, G. and Neviani, E. (2001). DNA-based, culture-independent strategies for evaluating microbial communities in food-associated ecosystems. *International Journal of Food Microbiology*, **67**, 19-34.

Giraffa, G. (2002). Enterococci from foods. *FEMS Microbiology Reviews*, **26**, 163–171.

Gomes, B.C., Esteves, C.T., Palazzo, I.C.V., Darini, A.L.C., Felis, G.E., Sechi, L.A., Franco, B.D.G.M. and De Martinis, E.C.P. (2008). Prevalence and characterization of *Enterococcus* spp. isolated from Brazilian foods. *Food Microbiology*, **25**, 668-675.

Guindon, S. and Gascuel, O. (2003). A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology*, **52**, 696-704.

Gran, H.M., Wetlesen, A., Mutukumira, A.N. and Narvhus, J. (2002). Smallholder dairy processing in Zimbabwe: Hygienic practices during milking and the microbiological quality of the milk at the farm and on delivery. *Food Control*, **13**, 41-47.

Gran, H.M., Wetlesen, A., Mutukumira, A.N., Rukure, G. and Narvhus, J.A. (2003). Occurrence of pathogenic bacteria in raw milk, cultured pasteurized milk and naturally soured raw milk produced at small-scale dairies in Zimbabwe. *Food Control*, **14**, 539-544.

Hall, T.A. (1999). BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, **41**, 95-98.

Halm, M., Sørensen, L.A. and Jakobsen, M. (1993). Microbiological and aromatic characteristics of fermented maize dough from *kenkey* production in Ghana. *International Journal of Food Microbiology*, **19**, 135-143.

Halm, M., Osei-Yaw, A., Hayford, A., Kpodo, K.A. and Amoa-Awua, W.K.A. (1996). Experiences with the use of starter culture in the fermentation of maize for '*kenkey*' production in Ghana. *World Journal of Microbiology and Biotechnology*, **12**, 531-536.

Hamacker, B.R., Kirleis, A.W., Mertz, E.T. and Axtell, J.D. (1986). Effect of cooking on the protein profiles and *in vitro* protein digestibility of sorghum and maize. *Journal of Agricultural and Food Chemistry*, **34**, 647-649.

Hassan, I.A.G. and El Tinay, A.H. (1995). Effect of fermentation on tannin content and *in vitro* protein and starch digestibilities of two sorghum cultivars. *Food Chemistry*, **53**, 149-151.

Heyndrickx, M., Väterin, L., Vandamme, P., Kersters, K. and De Vos, P. (1996). Applicability of combined amplified ribosomal DNA restriction analysis (ARDRA) patterns in bacterial phylogeny and taxonomy. *Journal of Microbiological Methods*, **26**, 247-259.

Hesseltine, C.W. (1979). Some important fermented foods in mid-Asia, the Middle-East and Africa. *Journal of the American Oil Chemists' Society*, **56**, 367-374.

Hounhouigan, D.J., Nout, M.J.R., Nago, C.M., Houben, J.H. and Rombouts, F.M. (1993). Characterization and frequency distribution of species of lactic acid bacteria involved in the processing of *mawe*, a fermented maize dough from Bénin. *International Journal of Food Microbiology*, **18**, 279-287.

Ibrahim, F.S., Babiker, E.E., Yousif, N.E. and El Tinay, A.H. (2005). Effect of fermentation on biochemical and sensory characteristics of sorghum flour supplemented with whey protein. *Food Chemistry*, **92**, 285-292.

Katoh, K., Misawa, K., Kuma, K. and Miyata, T.H. (2002). MAFFT: A novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research*, **30**, 3059-3066.

Katoh, K., Kuma, T.H. and Miyata, T. (2005). MAFFT version 5: Improvement in accuracy of multiple sequence alignment. *Nucleic Acids Research*, **33**, 511-518.

Kazanas, N. and Fields, M.L. (1981). Nutritional improvement of sorghum by fermentation. *Journal of Food Science*, **46**, 819-821.

Kingamkono, R.R., Sjögren, E., Svanberg, U. and Kaijser, B. (1995). Inhibition of different strains of enteropathogens in a lactic-fermenting cereal gruel. *World Journal of Microbiology and Biotechnology*, **11**, 299-303.

Kostinek, M., Pukall, R., Rooney, A.P., Schillinger, U., Hertel, C., Holzapfel, W.H. and Franz, C.M.A.P. (2005). *Lactobacillus arizonensis* is a later heterotypic synonym of *Lactobacillus plantarum*. *International Journal of Systematic and Evolutionary Microbiology*, **55**, 2485-2489.

Kuklinsky-Sobral, J., Araujo, W.L., Mendes, R., Geraldi, I.O., Pizzirani-Kleiner, A.A. and Azevedo, J.L. (2004). Isolation and characterization of soybean-associated bacteria and their potential for plant growth promotion. *Environmental Microbiology*, **6**, 1244-1251.

Kunene, N.F., Hastings, J.E. and von Holy, A. (1999). Bacterial populations associated with a sorghum-based fermented weaning cereal. *International Journal of Food Microbiology*, **49**, 75-83.

Kunene, N.F., Geornaras, I., von Holy, A. and Hastings, J.E. (2000). Characterization and determination of origin of lactic acid bacteria from a sorghum-based fermented weaning food by analysis of soluble proteins and amplified fragment length polymorphism fingerprinting. *Applied and Environmental Microbiology*, **66**, 1084-1092.

Krajčovič, J. and Ebringer, L. (1990). Different effects of eubacterial and eukaryotic DNA topoisomerase II inhibitors on chloroplasts of *Euglena gracilis*. *Origins of Life and Evolution of Biospheres*, **20**, 177-180.

Lane, D.L. (1991). 16S/23S rRNA sequencing. In: *Nucleic acid techniques in bacterial systematics*. Stackebrandt, E. and Goodfellow, M. (Eds.). John Wiley and Sons, New York, USA, pp. 115-175.

- Lin, C., Bolsen, K.K., Brent, B.E. and Fung, D.Y.C.** (1992). Epiphytic lactic acid bacteria succession during the pre-ensiling and ensiling periods of alfalfa and maize. *Journal of Applied Bacteriology*, **73**, 375-387.
- Liu, W.T., Marsh, T.L., Cheng, H. and Forney, L.J.** (1997). Characterization of microbial diversity by determining terminal-restriction fragment length polymorphisms of genes encoding 16S rRNA. *Applied and Environmental Microbiology*, **63**, 4516-4522.
- Lorri, W. and Svanberg, U.** (1995). An overview of the use of fermented foods for child feeding in Tanzania. *Ecology of Food and Nutrition*, **34**, 65-81.
- Martin, W.** (1999). A briefly argued case that mitochondria and plastids are descendants of endosymbionts, but that the nuclear compartment is not. *Proceedings of the Royal Society of London B*, **266**, 1387-1395.
- Martin, J.D. and Mundt, J.O.** (1972). Enterococci in insects. *Applied Microbiology*, **24**, 575-580.
- Meroth, C.B., Walter, J., Hertel, C., Brandt, M.J. and Hammes, W.P.** (2003). Monitoring the bacterial population dynamics in sourdough fermentation processes by using PCR-denaturing gradient gel electrophoresis. *Applied and Environmental Microbiology*, **69**, 475-482.
- Miambi, E., Guyot, J.P. and Ampe, F.** (2003). Identification, isolation and quantification of representative bacteria from fermented cassava dough using an integrated approach of culture-dependent and culture-independent methods. *International Journal of Food Microbiology*, **82**, 111-120.
- Mbugua, S.K.** (1984). Isolation and characterization of lactic acid bacteria during the traditional fermentation of *uji*. *East African Agriculture and Forestry Journal*, **50**, 36-43.
- Mohammed, S.I., Steenson, L.R. and Kirleis, A.W.** (1991). Isolation and characterization of microorganisms associated with the traditional sorghum fermentation for production of Sudanese *kisra*. *Applied and Environmental Microbiology*, **57**, 2529-2533.
- Morrison, D., Woodford, N. and Cookson, B.** (1997). Enterococci as emerging pathogens of humans. *Journal of Applied Microbiology Symposium Supplement*, **83**, 89S -99S.
- Mossel, D.A.A., van der Zee, H., Hardon, A.P. and van Netten, P.** (1986). The enumeration of thermotrophic types amongst the *Enterobacteriaceae* colonizing perishable foods. *Journal of Applied Bacteriology*, **60**, 289-295.
- Morgulis, S.** (1921). A study of the catalase reaction. *Journal of Biological Chemistry*, **47**, 341-375.

- Mukuru, S.** (1992). Traditional technologies in small grain processing. In: Utilisation of sorghum and millets. International Crops Research Institute for Semi-Arid Tropics. Gomez *et al.* (Eds). Patancheru, India, pp. 47-56.
- Mundt, J.O., Coggins, J.A. and Johnson, L.F.** (1962). Growth of *Streptococcus faecalis* var. *liquifaciens* on plants. Applied Microbiology, **10**, 552-555.
- Murray, B.E.** (1990). The life and times of the *Enterococcus*. Clinical Microbiology Reviews, **3**, 46-65.
- Muyanja, C.M.B.K., Narvhus, J.A., Treimo, J. and Langsrud, T.** (2003). Isolation, characterization and identification of lactic acid bacteria from *bushera*: A Ugandan traditional fermented beverage. International Journal of Food Microbiology, **80**, 201-210.
- Muyzer, G. and Smalla, K.** (1998). Application of denaturing gradient gel electrophoresis (DGGE) and temperature gradient gel electrophoresis (TGGE) in microbial ecology. Antonie van Leeuwenhoek, **73**, 127-141.
- Muyzer, G., De Waal, E.C. and Uitterlinden, A.G.** (1993). Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. Applied and Environmental Microbiology, **59**, 695-700.
- Naser, S.M., Thompson, F.L., Hoste, B., Gevers, D., Dawyndt, P., Vancanneyt, M. and Swings, J.** (2005). Application of multilocus sequence analysis (MLSA) for rapid identification of *Enterococcus* species based on *rpoA* and *pheS* genes. Microbiology, **151**, 2141-2150.
- Naser, S.M., Dawyndt, P., Hoste, B., Gevers, D., Vandemeulebroecke, K., Cleenwerk, I., Vancanneyt, M. and Swings, J.** (2007). Identification of lactobacilli by *pheS* and *rpoA* gene sequence analyses. International Journal of Systematic and Evolutionary Microbiology, **57**, 2777-2789.
- Neucere, N.J. and Sumrell, G.** (1979). Protein fractions from varieties of grain sorghum: Amino acid composition and solubility properties. Journal of Agricultural and Food Chemistry, **27**, 809-812.
- Nielsen, D.S., Teniola, O.D., Ban-Koffi, L., Owusu, M., Andersson, T.S. and Holzapfel, W.H.** (2007). The microbiology of Ghanaian cocoa fermentations analysed using culture-dependent and culture-independent methods. International Journal of Food Microbiology, **114**, 168-186.
- Nigatu, A. and Gashe, B.A.** (1994). Inhibition of spoilage and food-borne pathogens by lactic acid bacteria isolated from fermenting *tef* (*Eragrostis tef*) dough. Ethiopian Medical Journal, **32**, 223-229.

- Nout, M.J.R.** (1991). Ecology of accelerated natural lactic fermentation of sorghum-based infant-food formulas. *International Journal of Food Microbiology*, **12**, 217-224.
- Odunfa, S.A. and Adyele, S.** (1985). Microbiological changes during the production of *ogi-baba*, a West-African fermented sorghum gruel. *Journal of Cereal Science*, **3**, 175-180.
- Orita, M., Iwahana, H., Knazawa, H., Hayashi, K. and Sekiya, T.** (1989). Detection of polymorphisms of human DNA by gel electrophoresis as single strand conformation polymorphisms. *Proceedings of the National Academy of Sciences of the United States of America*, **86**, 2766-2770.
- Pitcher, D.G., Saunders, N.A. and Owen, R.J.** (1989). Rapid extraction of bacterial genomic DNA with guanidium thiocyanate. *Letters in Applied Microbiology*, **8**, 151-156.
- Pontes, D.S., Lima-Bittecourt, C.I., Chartone-Souza, E. and Nascimento, A.M.A.** (2007). Molecular approaches: Advantages and artifacts in assessing bacterial diversity. *Journal of Industrial Microbiology and Biotechnology*, **34**, 463-473.
- Posada, D. and Crandall, K.A.** (1998). Modeltest: Testing the model of DNA substitution. *Bioinformatics*, **14**, 817-818.
- Rantsiou, K. and Cocolin, L.** (2006). New developments in the study of the microbiota of naturally fermented sausages as determined by molecular methods: A review. *International Journal of Food Microbiology*, **108**, 255-267.
- Reasoner, D.J. and Geldreich, E.E.** (1985). A new medium for the enumeration and sub-culture of bacteria from potable water. *Applied and Environmental Microbiology*, **49**, 1-7.
- Rehman, S., Paterson, A. and Pigott, J.R.** (2006). Flavour in sourdough breads: A review. *Trends in Food Science and Technology*, **17**, 557-566.
- Renouf, V., Claisse, O., Miot-Sertier, C. and Lonvaud-Funel, A.** (2006). Lactic acid bacteria evolution during wine-making: Use of *rpoB* gene as target for PCR-DGGE analysis. *Food Microbiology*, **23**, 136-145.
- Ritchie, N.J., Schutter, M.E., Dick, R.P. and Myrold, D.D.** (2000). Use of length heterogeneity-PCR and fatty acid methyl ester profiles to characterize microbial communities in soil. *Applied and Environmental Microbiology*, **66**, 1668-1675.

Ruoff, K.L. (1990). Recent taxonomic changes in the genus *Enterococcus*. *European Journal of Clinical Microbiology and Infectious Disease*, **9**, 75-79.

Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989). *Molecular cloning: A laboratory manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, USA.

Sanni, A.I. (1993). The need for process optimization of African fermented foods and beverages. *International Journal of Food Microbiology*, **18**, 85-95.

Saitou, N. and Nei, M. (1987). Reconstructing phylogenetic trees. *Molecular Biology and Evolution*, **4**, 406-425.

Slanetz, L.W. and Bartley, C.H. (1957). Numbers of enterococci in water, sewage, and feces determined by the membrane filter technique with an improved medium. *Journal of Bacteriology*, **74**, 591-595.

Swofford, D.L. (2002). PAUP*. *Phylogenetic Analysis Using Parsimony (* and other methods)*. Version 4. Sinauer Associates, Sunderland, Massachusetts.

Tamura, K. and Nei, M. (1993). Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution*, **10**, 512-526.

Taylor, J.R.N. (2003). Overview: Importance of sorghum in Africa. (<http://www.Afripro.org.uk/papers/Paper01Taylor.pdf> cited 20/05/03).

Terzaghi, B.E. and Sandine, W.E. (1975). Improved medium for lactic Streptococci and their bacteriophages. *Applied Microbiology*, **29**, 807-813.

Torriani, S., Felis, G. and Dellaglio, F. (2001). Differentiation of *Lactobacillus plantarum*, *L. pentosus*, and *L. paraplantarum* by *recA* gene sequence analysis and multiplex PCR assay with *recA* gene-derived primers. *Applied and Environmental Microbiology*, **67**, 3450-3454.

Ueda, K., Seki, T., Kudo, T., Yoshida, T. and Kataoka, M. (1999). Two distinct mechanisms cause heterogeneity of 16S rRNA. *Journal of Bacteriology*, **181**, 78-82.

Van der Meulen, R., Scheirlinck, I., Van Schoor, A., Huys, G., Vancanneyt, M., Vandamme, P. and De Vuyst, L. (2007). Population dynamics and metabolite target analysis of lactic acid bacteria during laboratory fermentations of wheat and spelt sourdoughs. *Applied and Environmental Microbiology*, **73**, 4741-4750.

Von Wintzingerode, F., Göbel, U.B. and Stackebrandt, E. (1997). Determination of microbial diversity in environmental samples: Pitfalls of PCR-based RNA analysis. *FEMS Microbiology Reviews*, **21**, 213-229.

Wacher, C., Canas, A., Cook, P.E., Barzana, E. and Owens, J.D. (1993). Sources of microorganisms in *pozol*, a traditional Mexican fermented maize dough. *World Journal of Microbiology and Biotechnology*, **9**, 269-274.

Ward, B.B. (2002). How many species of prokaryotes are there? *Proceedings of the National Academy of Sciences of the United States of America*, **99**, 10234-10236.



CHAPTER FOUR

USE OF STARTER CULTURES OF LACTIC ACID BACTERIA IN THE PRODUCTION OF *TING*, A SOUTH AFRICAN FERMENTED FOOD

TABLE OF CONTENTS

4.1 INTRODUCTION	106
4.2 MATERIALS AND METHODS	107
4.2.1 Bacterial isolates	107
4.2.2 Preparation of starter cultures	107
4.2.3 Preparation of <i>ting</i>	108
4.2.4 pH and acidity determinations.....	108
4.2.5 Microbiological analysis	110
4.2.5.1 Bacterial and spore counts.....	110
4.2.5.2 Identification of colonies isolated from spore count plates.....	110
4.2.5.3 Analysis of yeasts and moulds	110
4.2.5.4 Analysis of food-borne bacterial pathogens.....	112
4.2.6 Antimicrobial studies	115
4.2.6.1 Test for microbial inhibition by LAB isolates	115
4.2.6.2 Test for possible bacteriocin production by LAB isolates	115
4.2.6.3 Test for possible bacteriocin production by LAB culture supernatants.....	115
4.2.6.4 Test for acid production as antimicrobial property of LAB isolates	116
4.2.7 Sensory evaluation of <i>ting</i>	116
4.2.8 Statistical analyses.....	116
4.3 RESULTS	117
4.3.1 Bacterial counts	117
4.3.2 Yeast and mould counts, and identification of isolates	117
4.3.3 Occurrence of food-borne bacterial pathogens	119
4.3.4 Antimicrobial interactions between LAB isolates used as starter cultures ...	119
4.3.5 Changes in pH and titratable acidity	122
4.3.6 Sensory evaluation of <i>ting</i>	122
4.4 DISCUSSION	124
4.5 REFERENCES	130

ABSTRACT

Ting, a porridge popular amongst southern Africans for its sour taste, is produced by cooking fermented sorghum into a thick porridge. With a view to selecting the most appropriate starter for use in the production of *ting*, trial fermentations were performed using different defined starter cultures. The lactic acid bacterial strains (*Lactococcus lactis*, *Lactobacillus fermentum*, *Lb. plantarum* and *Lb. rhamnosus*) used in the starters had been isolated previously from spontaneous sorghum fermentations. All the starter cultures showed an ability to ferment the sorghum, as evidenced from lowering the pH from 6.4 to 3.79 - 4.0 and increasing the titratable acidity from 0.18% to 0.72 - 0.95% (w/w, lactic acid) in 24 h of fermentation. However, the lowest pH (3.3) and highest lactic acid (1.44%) was produced the fastest in non-sterilized sorghum fermented by *L. lactis*. This fermentation was characterized by an increase in the number of lactic acid bacteria and yeasts, while the mean aerobic plate count, Gram-negative and bacterial spore counts, as well as fungal counts decreased. Moreover, bacterial food-borne pathogens detected in naturally fermented porridge, i.e. *Bacillus cereus* and *Escherichia coli* biotype I, were absent in the *L. lactis*-inoculated fermentation batch. *Ting* produced from this fermented gruel also had sensory properties preferred by panelists. The strong sour taste and strong aroma were especially appealing to the panelists. Conversely, *ting* prepared from naturally fermented sorghum in 54 h was the least preferred by panelists.

Key words: sorghum fermentation; *ting*; lactic acid bacteria; starter culture; pathogens; antimicrobial; sensory attributes

4.1 INTRODUCTION

Fermentation of local staple food, usually cereals such as maize, sorghum or millet, is a traditional technology in Africa (Mensah, 1997; Oyewole, 1997; Holzapfel, 2002). Lactic acid bacteria (LAB) and yeasts have been reported to be predominant microorganisms in most of the African indigenous fermented foods (Nout, 1991; Olasupo *et al.*, 1997; Kunene *et al.*, 1999). Lactic acid fermentation not only produces desirable changes in the aroma, flavour and texture of fermented foods (Kandler, 1983; Hugenholtz *et al.*, 2000; Axelsson, 2004), but studies have shown that the survival and growth of pathogenic and spoilage microorganisms are also adversely affected (Nout *et al.*, 1989; Holzapfel *et al.*, 1995; Kingamkono *et al.*, 1995; Adams and Nicolaidis, 1997; Magula *et al.*, 2003). Antimicrobial activity caused by growth of LAB is due to a decrease in pH to below 4 in the food products, depletion of nutrients and production of antimicrobial compounds, including bacteriocins (De Vuyst and Vandamme, 1994; Parente and Ricciardi, 1999) and various organic acids such as lactic and acetic acids (De Vuyst, 2000; Messens and De Vuyst, 2002; Mante *et al.*, 2003).

Ting, which is made from fermented sorghum, is consumed extensively in South Africa (Boling and Eisener, 1982). This food is usually prepared by soaking sorghum powder for two to three days in water. Microorganisms associated with the raw materials, equipment and local environments ferment the sorghum powder, resulting in a gruel that is mixed with boiling water, cooked and then consumed as a sour porridge. In rural and informal settlements of South Africa where electricity is not readily available, *ting* is frequently used as a weaning food for infants mainly because it is inexpensive to prepare, can be stored at ambient temperatures and does not require re-heating before consumption (Kunene *et al.*, 1999). Moreover, due to its appetizing taste, *ting* is consumed by adults at major ceremonies such as weddings and funerals. However, since the sorghum is allowed to undergo natural fermentation, variations in the sensory quality and acceptability of *ting* is frequently encountered.

The use of starter cultures may represent an appropriate approach for the control and optimization of the fermentation process in order to alleviate problems regarding the quality and acceptability of African indigenous fermented foods (Holzapfel *et al.*, 1995; Kimaryo *et al.*, 2000; Leroy and De Vuyst, 2004). The primary consideration before introducing starter cultures for traditional fermentations should be whether these would contribute significantly

to an improvement of processing conditions and product quality with respect to rapid and accelerated acidification, improved microbiological safety, an improved and more predictable fermentation process, and desirable sensory characteristics (Holzapfel, 2002; Leroy and De Vuyst, 2004).

Towards selecting an appropriate starter culture for the manufacture of *ting*, a study aimed at identifying the lactic acid flora present in the natural fermentation of sorghum had been undertaken previously. Using DNA-based molecular and bacteriological culturing approaches, the bacterial species *L. lactis*, *Lb. fermentum*, *Lb. plantarum* and *Lb. rhamnosus* were confirmed to be present during fermentation, but *L. lactis* appeared to be dominant during the final stages of fermentation (Chapter 3). Consequently, four starter cultures, each inclusive of *L. lactis*, were prepared by making use of the above-mentioned LAB species isolated from natural fermentations of sorghum. Thus, the aims of this study were to assess bacterial, yeast and mould populations found in sorghum before, during and after its preparation as a fermented cereal, to investigate possible antimicrobial interactions that may occur between the LAB species comprising the starter cultures, and to clarify the contribution of the LAB species with respect to the sensory characteristics of *ting*.

4.2 MATERIALS AND METHODS

4.2.1 Bacterial isolates

Starter cultures were prepared using LAB isolates that had been isolated previously from naturally fermented sorghum and of which their identity had been confirmed using 16S rRNA gene sequence analysis. The cultures of LAB (*L. lactis*, *Lb. fermentum*, *Lb. plantarum* and *Lb. rhamnosus*) had been stored at -20°C in sterile Eppendorf tubes, containing DeMan, Rogosa and Sharpe (MRS) broth (Merck, Darmstadt, Germany) with 20% (v/v) glycerol.

4.2.2 Preparation of starter cultures

The LAB starter cultures were prepared using a previously reported method (Mugula *et al.*, 2003a). Briefly, the LAB were individually streaked onto MRS agar (Merck) and incubated anaerobically in an anaerobic jar with anaerocult A (Merck) at 30°C for 24 h. A single colony was picked from each pure culture plate, grown overnight in MRS broth, followed by centrifugation at $655 \times g$ for 15 min in a benchtop centrifuge (Eppendorf, Hamburg, Germany). The cell pellets were washed twice in buffered peptone water (0.85% [w/v] NaCl,

0.1% [w/v] peptone, pH 7.2) and then resuspended in 1 ml of the solution. This procedure achieved a culture preparation containing 10^9 colony forming units (cfu)/ml, as determined by viable counts on MRS agar. Starter cultures E1, E2, E3 and E4 were considered. E1 was prepared from *L. lactis*; E2 was prepared from *L. lactis* and *Lb. fermentum*; E3 was prepared from *L. lactis*, *Lb. fermentum* and *Lb. plantarum*; and E4 was prepared from a combination of all LAB cultures (*L. lactis*, *Lb. fermentum*, *Lb. plantarum* and *Lb. rhamnosus*). Suspensions of each selected strain (1 ml) were combined, made up to 20 ml with buffered peptone water, mixed thoroughly and then inoculated into individual sorghum powder slurries used in the preparation of *ting*.

4.2.3 Preparation of *ting*

Six trial fermentations were prepared (Table 4.1). In addition to untreated (non-sterilized) sorghum powder inoculated with and without *L. lactis*, sterilized sorghum powders were inoculated separately with the respective starter cultures. Commercial packets of pure grain sorghum ('King Korn Mabele', King Food Corporation, Potchefstroom, South Africa) were purchased from a local supermarket. For natural fermentations, the sorghum powder was mixed with sterile luke-warm (ca. 40°C) water (1:1 [w/v]) in glass containers to make a slurry. The containers were covered and incubated at 25°C for 54 h. For inoculated fermentations, the sorghum powder was autoclaved at 121°C for 15 min prior to preparation of the slurry, as described above. The slurry was subsequently inoculated with the appropriate LAB starter culture, thoroughly mixed and incubated at 25°C for 24 h. The cooking process involved mixing 200 ml of the resultant fermented slurry with 600 ml of boiling (96°C) water, followed by cooking for 15 min. A further 100 ml of the fermented slurry was then added slowly until a thick porridge formed. The porridge was left to simmer for 15 min and stirred intermittently.

4.2.4 pH and acidity determinations

Samples of fermenting sorghum were withdrawn at 6-h intervals starting at $t = 0$ h until the end of fermentation ($t = 24$ h for inoculated fermentations and $t = 54$ h for natural fermentation). pH was measured using a Beckman model Ø 34 pH meter (Beckman Coulter, Fullerton, CA, USA) after calibration with standard buffers (Merck) at pH 4.0 and 7.0. For determination of titratable acidity of the fermented sorghum slurries, 10 g of sample was thoroughly drained and then titrated against 0.1 N NaOH to a pH of 8.5, using

Table 4.1 Experimental protocol for the six sorghum fermentations

Fermentation batch	Sorghum powder¹	Inoculum	Fermentation time (h)
NF	Untreated	Natural microflora	54
E1ns	Untreated	<i>L. lactis</i>	24
E1s	Sterilized	<i>L. lactis</i>	24
E2	Sterilized	<i>L. lactis</i> and <i>Lb. fermentum</i>	24
E3	Sterilized	<i>L. lactis</i> , <i>Lb. fermentum</i> and <i>Lb. plantarum</i>	24
E4	Sterilized	<i>L. lactis</i> , <i>Lb. fermentum</i> , <i>Lb. plantarum</i> and <i>Lb. rhamnosus</i>	24

¹ The respective starter cultures were inoculated directly into either slurries of untreated sorghum powder or slurries made from sorghum powder that was autoclaved at 121°C for 15 min.

phenolphthalein as indicator, as described previously (Nout *et al.*, 1989). One ml of 0.1 M NaOH was taken as equivalent to 0.0090 g lactic acid (Nout *et al.*, 1989). Measurements were repeated three times in each type of fermentation and the mean values recorded.

4.2.5 Microbiological analysis

4.2.5.1 Bacterial and spore counts

Sorghum powder, together with the fermented sorghum slurries and the corresponding cooked fermented *ting* porridges (Table 4.1), were aseptically processed and tested within 20 min of collection. For each sample, 10 g was homogenized in 90 ml of buffered peptone water, resulting in a 10^{-1} dilution. Ten-fold serial dilutions in the same diluent were plated onto different media and incubated for 48-72 h under either aerobic or anaerobic conditions (Table 4.2). Plates showing between 30 and 300 colony forming units (cfu), or the highest number if below 30, were counted following incubation. Bacterial spore counts were obtained by heating 10 ml of the 10^{-1} dilution at 80°C for 10 min in a water bath, followed by cooling on ice and spread plating onto Plate Count Agar (PCA) (Merck) (Becker *et al.*, 1994). All samples were processed in duplicate.

4.2.5.2 Identification of colonies isolated from spore count plates

A total of 65 colonies, isolated from plates of the highest dilution showing growth from sorghum powder (50 isolates) and the naturally fermented sorghum slurry (15 isolates), were purified on PCA agar (Table 4.2). These isolates were subjected to standard Gram stains and catalase tests. Gram-positive, catalase-positive isolates that showed phase bright spores were identified as *Bacillus* spp. (Ehrich, 1956), while *B. cereus* was identified with a culture-based approach, as described below.

4.2.5.3 Analysis of yeasts and moulds

For isolation of yeasts, the samples (1 g of sorghum powder and aliquots of 100 μ l of 10^{-1} dilutions of fermented sorghum slurries and *ting*) were plated onto Malt Extract Agar (MEA) (Merck), containing 125 mg/l of chloramphenicol (Sigma, St. Louis, MO, USA), followed by incubation at 25°C for 3-5 days. The yeast isolates were identified based on morphological characteristics (Jespersen *et al.*, 2005) and by using ID32C diagnostic test kits (bioMérieux, France), assisted by computer software (bioMérieux). This identification system was based

Table 4.2 Culture conditions used for microbiological analysis of sorghum samples

Enumeration type	Incubation temperature (°C)	Plating technique	Incubation period (h)	Atmospheric conditions	Growth medium
Aerobic plate count (APC)	30	Pour-plate	72	Aerobic	Plate Count Agar
Gram-negative count (GNC)	37	Pour-plate	24-48	Aerobic	Violet Red Bile Glucose Agar
Spore count (SC)	37	Spread-plate	24-48	Aerobic	Plate Count Agar
Lactic acid bacteria count (LABC)	30	Spread-plate	48-72	Anaerobic	MRS agar with 0.1% L-cysteine (Dykes <i>et al.</i> , 1991) and 0.01% cycloheximide (Nout <i>et al.</i> , 1989)
Yeasts	25	Spread-plate	120	Aerobic	Malt Extract Agar (MEA) with 125 mg/l chloramphenicol
Moulds	25	Direct plating and spread plating	360	Aerobic	Potato Dextrose Agar (PDA), Malt Salt Agar (MSA), Pentochloronitrobenzene (PCNB) agar. PDA medium was supplemented with 125 mg/l chloramphenicol

on 29 carbohydrate assimilation profiles (carbohydrates, organic acids, amino acids), one susceptibility test (cycloheximide) and an esculin hydrolysis colorimetric test.

Moulds were isolated on Potato Dextrose Agar (PDA), Malt Salt Agar (MSA) and Pentachloronitrobenzene (PCNB) agar (Merck), as described previously (Rabie and Lübben, 1984; Rabie *et al.*, 1997). Sorghum powder (1 g) was inoculated at almost equidistant positions on five points in Petri dishes containing the respective agar media (10 for each type of medium) and incubated at 25°C for 15 days. Aliquots of 1 ml of 10⁻¹ dilutions of fermented sorghum slurries and *ting* were also streaked onto the above agar media and incubated at 25°C for 15 days. Moulds were microscopically identified based on morphological characteristics to either genus or species level, depending on the ability of fungi to form fruiting structures and spores (Pitt, 1988). The identifications were confirmed by qualified mycologists.

4.2.5.4 Analysis of food-borne bacterial pathogens

Bacillus cereus

Robertson's heated cooked meat (100 ml) (Merck) was mixed with 50 ml of the 10⁻¹ dilution and incubated at 37°C for 48 h (Kramer and Gilbert, 1989). The broth (100 µl) was spread onto Columbia blood agar (Oxoid, Basingstoke, UK) and Polymyxin-Pyruvate-Egg Yolk Mannitol-Bromothymol Blue agar (PEMBA) (Oxoid), and incubated at 37°C for 24 h (ICMSF, 1996). Thereafter, PEMBA plates were examined for discoloration and blood agar plates for haemolysis, which is suggestive of *B. cereus*.

Campylobacter spp.

Preston broth (225 ml) (Oxoid) was mixed with 25 g of sample and incubated at 42°C for 24 h. The broth (100 µl) was plated onto blood-free *Campylobacter* agar (Oxoid) and Columbia blood agar (Oxoid), and incubated at 42°C for 48 h under microaerophilic conditions (Beuchat, 1987).

Clostridium perfringens

Aliquots of 100 µl of the 10⁻¹ dilution were spread onto Clostridial agar (Oxoid) and Salicin Tryptic Soy agar (STSA) (Difco, Detroit, MI, USA). The plates were incubated at 37°C for

48 h aerobically and anaerobically (anaerocult A) to test for scant or no growth on aerobic plates and for growth on the anaerobic plates (SABS, 1975a, method 761).

***Escherichia coli* biotype I**

An aliquot of 1 ml of the 10^{-1} dilution was added into test tubes, containing 10 ml of MacConkey broth purple (Merck) and sterile Durham tubes, and incubated at 37°C for 24 h. Aliquots of 100 µl of the broth from test tubes showing gas and acid production (yellow colour) was inoculated into Brilliant green bile broth (Oxoid) and Tryptone water (Oxoid), and incubated at 44°C for 18 h. Production of gas in Brilliant green bile broth and indole in tryptone water (red colour after addition of Kovac's reagent [Merck]) indicated the presence of *E. coli* biotype I (SABS, 1975b, method 758). Gram-negative lactose fermenting colonies were subjected to further biochemical tests, as described by Schlegel (1988), and all tests were performed in duplicate.

***Escherichia coli* 0157:H7**

An aliquot of 100 µl of the 10^{-1} dilution was spread onto MacConkey sorbitol agar (Oxoid) and incubated at 37°C for 24 h to select for sorbitol-negative *E. coli* 0157:H7 (Vernozy-Rozand, 1997).

Listeria monocytogenes

Listeria enrichment broth (225 ml) (Difco) was inoculated with 25 g of sample and incubated at 37°C for five weeks. At the end of each week, the broth (100 µl) was spread onto *Listeria* selective agar (Oxoid), incubated at 37°C for 24 h and examined for the presence of black colonies (Van Netten *et al.*, 1988).

***Salmonella* spp.**

Pre-enrichment was carried out by homogenizing 25-g samples of sorghum powder, fermented sorghum slurries or *ting* in 225 ml of buffered peptone water, followed by incubation at 37°C for 24 h (Arvanitidou *et al.*, 1998). Selective enrichment for *Salmonella* spp. was done by inoculation of 10 ml of the buffered peptone water into 100 ml of Tetrathionate medium (Oxoid) and incubation at 43°C for 48 h. In addition, buffered peptone water (10 ml) was inoculated into 100 ml of Selenite broth (Difco) and Gram-negative medium (Merck), and then incubated at 37°C for 48 h (Arvanitidou *et al.*, 1998). The respective broths were streaked onto Xylose, Lysine, Desoxycholate (XLD) (Difco) and

Brilliant Green Agar (BGA) (Oxoid), and incubated at 37°C for 24 h. Lactose-negative colonies are indicative of the presence of *Salmonella* spp.

***Shigella* spp.**

Gram-negative medium (100 ml) (Merck) was mixed with 25 ml of the 10⁻¹ dilution and incubated at 37°C for 24 h. The broth (100 µl) was streaked onto XLD agar plates and incubated at 37°C for 24 h. Translucent colonies are suggestive of the presence of *Shigella* spp. (Taylor, 1965).

Staphylococcus aureus

Three bottles, each containing 20 ml of *Staphylococcus* enrichment broth (Difco), were mixed with 1 ml of the 10⁻¹ dilution and incubated at 37°C for 48 h after which they were examined for gas production (SABS, 1975c, method 760). The broth (100 µl) was streaked onto Baird-Parker agar (Merck) and incubated at 37°C for 24 h, after which plates were examined for black colonies surrounded by a clear zone that are indicative of *S. aureus*.

***Vibrio* spp.**

A 25-g sample was homogenized in 225 ml of buffered peptone water and an aliquot of 50 ml was then inoculated into two bottles containing 50 ml of *Vibrio* enrichment broth (Oxoid), followed by incubation for 24 h, one at 37°C and one at 42°C. Loopfuls from the surface of the broths were streaked onto Thiosulphate Citrate Bile Sucrose (TCBS) agar (Difco) and incubated at 37°C for 24 h. Thereafter plates were examined for flat yellow colonies and/or smooth green colonies, which are indicative of *Vibrio* spp. (Madden and McCardell, 1989).

***Yersinia* spp.**

A 10-g sample was homogenized in 100 ml of phosphate-buffered saline (3.2 mM Na₂HPO₄, 0.5 mM KH₂PO₄, 1.3 mM KCl, 135 mM NaCl, pH 7.4) and incubated at 4°C for four weeks (Sutherland and Varnam, 1977). Loopfuls from the broth were streaked onto *Yersinia* selective agar (Difco) and incubated aerobically at 30°C for 24-48 h. *Yersinia* spp. ferment mannitol that result in acid production and thus are indicated by red discolouration of the colonies surrounded by a transparent zone (Schiemann, 1979).

4.2.6 Antimicrobial studies

4.2.6.1 Test for microbial inhibition by LAB isolates

The inhibitory potential of LAB cultures used in starter culture formulations were investigated *in vitro* using the Agar Well Assay method (Schillinger and Lücke, 1989; Olsen *et al.*, 1995). For this purpose, MRS agar and M17 agar (Merck) were poured into Petri dishes and left to solidify for two days. A sterile cork borer was used to make circular wells into the agar. *L. lactis*, *Lb. fermentum*, *Lb. plantarum* and *Lb. rhamnosus* were each cultured separately in MRS broth (Merck) at 30°C for 24 h, and 100 µl of the cultures (1×10^8 cfu/ml) were transferred into the wells and left to diffuse into the agar for 5 h. The wells were overlaid with ca. 10 ml of soft MRS agar (0.7% agar) containing the above LAB isolates as indicator strains. The indicator lawns were prepared by adding 250 µl of a 10^{-1} dilution of an overnight culture of the indicator bacterium to 10 ml of MRS agar (Merck). The plates were incubated at 30°C for 24-48 h and the diameter of clear zones around the wells were measured and regarded as inhibitory reactions (Mante *et al.*, 2003). Individual tests were carried out in duplicate.

4.2.6.2 Test for possible bacteriocin production by LAB isolates

Volumes of 100 µl of overnight LAB cultures were transferred separately into MRS agar wells and overlaid with 10 ml of MRS soft agar containing *Listeria monocytogenes* ATCC 19115, as bacteriocin indicator organism. The agar plates were incubated aerobically at 37°C for 24 h and examined for inhibitory reactions, as evidenced by zones of inhibition. The experiments were carried out in duplicate.

4.2.6.3 Test for possible bacteriocin production by LAB culture supernatants

For each of the LAB cultures tested, 10 ml of an overnight culture was centrifuged at $3\,000 \times g$ for 15 min at 4°C, and the cell-free supernatants filtered using a 0.20-µm sterile filter (Sartorius Minisart, Cottingen, Germany). Cell-free extracts (100 µl) were digested separately with Proteinase K (Merck) and Pronase E (Sigma), at a concentration of 1 mg/ml at 30°C for 2 h, and then tested against the *L. monocytogenes* indicator strain using the Agar Well Assay method. The agar plates were incubated aerobically at 37°C for 24 h.

4.2.6.4 Test for acid production as antimicrobial property of LAB isolates

Cell-free extracts (100 µl) of the LAB cultures were neutralized by adding 0.1 M NaOH to the supernatants to raise their pH to 6.8. Both neutralized and non-neutralized supernatants were filter-sterilized through a 0.20-µm filter (Sartorius Minisart). The neutralized and non-neutralized supernatants were tested separately against *Listeria monocytogenes* using the Agar Well Assay method.

4.2.7 Sensory evaluation of *ting*

Sensory evaluation of the different *ting* porridges was done at the sensory laboratory of the Department of Food Science, University of Pretoria, South Africa. Randomly selected three-digit numbers were used for coding the *ting* samples that were presented in random order to the consumer panel for sensory evaluation. The acceptability of aroma, appearance, mouth feel, taste, aftertaste and overall acceptance of *ting* was evaluated. The panelists scored the seven characteristics using a 9-point hedonic scale (1 = dislike extremely; 2 = dislike very much; 3 = dislike moderately; 4 = dislike slightly; 5 = neither like nor dislike; 6 = like slightly; 7 = like moderately; 8 = like very much; and 9 = like extremely) (Stone and Siddel, 1992). The panelists took between 15 and 20 min to rate the samples. A total of 50 panelists (23 men and 27 women), who were all familiar with the taste of *ting*, participated in the study. The panelists were given water and a piece of carrot before and in between tasting the *ting* samples in order to cleanse their palates. Data was recorded using Compusense[®] version 5.4.6 (Compusense Inc., Guelph, ON, Canada).

4.2.8 Statistical analyses

The results of the sensory analysis were subjected to analysis of variance (ANOVA), using a significance level of $P < 0.05$. Fischer's least significant difference (LSD) test was also applied to ascertain the significance of differences in the average values for each of the attributes for the different *ting* samples. Pearson's product-moment correlation coefficient (r) was used to test the correlation between pH, titratable acidity and acceptance of other sensory attributes at $P < 0.05$. All statistical analyses were performed using Statistica version 7.1 (StatSoft Inc., Tulsa, OK, USA).

4.3 RESULTS

4.3.1 Bacterial counts

Bacterial counts were determined for the sorghum powder, the fermented sorghum slurries ($t = 24$ h for inoculated fermentations and $t = 54$ h for natural fermentation) and for the corresponding cooked fermented porridges. For the sorghum powder, aerobic plate counts (APC) were the highest, followed by counts of lactic acid bacteria (LABC) and Gram-negative counts (GNC), while spore counts (SC) were the lowest. Compared to sorghum powder, the mean APC and GNC of the naturally fermented sorghum slurry increased by 2.1 and 2.07 log cfu/g, respectively, while the SC decreased by 1 log cfu/g. In contrast, the mean APC of inoculated fermented sorghum slurries decreased by more than 2 log cfu/g, while the mean SC and GNC decreased to below the detection limit (<2.00 log cfu/g). The LABC exhibited increases in excess of 5 log cfu/g in both natural and inoculated fermented sorghum slurries. In cooked fermented porridge samples, the bacterial counts (APC, SC, GNC and LABC) were below the detection limit (Table 4.3).

4.3.2 Yeast and mould counts, and identification of isolates

Yeast and mould counts were determined concurrently with bacterial counts. Compared to sorghum powder, the mean yeast and mould counts of the naturally fermented sorghum slurry increased by 1.40 and 3.19 log cfu/g, respectively. The mould count of inoculated fermented sorghum slurries decreased to below the detection limit (<2.00 log cfu/g) as did the yeast count, except for *L. lactis*-inoculated untreated sorghum which exhibited an increase of 2.58 log cfu/g in the mean yeast count. In cooked fermented porridge samples, the yeast and mould counts were below the detection limit (Table 4.3).

Yeasts isolated from sorghum powder and fermented sorghum slurries were identified by the use of ID32C diagnostic kits. The 16 isolates were tentatively identified as *Cryptococcus laurentii* (75%) and *Cryptococcus humicolus* (25%). In contrast, various different moulds were isolated from sorghum powder used in the respective fermentations. The predominant species were *Alternaria* spp., *Cladosporium* spp., *Fusarium andiyazi*, *Penicillium* spp., *Mucor* spp. and *Phoma sorghina*. Fungi that occurred at low prevalence included *Trichoderma* spp., *Rhizopus oryzae*, *F. equiseti*, *Drechslera sativus*, *Drechslera halodes*, *Aspergillus terreus*, *Curvularia eragrostidis*, *Euroteum repens*, *Nigrospora* spp. and *F. chlamydosporum*. Of these fungi, however, *Mucor* spp. and *Penicillium* spp. were detected in the naturally fermented sorghum slurry at the end of fermentation only.

Table 4.3 Microbial counts (log cfu/g) of sorghum powder, fermented sorghum slurries and corresponding cooked fermented porridge (*ting*)

Analysis	Sorghum powder	Fermentation batch ¹						Cooked <i>ting</i>
		NF	E1ns	E1s	E2	E3	E4	
APC	5.84 (± 0.02)	7.90 (± 0.09)	2.91 (± 0.02)	2.18 (± 0.02)	2.18 (± 0.03)	2.32 (± 0.03)	2.50 (± 0.02)	<2.00
SC	2.04 (± 0.04)	<2.00	<2.00	<2.00	<2.00	<2.00	<2.00	<2.00
GNC	2.94 (± 0.01)	5.01 (± 0.02)	<2.00	<2.00	<2.00	<2.00	<2.00	<2.00
LABC	3.85 (± 0.01)	9.21 (± 0.02)	9.83 (± 0.03)	9.37 (± 0.04)	9.59 (± 0.02)	9.70 (± 0.02)	9.82 (± 0.02)	<2.00
Yeasts	2.83 (± 0.02)	4.23 (± 0.62)	5.41 (± 0.07)	<2.00	<2.00	<2.00	<2.00	<2.00
Moulds	2.31 (± 0.02)	5.50 (± 0.13)	<2.00	<2.00	<2.00	<2.00	<2.00	<2.00

¹ Refer to Table 4.1 for details of fermentation batches.

4.3.3 Occurrence of food-borne bacterial pathogens

Of the different food-borne pathogens tested for, only *B. cereus* and *E. coli* biotype I were detected in the sorghum powder and in the naturally fermented sorghum slurry. None of the pathogens tested for were detected in any of the inoculated sorghum slurries or in any of the cooked, fermented porridge samples (Table 4.4). Based on biochemical tests, isolates belonging to *E. coli* biotype I was identified as *Klebsiella* spp., *Escherichia* spp. and *Enterobacter* spp. The 65 isolates from spore count plates of sorghum powder and the naturally fermented sorghum slurry were all catalase-positive, rod-shaped and spore-forming, and, of these, only five (7.7%) were identified as *B. cereus*.

4.3.4 Antimicrobial interactions between LAB isolates used as starter cultures

The four LAB isolates used in the formulation of starter cultures were tested for inhibitory reactions against each other, the results of which are shown in Table 4.5. Of the different LAB isolates tested, inhibitory reactions were only observed for *L. lactis* and *Lb. plantarum*. Although *L. lactis* showed strong inhibitory reactions against both *Lb. fermentum* and *Lb. rhamnosus*, no inhibitory reaction was observed against *Lb. plantarum*. Similarly, *Lb. plantarum* showed strong inhibitory reaction against *Lb. fermentum*, but exhibited less interaction with *Lb. rhamnosus* and no inhibitory activity against *L. lactis*.

To determine the mechanism of *L. lactis* and *Lb. plantarum* inhibitory activity, the culture supernatants of the LAB isolates were tested against *L. monocytogenes* as an indicator of bacteriocin production. Both of the LAB isolates showed inhibitory reactions against *L. monocytogenes*. However, after neutralization of the culture supernatants, no inhibitory reactions were observed for the respective cultures. The addition of the proteolytic enzymes Proteinase K and Pronase to the supernatants did not have an effect on their inhibitory activity against the *L. monocytogenes* indicator bacterium (results not shown). Thus, it was concluded that the inhibitory action of *L. lactis* and *Lb. plantarum* was due to acids produced rather than the production of bacteriocins or other antimicrobial compounds.

Table 4.4 Occurrence of bacterial food-borne pathogens in sorghum powder, fermented sorghum slurries and cooked *ting*

Pathogen	Sorghum powder	Fermented slurries		<i>Ting</i>
		Natural	Inoculated sorghum slurries	
<i>B. cereus</i>	+	+	-	-
<i>Campylobacter</i> spp.	-	-	-	-
<i>C. perfringens</i>	-	-	-	-
<i>L. monocytogenes</i>	-	-	-	-
<i>Salmonella</i> spp.	-	-	-	-
<i>Shigella</i> spp.	-	-	-	-
<i>E. coli</i> biotype I	+	+	-	-
<i>S. aureus</i>	-	-	-	-
<i>Vibrio</i> spp.	-	-	-	-
<i>Yersinia</i> spp.	-	-	-	-
<i>E. coli</i> 0157:H7	-	-	-	-

-: not detected; +: detected.



Table 4.5 Antimicrobial interactions among lactic acid bacteria used as starter cultures for sorghum fermentation

LAB test strains	LAB indicator strains			
	<i>L. lactis</i>	<i>Lb. fermentum</i>	<i>Lb. plantarum</i>	<i>Lb. rhamnosus</i>
<i>L. lactis</i>	-	++	-	++
<i>Lb. fermentum</i>	-	-	-	-
<i>Lb. plantarum</i>	-	++	-	+
<i>Lb. rhamnosus</i>	-	-	-	-

-: no inhibition; +: 5-8 mm inhibition zone; ++ : >8 mm inhibition zone.


4.3.5 Changes in pH and titratable acidity

The changes in pH and titratable acidity of sorghum slurries fermented naturally for 54 h and of inoculated slurries fermented for 24 h are shown in Fig. 4.1. For all six fermentations, a decrease in pH was accompanied by a simultaneous increase in titratable acidity. In the naturally fermented sorghum slurry, the pH decreased from 6.64 to 4.64 in 24 h and to pH 3.75 in 54 h, with a simultaneous increase in titratable acidity (% w/w, lactic acid) from 0.13% to 0.65% and to 0.84% in 24 and 54 h of fermentation, respectively (Fig. 4.1). After 24 h, the pH of sorghum powder slurries prepared from sterilized sorghum powder and inoculated separately with the starter cultures E1, E2, E3 and E4 decreased from 6.4 to 3.79 - 4.0, with a simultaneous increase in titratable acidity from 0.18% to 0.72% for fermentations by E2 and E3, and to 0.90% and 0.95% for fermentations by E4 and E1, respectively. Inoculation of slurry prepared from untreated (non-sterilized) sorghum powder with *L. lactis*, however, resulted in the most significant and rapid decrease in pH (3.3) and increase in acidity (1.44%) in 24 h of fermentation.

4.3.6 Sensory evaluation of *ting*

Attributes such as aroma and sourness, especially, makes *ting* very popular amongst consumers. The results of the sensory analysis (Table 4.6) indicated that for aroma, *ting* prepared from untreated sorghum powder inoculated with *L. lactis* as starter culture (batch E1ns) was more acceptable than *ting* prepared from both naturally fermented sorghum powder (batch NF) and from sterilized sorghum powder inoculated with the same starter culture (batch E1s). In addition, these samples differed significantly ($P < 0.05$) with respect to acceptability of aroma. The acceptability of *ting* prepared from batches E2, E3 and E4 did not differ significantly with respect to aroma. *Ting* prepared from batch E1ns was clearly most acceptable compared to the rest of the samples as far as sourness was concerned. On the contrary, *ting* prepared from batches E1s, E2, E3, E4 and NF did not differ significantly ($P > 0.05$) with respect to acceptability of sourness. Although there was a strong negative correlation between pH and titratable acidity ($r = -0.8494$; Table 4.7), there were no significant correlations ($P > 0.05$) between pH and acceptance of any of the sensory attributes. However, titratable acidity had a strong positive correlation with sourness ($r = 0.8389$; Table 4.7). Indeed, sourness ($r = 0.8776$) was among other attributes (aroma, appearance and taste) that exhibited a strong positive correlation with overall acceptability of *ting*. Although there were no significant differences ($P > 0.05$) regarding acceptability of the taste and aftertaste of

(A)



(B)




Figure 4.1 Changes in pH (A) and titratable acidity (B) in sorghum powder slurries during natural fermentation (NF) for 54 h and inoculated fermentations with different LAB starter cultures (E1, E2, E3 and E4) for 24 h. LAB starter cultures were as follow: E1, *L. lactis*; E2, *L. lactis* + *Lb. fermentum*; E3, *L. lactis* + *Lb. fermentum* + *Lb. plantarum*; E4, *L. lactis* + *Lb. fermentum* + *Lb. plantarum* + *Lb. rhamnosus*. Prior to inoculation with *L. lactis* the sorghum powder was left untreated (non-sterile; E1ns) or autoclaved (sterilized; E1s). Averaged pH and titratable acidity are indicated.

the respective *ting* samples, there were, however, significant differences ($P < 0.05$) for acceptability of mouth feel, as well as overall acceptability of the *ting* samples (Table 4.6). Based on the results obtained, *ting* prepared from fermentation batch E1ns was the most acceptable amongst the panelists. However, the results also indicated that the overall acceptability of *ting* prepared from sterilized sorghum using starter culture E3 did not differ significantly from *ting* prepared from batch E1ns.

4.4 DISCUSSION

The use of spontaneous fermentation, albeit a simple way to achieve *ting* fermentation, involves a complex microbial process that results in a product of variable quality. It would therefore be desirable to change the fermentation from a wholly natural and unpredictable process to a process that can be manipulated and improved to yield more reliable results. Since lactic acid-fermented porridges have been reported to improve the microbiological stability, as well as nutritional and sensory properties of the product (Chavan and Kadam, 1989; Mugula *et al.*, 2003a; Hammes *et al.*, 2005), an attempt was therefore made to replace the natural fermentation with fermentations conducted with defined starter cultures comprising of different lactic acid bacterial isolates.

The sorghum powder used in the present study was found to be contaminated with Gram-negative bacteria and bacterial spores at levels that are in agreement with previously reported contamination levels of sorghum powder before fermentation (Nout, 1991; Kunene *et al.*, 1999). The bacterial populations were reduced to below the detection limit (< 2.00 cfu/g) in fermented porridge samples prepared from sorghum powder that had been sterilized by autoclaving prior to inoculation with different LAB starter cultures. Despite the non-sterile character of the fermentation, similar results were obtained for fermented porridge prepared from untreated sorghum powder inoculated with *L. lactis* only. In contrast, for fermented porridge samples prepared by natural fermentation, the bacterial spore count was reduced (< 2.00 cfu/g), but both the aerobic plate count (7.9 log cfu/g) and Gram-negative count (5.01 log cfu/g) was increased compared to sorghum powder. Indeed, cooking of the fermented porridge for 30 min was required to reduce the bacterial populations to below the detection limit. However, in all of the fermented porridge samples, lactic acid bacteria were dominant (9.21 - 9.83 log cfu/g), reinforcing results obtained by Rombouts and Nout (1995) who reported LABC of 9.5 - 9.7 cfu/g in fermented sorghum.

Table 4.6 Acceptability of the sensory characteristics of *ting* produced, using different starter cultures, based on average hedonic ratings

Fermentation batch ¹	Sensory attribute scores ²						
	Aroma	Appearance	Sourness	Mouth feel	Taste	Aftertaste	Overall acceptance
NF	5.0 (±2.68)c	6.0 (±2.01)c	5.0 (±2.23)b	5.7 (±1.90)bc	5.3 (±2.27)	5.5 (±2.18)	5.4 (±2.31)c
E1ns	6.7 (±1.99)a	6.8 (±1.60)a	6.4 (±2.11)a	6.4 (±1.85)a	6.2 (±2.30)	6.3 (±1.82)	6.5 (±2.08)a
E1s	5.7 (±2.31)b	6.5 (±1.66)abc	5.3 (±2.21)b	6.1 (±2.03)abc	5.7 (±2.19)	5.8 (±1.83)	5.8 (±2.07)bc
E2	6.1 (±2.20)ab	6.3 (±1.76)bc	5.0 (±2.50)b	5.7(±2.03)c	5.7 (±2.13)	6.0 (±2.09)	5.8 (±2.40)bc
E3	6.2 (±1.94)ab	6.6 (±1.40)ab	5.6 (±2.38)b	6.4 (±1.65)a	6.1(±2.13)	6.0 (±1.91)	6.2 (±1.99)ab
E4	6.0 (±2.00)ab	6.2(±1.82)bc	5.5 (±2.28)b	6.3(±1.76)ab	5.5 (±2.36)	5.8 (±1.90)	5.7 (±2.27)bc

¹ See Table 4.1 for inoculum information.

² Sensory attributes of the various *ting* preparations are indicated on a Hedonic scale (1 = Dislike extremely, 9 = Like extremely). The standard deviation of each score is indicated in parentheses. Average score values in the same column that differ significantly ($P < 0.05$) are indicated with different letters (a, b or c).

Table 4.7 Pearson's product-moment correlation coefficients (r)^a for pH, titratable acidity and acceptability of sensory attributes associated with *ting* produced using different starter cultures

	Titratable acidity	Aroma	Appearance	Sourness	Mouth feel	Taste	Aftertaste	Overall acceptance
pH	(-0.8484) $P = 0.032$	(-0.4899) $P = 0.324$	(-0.4174) $P = 0.410$	(-0.7944) $P = 0.059$	(-0.36030) $P = 0.483$	(-0.4390) $P = 0.384$	(-0.5468) $P = 0.262$	(-0.5788) $P = 0.229$
Titratable acidity		(0.4967) $P = 0.316$	(0.5857) $P = 0.222$	(0.8389) $P = 0.037$	(0.4642) $P = 0.354$	(0.4628) $P = 0.355$	(0.5546) $P = 0.253$	(0.6003) $P = 0.208$
Aroma			(0.8257) $P = 0.043$	(0.7779) $P = 0.069$	(0.6746) $P = 0.142$	(0.8620) $P = 0.027$	(0.9705) $P = 0.01$	(0.9026) $P = 0.014$
Appearance				(0.8091) $P = 0.051$	(0.7139) $P = 0.111$	(0.9602) $P = 0.002$	(0.8744) $P = 0.023$	(0.9558) $P = 0.003$
Sourness					(0.8188) $P = 0.046$	(0.7801) $P = 0.067$	(0.7736) $P = 0.071$	(0.8776) $P = 0.022$
Mouth feel						(0.6880) $P = 0.131$	(0.5670) $P = 0.241$	(0.7337) $P = 0.097$
Taste							(0.9075) $P = 0.012$	(0.9815) $P = 0.001$
Aftertaste								(0.9368) $P = 0.006$

^a Correlation coefficients (r) are given in brackets. Significant correlations are indicated by $P < 0.05$.

The results presented here indicated that not only does natural fermentation appear to be ineffective in reducing different bacterial populations, but also that addition of *L. lactis* to untreated sorghum slurry is responsible for reducing the bacterial populations effectively during fermentation. Reductions in bacterial populations in LAB-based fermentations have been attributed to a decrease in pH of the fermented gruel (Mante *et al.*, 2003) and to the production of organic acids by LAB (Messens and De Vuyst, 2002) and possibly other inhibitory substances (De Vuyst and Vandamme, 1994). Specifically, rapid acidification has been reported to be desirable to not only enhance the sensory characteristics of traditional fermented foods, but also to inhibit growth of pathogens and spoilage bacteria (Kingamkono *et al.*, 1995; Mugula *et al.*, 2003a; Ammor and Mayo, 2007). In this regard, it is important to note that the lowest pH (3.3) and highest levels of lactic acid (1.44%) were produced the fastest in untreated sorghum inoculated with *L. lactis* (t = 24 h). In contrast, the decrease in pH observed in naturally fermented sorghum was slow and gradual, reaching a pH of 4.6 after 24 h and a final pH of 3.75 at the end of fermentation (t = 54 h) (Fig. 4.1). Due to the changing conditions brought about by the fermentation processes, the bacterial populations in the fermented porridge samples are likely not to have been the same compared to the sorghum powder. Thus, it is possible that the gradual decrease in pH observed during natural fermentation might have selected for and supported the proliferation of acid-tolerant bacteria, thereby accounting for the increase in the aerobic plate count and Gram-negative count. Moreover, the above-mentioned fermentation processes were both found to result in a decrease in bacterial spore counts to below the detection limit (< 2.00 log cfu/g). Previous studies on the survival of spores at low pH have reported a decrease in the counts of *B. cereus* in media with pH values below 5 (Wong and Chen, 1988).

With the exception of *B. cereus* and *E. coli* biotype I, which were both detected in sorghum powder and in naturally fermented porridge samples only, none of the bacterial pathogens tested for were detected in any of the samples analyzed. Both *B. cereus* (Kramer and Gilbert, 1989) and members of the *Enterobacteriaceae* family (Gassem, 1999; Nout, 1991; Mohammed *et al.*, 1991) are common contaminants of agricultural commodities, including sorghum crops, and their presence in the samples could have resulted from contamination of the sorghum powder during manufacture and packaging. The absence of *B. cereus* in fermented porridge samples from inoculated fermentations is in agreement with a related study of Kingamkono *et al.* (1994), who also reported an absence of *B. cereus* in lactic acid-fermenting cereal gruels after 24 h. A number of investigations have reported on the survival

of *Enterobacteriaceae* in some traditional African fermented food. Although some of these have reported inhibition of the bacteria by LAB (Nout, 1991; Kunene *et al.*, 1999; Mugula *et al.*, 2003a), others have reported either a reduction in their numbers (Kingamkono *et al.*, 1995) or an increase in the number of coliforms and *E. coli*, especially when non-LAB starter cultures were used (Gran *et al.*, 2002; 2003). The survival and growth of *Enterobacteriaceae* at the end of the fermentation process has been ascribed to acid resistance of some strains and/or the presence of microenvironments in the food matrix that support the growth of these bacteria (Arnold and Kasper, 1995; Tomicka *et al.*, 1997). Furthermore, the ability of members of *Enterobacteriaceae* to attach to surfaces also results in an increased acid resistance of these bacteria (Rowbury, 1995). In the present study, the *Enterobacteriaceae* identified included *Escherichia* spp., *Klebsiella* spp. and *Enterobacter* spp.

The infestation of sorghum powder by a variety of moulds is in agreement with previous reports (Rabie and Lübben, 1984; Saubois *et al.*, 1999; Lefyedi *et al.*, 2005). Fungi such as *P. sorghina*, *Alternaria* spp. and *Cladosporium* spp., identified in the present study, may pose possible health risks due to the production of mycotoxins (Rabie *et al.*, 1975; Mazur and Kim, 2006). Moreover, elevated levels of moulds in raw materials used for food preparation may render the food unpalatable (Lefyedi *et al.*, 2005). However, different LAB species such as *L. lactis* (Roy, 1996), as well as *Lb. plantarum*, *Lb. rhamnosus* and *Lb. paracasei* have been reported to inhibit the growth of certain moulds (Corsetti *et al.*, 1998; Meroth *et al.*, 2003). Isolates of these species produce combinations of lactic acid, acetic acid, phenyllactic acid (PLA), hydrogen peroxide and other low molecular weight compounds that inhibit mold growth (Batish *et al.*, 1997; Lavermicocca *et al.*, 2003). Indeed, the naturally occurring moulds appeared unable to compete under the conditions prevailing in fermenting sorghum inoculated with *L. lactis* as their viable numbers were below the detection limit ($< 2.00 \log \text{ cfu/g}$), whilst only *Mucor* spp. and *Penicillium* spp. were detected in fermented porridge samples prepared by natural fermentation. This implies that there is less of a risk of mycotoxins being formed during the fermentation process, but the risk of raw materials being contaminated with mycotoxins can, however, not be excluded.

In addition to LAB, yeasts have also been reported to be common in a wide range of traditional African food and beverage fermentations (Olasupo *et al.*, 1997; Mugula *et al.*, 2003b; Kebede *et al.*, 2007). They have been reported to make a useful contribution to the improvement of flavour and acceptability of fermented cereal gruels (Akinrele, 1970; Odunfa

and Adeyele, 1985; Brauman *et al.*, 1996; Mugula *et al.*, 2003a). In the present study, yeasts were detected in both sorghum powder (2.83 log cfu/g) and in fermented sorghum slurries prepared from untreated sorghum inoculated without or with *L. lactis* (4.23 and 5.41 log cfu/g, respectively). The yeast isolates, *C. laurentii* and *C. humicola*, identified in this study, have been isolated previously from acidic fermentations of cereal substrates (Nout, 1991; Nago *et al.*, 1998; De Vuyst and Neysens, 2005). A co-metabolism between yeasts and lactic acid bacteria has been suggested, whereby the bacteria provide the acid environment, which selects for the growth of yeasts, and the yeasts provide vitamins and other growth factors to the bacteria (Alexander, 1971; Gobetti *et al.*, 1994; Steinkraus, 1996). However, despite low levels of yeasts being present in fermented sorghum slurries (<2.00 log cfu/g) prepared from sterilized sorghum powder inoculated with different LAB starter cultures (E1, E2, E3, and E4), the LAB were nevertheless able to proliferate to levels comparable with that determined in fermented sorghum slurries prepared from untreated sorghum powder inoculated without and with *L. lactis* (Table 4.3). Moreover, the lactic acid produced in these fermentations (0.72-0.95%) was comparable to that produced during natural fermentation (0.84%) (Fig. 4.1B). These results suggest that the role of yeasts during lactic acid fermentation is negligible (Martineau and Henick-Kling, 1995), but does not exclude a role in flavour development.

The results from sensory analysis of cooked fermented *ting* porridges prepared from untreated sorghum without or with *L. lactis*, respectively, showed that *ting* from the latter fermentation batch (E1ns) was rated by the panelists as being more acceptable than *ting* prepared from natural fermentation (NF). For *ting* production using untreated sorghum, it therefore appears that the contribution to the fermentation process by *L. lactis* is important. Not only does the presence of this species inhibit growth of fungi and bacteria (Table 4.3), possibly through an accelerated decrease in pH of the fermenting gruel and/or the production of organic acids, but it also exerts a positive influence on the sensory properties of *ting*. This is evidenced by *ting* prepared from *L. lactis*-inoculated untreated sorghum powder being awarded the highest sensory attribute scores. Although *L. lactis* produces diacetyl and acetylaldehyde, which are responsible for pleasant flavour in fermented foods (Hugenholtz *et al.*, 2000; De Vos and Hugenholtz, 2004), it is possible that yeasts present in the fermenting gruel may also contribute to flavour acceptability. This is evidenced by a significant difference ($P < 0.05$) with regards to the acceptability of the aroma of *ting* prepared from fermentation batches using untreated (E1ns) or sterilized (E1s) sorghum powder inoculated with *L. lactis* (Table 4.6). Conversely, *ting* prepared by natural fermentation earned the lowest attribute scores and

was the least acceptable of all the fermented *ting* porridges. The presence of microorganisms other than lactic acid bacteria during fermentation, especially fungi, may account for the unpalatability of this *ting*, as evidenced by low attribute scores for especially aroma, mouth-feel, taste and aftertaste.

The sensory attribute scores awarded to the cooked fermented *ting* porridges prepared from sterilized sorghum powder inoculated with starter cultures E1, E2, E3 and E4 (Table 4.2) fell in between the scores for the *ting* prepared from untreated sorghum inoculated without (least preferred) or with (most preferred) *L. lactis*. Even so, it is interesting to note that the sensory attribute scores for *ting* prepared from fermentation batch E3 always came second and its overall acceptability amongst panelist did not differ significantly ($P < 0.05$) from that of fermentation batch E1ns, which was the most acceptable (Table 4.6). Since both *L. lactis* and *Lb. plantarum* showed inhibitory reactions against *Lb. fermentum* and *L. rhamnosus* (Table 4.5), it is tempting to speculate that these two LAB strains may contribute only slightly to the sensory characteristics of *ting*. It thus follows that the contribution of *Lb. plantarum*, in addition to *L. lactis* mentioned above, is important.

Traditional natural fermentations in the alcoholic beverage industry have been replaced with defined inocula, high-quality raw materials, strict control of fermentation and better treatment of the final product, thereby resulting in diversification of the market. Traditional African fermented foods, including *ting*, have a long way to go before they reach a similar stage of development. Nevertheless, this is the first report on the use of defined starter cultures for *ting* fermentations. Of the different strains used, *L. lactis* was identified as the most promising candidate *ting* starter culture. This species can be used either in co-culture with *Lb. plantarum* for use in the fermentation of sterilized sorghum or as a monoculture to inoculate untreated sorghum, thus yielding *ting* with better sensory properties and higher microbiological quality than that prepared by natural fermentation. By setting these initial conditions, defined inocula can be considered a means whereby *ting* of more reliable and better quality can be produced and it may possibly be exploited to produce different varieties of *ting*.

4.5 REFERENCES

Adams, M.R. and Nicolaides, L. (1997). Review of the sensitivity of different pathogens to fermentation. *Food Control*, **8**, 227-239.

- Akinrele, I.A.** (1970). Fermentation studies on maize during the preparation of a traditional African starch-cake food. *Journal of the Science of Food and Agriculture*, **21**, 619-625.
- Alexander, M.** (1971). *Microbial ecology*. John Wiley and Sons, Inc., London, UK.
- Ammor, M.S. and Mayo, B.** (2007). Selection criteria for lactic acid bacteria to be used as starter cultures in dry sausage production: An update. *Meat Science*, **76**, 138-146.
- Arnold, K.W. and Kaspar, C.W.** (1995). Starvation and stationary phase-induced acid-tolerance in *Escherichia coli* 0157:H7. *Applied and Environmental Microbiology*, **61**, 2037-2039.
- Arvanitidou, M., Tsakris, A., Sofianou, D. and Katsouyannopolous, V.** (1998). Antimicrobial resistance and R-factor transfer of salmonellae isolated from chicken carcasses in Greek hospitals. *International Journal of Food Microbiology*, **40**, 197-201.
- Axelsson, L.** (2004). Lactic acid bacteria: Classification and physiology. In: *Lactic acid bacteria. Microbiological and functional aspects*. Salimen, S., von Wright, A. and Ouwehand, A. (Eds.), Marcel Dekker, Inc., New York, USA, pp. 1-66.
- Batish, V.K., Roy, U., Lal, R. and Grover, S.** (1997). Antifungal attributes of lactic acid bacteria - A review. *Critical Reviews in Biotechnology*, **17**, 209-225.
- Becker, H., Schaller, G., Von Wiese, W. and Terplan, G.** (1994). *Bacillus cereus* in infant foods and dried milk products. *International Journal of Food Microbiology*, **23**, 1-15.
- Beuchat, L.R.** (1987). Efficacy of media and methods for detecting and enumerating *Campylobacter jejuni* in refrigerated chicken meat. *Applied and Environmental Microbiology*, **50**, 934-939.
- Boling, M.B. and Eisener, N.** (1982). *Bogobe*: Sorghum porridge of Botswana. In: *Proceedings of the International Symposium on Sorghum Grain Quality*. International Crops Research Institute for the Semi-Arid Tropics. Mertin, J.V. (Ed.). Patancheru, India, pp. 32-35.
- Brauman, A., Keleke, S., Malonga, M., Miambi, E. and Ampe, F.** (1996). Microbiological and biochemical characterization of cassava retting, a traditional lactic acid fermentation for *foo-foo* (cassava powder) production. *Applied and Environmental Microbiology*, **62**, 2854-2858.
- Chavan, J.K. and Kadam, S.S.** (1989). Nutritional improvement of cereals by fermentation. *Critical Reviews in Food Science and Nutrition*, **28**, 349-400.

- Corsetti, A., Gobetti, M., Rossi, J. and Damiani, P.** (1998). Antimould activity of sourdough lactic acid bacteria: Identification of a mixture of organic acids produced by *Lactobacillus sanfrancisco* CB1. *Applied Microbiology and Biotechnology*, **50**, 253-256.
- De Vos, W.M. and Hugenholtz, J.** (2004). Engineering metabolic highways in lactococci and other lactic acid bacteria. *Trends in Biotechnology*, **22**, 72-79.
- De Vuyst, L.** (2000). Technology aspects related to the application of functional starter cultures. *Food Technology and Biotechnology*, **38**, 105-112.
- De Vuyst, L. and Neysens, P.** (2005). The sourdough microflora: Biodiversity and metabolic interactions. *Trends in Food Science and Technology*, **16**, 43-56.
- De Vuyst, L. and Vandamme, E.J.** (1994). *Bacteriocins of lactic acid bacteria*. Blackie Academic, London, UK, pp. 91-141.
- Dykes, G., Cloete, T.E. and von Holy, A.** (1991). Quantification of microbial populations associated with the manufacture of vacuum-packaged, smoked vienna sausages. *International Journal of Food Microbiology*, **13**, 239-248.
- Ehrlich, P.** (1956). The requirement of the organism for oxygen. An analytical study with the aid of dyes. In: *Collected papers of Paul Ehrlich*. Himmelweit, F., Marquardt, M. and Dale, H. (Eds.). Pergamon Press, London, UK, pp. 433-496.
- Gassem, M.A.A.** (1999). Study of the microorganisms associated with the fermented bread (*Khamir*) product produced from sorghum in Gizan region, Saudi Arabia. *Journal of Applied Microbiology*, **86**, 221-225.
- Gobetti, M., Corsetti, A. and Rossi, J.** (1994). The sourdough microflora interactions between lactic acid bacteria and yeasts: Metabolism of carbohydrates. *Applied Microbiology and Biotechnology*, **41**, 456-460.
- Gourama, M. and Bullerman, L.B.** (1995). Antimycotic and antiaflatoxigenic effect of lactic acid bacteria - A review. *Journal of Food Protection*, **57**, 1275-1280.
- Gran, H.M., Wetlesen, A., Mutukumira, A.N. and Narvhus, J.A.** (2002). Smallholder dairy processing in Zimbabwe: The production of fermented milk products with particular emphasis on sanitation and microbiological quality. *Food Control*, **13**, 161-168.

Gran, H.M., Wetlesen, A., Mutukumira, A.N., Rukure, G. and Narvhus, J.A. (2003). Occurrence of pathogenic bacteria in raw milk, cultured pasteurized milk and naturally soured raw milk produced at small-scale dairies in Zimbabwe. *Food Control*, **14**, 539-544.

Hammes, W.P., Brandt, M.J., Francis, K.L., Rosenheim, J., Seitter, M.F.H. and Vogelmann, S.A. (2005). Microbial ecology of cereal fermentation. *Trends in Food Science and Technology*, **16**, 4-11.

Health Protection Agency (HPA). (2004). Enumeration of *Bacillus cereus* and other *Bacillus* species. National Standard Method F15, Issue 1.

Holzappel, W.H. (2002). Appropriate starter culture technologies for small-scale fermentation in developing countries. *International Journal of Food Microbiology*, **75**, 197-212.

Holzappel, W.H., Geisen, R. and Schillinger, U. (1995). Biological preservation of foods with reference to protective cultures, bacteriocins and food-grade enzymes. *International Journal of Food Microbiology*, **24**, 343-362.

Hugenholtz, J., Kleerebezem, M., Starrenburg, M., Delcour, J., de Vos, W. and Pascal, H. (2000). *Lactococcus lactis* as a cell factory for high-level diacetyl production. *Applied and Environmental Microbiology*, **66**, 4112-4114.

International Commission on Microbiological Specifications for Foods (ICMSF). (1996). Microorganisms in foods: Microbiological specifications of food pathogens. Blackie Academic, London, UK.

Jacobsen, C.N., Rosenfeldt Nielsen, V., Hayford, A.E., Moller, P.L., Michaelsen, K.F., Paerregaard, A., Sandstrom, B., Tvede, M. and Jakobsen, M. (1999). Screening of probiotic activities of forty-seven strains of *Lactobacillus* spp. by *in vitro* techniques and evaluation of the colonization ability of five selected strains in humans. *Applied and Environmental Microbiology*, **65**, 4949-4956.

Jespersen, L., Nielsen, D.S., Hønholt, S. and Jakobsen, M. (2005). Occurrence and diversity of yeasts involved in fermentation of West African cocoa beans. *FEMS Yeast Research*, **5**, 441-453.

Kandler, O. (1983). Carbohydrate metabolism in lactic acid bacteria. *Antonie Van Leeuwenhoek*, **49**, 209-224.

Keddebe, A., Viljoen, B.C., Gadaga, T.H. and Narvhus, J.A. (2007). The effect of container type on the growth of yeast and lactic acid bacteria during production of *Sethemi*, a South African spontaneously fermented milk. *Food Research International*, **40**, 33-38.

- Kimaryo, V.M., Massawe, G.A., Olasupo, N.A. and Holzapfel, W.H.** (2000). The use of starter culture in the fermentation of cassava for the production of 'kivunde', a traditional Tanzanian food product. *International Journal of Food Microbiology*, **56**, 179-190.
- Kingamkono, R., Sjögren, E., Svanberg, U. and Kaijser, B.** (1994). pH and acidity in lactic-fermenting cereal gruels - effects on viability of enteropathogenic microorganisms. *World Journal of Microbiology and Biotechnology*, **10**, 664-669.
- Kingamkono, R., Sjögren, E., Svanberg, U. and Kaijser, B.** (1995). Inhibition of different strains of enteropathogens in a lactic-fermenting cereal gruel. *World Journal of Microbiology and Biotechnology*, **11**, 299-303.
- Kramer, J.M. and Gilbert, R.J.** (1989). *Bacillus cereus* and other *Bacillus* species. In: *Foodborne Pathogens*. Doyle, M.P. (Ed.). Marcel Dekker Inc., New York, USA, pp. 21-70.
- Kunene, N.F., Hastings, J.W. and von Holy, A.** (1999). Bacterial populations associated with a sorghum-based fermented weaning cereal. *International Journal of Food Microbiology*, **49**, 75-83.
- Lavermicocca, P., Valerio, F. and Visconti, A.** (2003). Antifungal activity of phenyllactic acid against molds isolated from bakery products. *Applied and Environmental Microbiology*, **69**, 634-640.
- Lefyedi, M.L., Marais, G.J., Dutton, M.F. and Taylor, J.R.N.** (2005). The microbial contamination, toxicity and quality of turned and unturned outdoor floor malted sorghum. *Journal of the Institute of Brewing*, **111**, 190-196.
- Leroy, F. and De Vuyst, L.** (2004). Lactic acid bacteria as functional starter cultures for the food fermentation industry. *Trends in Food Science and Technology*, **15**, 67-78.
- Madden, J.M. and McCardell, B.A.** (1989). *Vibrio cholerae*. In: *Foodborne Pathogens*. Doyle, M.P. (Ed.). Marcel Dekker, Inc., New York, USA, pp. 525-542.
- Mante, E.S., Skyi-Dawson, E. and Amoa-Awuba, W.K.** (2003). Antimicrobial interactions of microbial species involved in the fermentation of cassava dough into agbelima with particular reference to the inhibitory effect of lactic acid bacteria on enteropathogens. *International Journal of Food Microbiology*, **89**, 41-50.
- Mazur, L.J. and Kim, J.** (2006). Spectrum of noninfectious health effects from molds. *Pediatrics*, **118**, 1909-1926.

Mensah, P. (1997). Fermentation - the key to food safety assurance in Africa? *Food Control*, **8**, 271-278.

Meroth, B., Walter, J., Hertel, C. and Brandt, M. (2003). Monitoring the bacterial population dynamics in sourdough fermentation processes by using PCR-denaturing gradient gel electrophoresis. *Applied and Environmental Microbiology*, **69**, 475-482.

Messens, W. and De Vuyst, L. (2002). Inhibitory substances produced by lactobacilli isolated from sourdoughs - A review. *International Journal of Food Microbiology*, **72**, 31-43.

Mohammed, S.I., Steenson, L.R. and Kirleis, A.W. (1991). Isolation and characterization of microorganisms associated with the traditional sorghum fermentation for production of Sudanese *kisra*. *Applied and Environmental Microbiology*, **8**, 35-46.

Mugula, J.K., Narvhus, J.A. and Sørhaug, T. (2003a). Use of starter cultures of LAB and yeasts in the preparation of *togwa*, a Tanzanian fermented food. *International Journal of Food Microbiology*, **83**, 307-318.

Mugula, J.K., Nnko, S.A.M., Narvhus, J.A. and Sørhaug, T. (2003b). Microbiological and fermentation characteristics of *togwa*, a Tanzanian fermented food. *International Journal of Food Microbiology*, **80**, 187-199.

Nago, M.C., Hounhouigan, J.D., Akissoe, N., Zanou, E. and Mestres, C. (1998). Characterization of the Beninese traditional *ogi*, a fermented maize slurry: Physicochemical and microbiological aspects. *International Journal of Food Science and Technology*, **33**, 307-315.

Nout, M.J.R. (1991). Ecology of accelerated natural lactic fermentation of sorghum-based weaning infant food formulas. *International Journal of Food Microbiology*, **12**, 217-224.

Nout, M.J.R., Rombouts, F.M. and Havelaar, A. (1989). Effect of accelerated natural lactic fermentation of infant food ingredients on some pathogenic microorganisms. *International Journal of Food Microbiology*, **8**, 351-361.

Odunfa, S.A. and Adeyele, S. (1985). Microbiological changes during the traditional production of *ogi-baba*, a West African fermented sorghum gruel. *Journal of Cereal Science*, **3**, 173-180.

Olasupo, N.A., Olukoya, D.K. and Odunfa, S.A. (1997). Identification of *Lactobacillus* species associated with selected African fermented foods. *Zeitschrift für Naturforschung*, **52**, 105-108.

- Olsen, A., Halm, M. and Jakobsen, M.** (1995). The antimicrobial activity of lactic acid bacteria from fermented maize (*kenkey*) and their interactions during fermentation. *Journal of Applied Bacteriology*, **79**, 506-512.
- Oyewole, O.B.** (1997). Lactic fermented foods in Africa and their benefits. *Food Control*, **8**, 289-297.
- Papamanoli, E., Tzanetakis, N., Litopoulou-Tzanetaki, E., Kotzekidou, P. and Parente, E.** (2003). Characterization of lactic acid bacteria isolated from a Greek dry-fermented sausage in respect of their technological and probiotic properties. *Meat Science*, **65**, 859-867.
- Parente, E. and Ricciardi, A.** (1999). Production, recovery and purification of bacteriocins from lactic acid bacteria. *Applied Microbiology and Biotechnology*, **52**, 628-638.
- Peryam, D.R. and Pilgrim, F.J.** (1957). Hedonic scale method of measuring food preference. *Food Technology*, **11**, 9-14.
- Pitt, J.I.** (1988). A laboratory guide to common *Penicillium* species, Second Edition. Commonwealth Scientific and Industrial Research Organisation, North Ryde, Australia.
- Rabie, C.J. and Lübben, A.** (1984). The mycoflora of sorghum malt. *South African Journal of Botany*, **3**, 251-255.
- Rabie, C.J., Lübben, A., Marais, G.J. and Van Vuuren, H.J.** (1997). Enumeration of fungi in barley. *International Journal of Food Microbiology*, **35**, 117-127.
- Rabie, C.J., Van Rensburg, S.J., Van der Walt, J.J. and Lübben, A.** (1975). Onyalai - the possible involvement of a mycotoxin produced by *Phoma sorghina* in the aetiology. *South African Medical Journal*, **57**, 1647-1650.
- Rehman, S., Paterson, A. and Pigott, J.R.** (2006). Flavour in sourdough breads: A review. *Trends in Food Science and Technology*, **17**, 557-566.
- Rombouts, F.M. and Nout, M.J.R.** (1995). Microbial fermentation in the production of plant foods. *Journal of Applied Bacteriology*, **79**, 108S-117S.
- Rowbury, R.J.** (1995). An assessment of environmental factors influencing acid tolerance and sensitivity in *Escherichia coli*, *Salmonella* spp. and other enterobacteria. *Letters in Applied Microbiology*, **20**, 333-337.
- Roy, U., Batish, V.K., Grover, S. and Neelakantan, S.** (1996). Production of antifungal substance by *Lactococcus lactis* subsp. *lactis* CHD-28.3. *Food Microbiology*, **32**, 27-34.

- Saubois, A., Laforet, E.P., Nepote, M.C. and Wagner, M.L.** (1999). Mycological evaluation of a sorghum grain of Argentina, with emphasis on the characterization of *Fusarium* species. *Food Microbiology*, **16**, 435-445.
- Schiemann, D.A.** (1979). Synthesis of a selective agar medium for *Yersinia enterocolitica*. *Canadian Journal of Microbiology*, **25**, 1298-1304.
- Schillinger, U. and Lücke, F.K.** (1989). Antibacterial activity of *Lactobacillus sake* isolated from meat. *Applied and Environmental Microbiology*, **55**, 1901-1906.
- Schlegel, H.G.** (1984). *General Microbiology*, Sixth Edition, Cambridge University Press, Cambridge, UK, pp. 286-288.
- South African Bureau of Standards.** (1975a). Examination for the presence of viable *Clostridium perfringens* organisms in foods, SABS Method Number 761. South African Bureau of Standards, South Africa.
- South African Bureau of Standards.** (1975b). Examination for the presence of viable *Escherichia coli* I in foods, SABS Method Number 758. South African Bureau of Standards, South Africa.
- South African Bureau of Standards.** (1975c). Examination for the presence of viable *Staphylococcus aureus* organisms in foods, SABS Method Number 760. South African Bureau of Standards, South Africa.
- Steinkraus, K.H.** (1996). *Handbook of indigenous fermented food*, Second Edition, Marcel Dekker, Inc., New York, USA.
- Stone, H. and Sidel, J.L.** (1992). *Sensory evaluation practices*. Academic Press, New York, USA, pp. 336.
- Sutherland, J.P. and Varnam, A.H.** (1977). Methods of isolation of potential importance of *Yersinia enterocolitica* in foods stored at low temperatures. *Journal of Applied Bacteriology*, **43**, 304-305.
- Taylor, W.J.** (1965). Isolation of shigellae. I. Xylose lysine agars: New media for isolation of enteric pathogens. *American Journal of Clinical Pathology*, **44**, 471-475.
- Tomicka, A., Chen, J., Barbut, S. and Griffiths, M.W.** (1997). Survival of bioluminescent *Escherichia coli* 0157:H7 in a model system representing fermented sausage production. *Journal of Food Protection*, **60**, 1487-1492.

van Netten, P., van de Ven, A., Perales, I. and Mossel, D.A.A. (1988). A selective and diagnostic medium for use in the enumeration of *Listeria* spp. in foods. *International Journal of Food Microbiology*, **6**, 187-198.

Vernozy-Rozand, C. (1997). Detection of *Escherichia coli* 0157:H7 and other verotoxin-producing *E. coli* (VTEC) in food. *Journal of Applied Microbiology*, **82**, 537-551.

Wong, H.C. and Chen, Y.L. (1988). Effects of lactic acid bacteria and organic acids on growth and germination of *Bacillus cereus*. *Applied and Environmental Microbiology*, **54**, 2179-2184.



CHAPTER FIVE

GENERAL DISCUSSION AND CONCLUSIONS

The objective of this study was essentially to generate new scientific knowledge regarding the microbial diversity and dynamics during sorghum fermentations, using a systematic approach that combined both culture-dependent and culture-independent techniques. Other aims included evaluating microbial populations present in sorghum before, during and after preparation of *ting* in order to elucidate the role of LAB strains with regards to microbiological safety and acceptability of sensory characteristics of *ting*. This research represents the first comprehensive study investigating sorghum fermentations used to produce *ting*. The information generated is significant not only from an academic perspective, but is also relevant for conservation of indigenous knowledge, which may help to bridge the gap between universities and communities by attempting to upgrade the status of indigenous foods.

One of the specific aims of the study was to characterize six representative LAB isolates from spontaneous sorghum fermentations, using a polyphasic approach that encompassed both classical microbiological and DNA-based methods. Sequencing of genes encoding *pheS* clearly assigned the LAB to *Lb. plantarum*, *Lb. rhamnosus* and *Lb. fermentum*. Although sequencing of the 16S rRNA-encoding genes gave similar results, *Lb. plantarum* could not be clearly distinguished from its close phylogenetic neighbours, *Lb. paraplantarum* and *Lb. pentosus*. Likewise, 16S rRNA gene sequencing failed to distinguish *Lb. rhamnosus* from its close relative, *Lb. casei*. These results indicate that a polyphasic approach, which includes both biochemical and DNA-based techniques, should be followed for robust genotypic and phenotypic characterization of LAB obtained from spontaneous sorghum fermentations.

Another aim of this study was to systematically and effectively describe the microbial populations during spontaneous fermentations of two sorghum types at 20, 25 and 30°C using culture-independent PCR-DGGE, as well as culture-dependent methods aided by DNA-based techniques (sequencing of genes encoding 16S rRNA and *pheS*). *Lb. curvatus* was not detected in culture media, but it was predominant in DGGE gels. This is probably due to these bacteria having entered a viable but non-cultivable state, which is characterized by metabolically active cells that do not produce colonies on both selective and non-selective media (Giraffa and Neviani, 2001). This demonstrates one of the virtues of culture-independent approaches over culture-dependent methods (Muyzer and Smalla, 1998; Giraffa and Neviani, 2001). In contrast, *Lb. fermentum*, *Lb. plantarum*, *Lb. rhamnosus*, *E. faecalis* and *E. mundtii* were all isolated in culture media, but they were absent in DGGE gels. This

might be due to either the bacteria occurring in numbers below the detection limits of PCR-DGGE (Muyzer and Smalla, 1998), and/or due to biases introduced by DNA extraction and PCR (Ercolini, 2004; De Vero *et al.*, 2006; Camu *et al.*, 2007). Indeed, related studies showed that PCR-DGGE was unable to detect some bacterial species that were isolated on culture media (Meroth *et al.*, 2003; Ercolini *et al.*, 2004). Therefore, the combined approach used in this study was effective in revealing the microbial diversity and dynamics during spontaneous sorghum fermentations.

Other aims of this study included assessing bacterial, yeast and mould populations found in sorghum before, during and after preparation of *ting*, and clarifying the role of selected LAB strains with regards to their contribution to the microbiological safety and acceptability of sensory characteristics of *ting*. Commercial sorghum powder was shown to harbour microorganisms such as LAB, *Bacillus* spp., *Enterobacteriaceae*, yeasts and moulds. Natural fermentation alone did not completely reduce the numbers of pathogens such as *Enterobacteriaceae* members and *Bacillus* spp. below the detection limit, despite the low pH. This may be due to the presence of acid-resistant strains or microenvironments within the fermented sorghum slurries that allowed these bacteria to survive. However, the addition of a *L. lactis* starter culture to untreated sorghum reduced the numbers of *B. cereus*, *Enterobacteriaceae* and moulds to undetectable levels, illustrating the potential that this strain has in improving the microbiological safety of *ting*. Indeed, the addition of *L. lactis* to untreated sorghum resulted in *ting* that was most preferred by panellists. This may be due to the ability of *L. lactis* to control fermentation, as illustrated by its dominance over other LAB in DGGE gels and its ability to reduce the growth of other LAB in antimicrobial studies. For this reason, *L. lactis* may be used either as a monoculture to inoculate untreated sorghum, or in co-culture with *Lb. plantarum* for use in the fermentation of sterilized sorghum, thus yielding *ting* with better sensory properties than that prepared by natural fermentation. These LAB may possibly be exploited to produce different varieties of *ting* in future.

REFERENCES

Camu, N., De Winter, T., Verbrugghe, K., Cleenwerck, I., Vandamme, P., Takrama, J.S., Vancanneyt, M. and De Vuyst, L. (2007). Dynamics and biodiversity of populations of lactic acid bacteria involved in spontaneous heap fermentation of cocoa beans in Ghana. *Applied and Environmental Microbiology*, **73**, 1809-1824.

De Vero, L., Gala, E., Gullo, M., Solieri, L., Landi, S. and Giudici, P. (2006). Application of denaturing gradient gel electrophoresis (DGGE) analysis to evaluate acetic acid bacteria in traditional balsamic vinegar. *Food Microbiology*, **23**, 809-813.

Ercolini, D. (2004). PCR-DGGE fingerprinting: Novel strategies for detection of microbes in food. *Journal of Microbiological Methods*, **56**, 297-314.

Ercolini, D., Mauriello, G., Blaiotta, G., Moschetti, G. and Coppola, S. (2004). PCR-DGGE fingerprints of microbial succession during a manufacture of traditional water buffalo mozzarella cheese. *Journal of Applied Microbiology*, **96**, 263-270.

Giraffa, G. and Neviani, E. (2001). DNA-based, culture-independent strategies for evaluating microbial communities in food-associated ecosystems. *International Journal of Food Microbiology*, **67**, 19-34.

Muyzer, G. and Smalla, K. (1998). Application of denaturing gradient gel electrophoresis (DGGE) and temperature gradient gel electrophoresis (TGGE) in microbial ecology. *Antonie Van Leeuwenhoek*, **73**, 127-141.

Meroth, C.B., Walter, J., Hertel, C., Brandt, M.J. and Hammes, W.P. (2003). Monitoring the bacterial population dynamics in sourdough fermentation processes by using PCR-denaturing gradient gel electrophoresis. *Applied and Environmental Microbiology*, **69**, 475-482.



APPENDICES

Appendix A - Current contents of the order *Lactobacillales*, showing all valid species in the genera associated with food technology.....145

Appendix B - Current contents of the order *Lactobacillales*, showing all valid species other than *Lactacillaceae* family.....147

Appendix C - Multiple *pheS* gene sequence alignment for LAB isolated from natural sorghum fermentations and type strains (Chapter 3).....150

Appendix D - Multiple 16S rRNA gene sequence alignment for LAB isolated from natural sorghum fermentations and type strains (Chapter 3).....152

Appendix E - Multiple 16S rRNA gene sequence alignment for *Lactobacillus* and *Weissella* species isolated from natural sorghum fermentations and type strains (Chapter 4).....158

Appendix F - Multiple *pheS* gene sequence alignment for *Lactobacillus* species isolated from spontaneous sorghum fermentations and type strains (Chapter 4).....166

Appendix G - Multiple 16S rRNA gene sequence alignment for *Lactococcus* and *Enterococcus* species isolated from spontaneous sorghum fermentations and type strains (Chapter 4).....167

Appendix H - References.....172

Appendix A - Current contents of the order *Lactobacillales*, showing all valid species in the genera associated with food technology.

Genus	Species	Author	Phylogenetic group ¹	Metabolism ²
<i>Lactobacillus</i>	<i>acetotolerans</i>	Entani <i>et al.</i> , 1986	<i>Lb. delbrueckii</i>	B
<i>Lactobacillus</i>	<i>acidifarinae</i>	Vancanneyt <i>et al.</i> , 2005	<i>Lb. buchneri</i>	C
<i>Lactobacillus</i>	<i>acidipiscis</i>	Tanasupawat <i>et al.</i> , 2000	<i>Lb. salivarius</i>	B
<i>Lactobacillus</i>	<i>acidophilus</i>	(Moro, 1900) Hansen and Mocoquot, 1970	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>agilis</i>	Weiss <i>et al.</i> , 1982	<i>Lb. salivarius</i>	B
<i>Lactobacillus</i>	<i>algidus</i>	Kato <i>et al.</i> , 2000	<i>Lb. salivarius</i> -single species	B
<i>Lactobacillus</i>	<i>alimentarius</i>	(Ex Reuter, 1970) Reuter, 1983	<i>Lb. alimentarius</i> - <i>Lb. farciminis</i>	B
<i>Lactobacillus</i>	<i>amyolyticus</i>	Bohak <i>et al.</i> , 1999	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>amylophilus</i>	Nakamura and Crowell, 1981	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>amylotrophicus</i>	Naser <i>et al.</i> , 2006	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>amylovorus</i>	Nakamura, 1981	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>animalis</i>	Dent and Williams, 1983	<i>Lb. salivarius</i>	A
<i>Lactobacillus</i>	<i>antri</i>	Roos <i>et al.</i> , 2005	<i>Lb. reuteri</i>	C
<i>Lactobacillus</i>	<i>apodemi</i>	Osawa <i>et al.</i> , 2006	<i>Lb. salivarius</i>	B
<i>Lactobacillus</i>	<i>arizonensis</i>	Swezey <i>et al.</i> , 2000 ^a		
<i>Lactobacillus</i>	<i>aviaries</i>	Fujisawa <i>et al.</i> , 1985	<i>Lb. salivarius</i>	A
<i>Lactobacillus</i>	<i>aviarius</i> subsp. <i>araffinosus</i>	Fujisawa <i>et al.</i> , 1986	<i>Lb. salivarius</i>	A
<i>Lactobacillus</i>	<i>aviarius</i> subsp. <i>aviaries</i>	Fujisawa <i>et al.</i> , 1985	<i>Lb. salivarius</i>	A
<i>Lactobacillus</i>	<i>bavaricus</i>	Stetter and Stetter, 1980		
<i>Lactobacillus</i>	<i>bifermentans</i>	Kandler <i>et al.</i> , 1983	<i>Lb. coryneformis</i>	B
<i>Lactobacillus</i>	<i>brevis</i>	(Orla-Jensen, 1919) Bergey <i>et al.</i> , 1934	<i>Lb. brevis</i>	C
<i>Lactobacillus</i>	<i>buchneri</i>	(Henneberg, 1903) Bergey <i>et al.</i> , 1923	<i>Lb. buchneri</i>	C
<i>Lactobacillus</i>	<i>casei</i>	(Orla-Jensen, 1916) Hansen and Lessel, 1971	<i>Lb. casei</i>	B
<i>Lactobacillus</i>	<i>catenaformis</i>	(Eggert, 1935) Moore and Holdeman, 1970		A
<i>Lactobacillus</i>	<i>coleshominis</i>	Nikolaitchouk <i>et al.</i> , 2001	<i>Lb. reuteri</i>	B
<i>Lactobacillus</i>	<i>collinooides</i>	Carr and Davis, 1972		C
<i>Lactobacillus</i>	<i>composti</i>	Endo and Okada, 2007	<i>Lb. coryneformis</i>	B
<i>Lactobacillus</i>	<i>concaus</i>	Tong and Dong, 2005	<i>P. dextrinicus</i>	A
<i>Lactobacillus</i>	<i>coryniformis</i> subsp. <i>coryniformis</i>	Abo-Elnaga and Kandler, 1965	<i>Lb. coryniformis</i>	B
<i>Lactobacillus</i>	<i>coryniformis</i> subsp. <i>torquens</i>	Abo-Elnaga and Kandler, 1965	<i>Lb. coryniformis</i>	B
<i>Lactobacillus</i>	<i>crispatus</i>	(Brygoo and Aladame, 1953) Moore and Holdeman, 1970	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>curvatus</i>	(Troili-Petersson, 1903) Abo-Elnaga and Kandler, 1965; Klein <i>et al.</i> , 1996	<i>Lb. sakei</i>	B
<i>Lactobacillus</i>	<i>curvatus</i>	(Troili-Petersson, 1903) Abo-Elnaga and Kandler, 1965 emend. Klein <i>et al.</i> , 1996	<i>Lb. sakei</i>	B
<i>Lactobacillus</i>	<i>delbrueckii</i> subsp. <i>bulgaricus</i>	(Orla-Jensen, 1919) Weiss <i>et al.</i> , 1984	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>delbrueckii</i> subsp. <i>delbrueckii</i>	(Leichmann, 1896) Beijerinck, 1901	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>delbrueckii</i> subsp. <i>indicus</i>	Dellaglio <i>et al.</i> , 2005	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>delbrueckii</i> subsp. <i>lactis</i>	(Orla-Jensen, 1919) Weiss <i>et al.</i> , 1984	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>diolivorans</i>	Krooneman <i>et al.</i> , 2002	<i>Lb. buchneri</i>	C
<i>Lactobacillus</i>	<i>equi</i>	Morotomi <i>et al.</i> , 2002	<i>Lb. salivarius</i>	A
<i>Lactobacillus</i>	<i>farciminis</i>	(Ex Reuter, 1970) Reuter, 1983	<i>Lb. alimentarius</i> - <i>Lb. farciminis</i>	A
<i>Lactobacillus</i>	<i>farraginis</i>	Endo and Okada, 2007	<i>Lb. buchneri</i>	B
<i>Lactobacillus</i>	<i>fermentum</i>	Beijerinck, 1901 emend. Dellaglio <i>et al.</i> , 2004	<i>Lb. reuteri</i>	C
<i>Lactobacillus</i>	<i>formicalis</i>	Dicks <i>et al.</i> , 2000	<i>Lb. delbrueckii</i>	B
<i>Lactobacillus</i>	<i>fructivorans</i>	Charlton <i>et al.</i> , 1934	<i>Lb. fructivorans</i>	C
<i>Lactobacillus</i>	<i>frumenti</i>	Müller <i>et al.</i> , 2000	<i>Lb. reuteri</i>	C
<i>Lactobacillus</i>	<i>fuchuensis</i>	Sakala <i>et al.</i> , 2002	<i>Lb. sakei</i>	B
<i>Lactobacillus</i>	<i>gallinarum</i>	Fujisawa <i>et al.</i> , 1992	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>gasseri</i>	Lauer and Kandler, 1980	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>gastricus</i>	Roos <i>et al.</i> , 2005	<i>Lb. reuteri</i>	C
<i>Lactobacillus</i>	<i>graminis</i>	Beck <i>et al.</i> , 1989	<i>Lb. sakei</i>	B
<i>Lactobacillus</i>	<i>hammesii</i>	Valcheva <i>et al.</i> , 2005	<i>Lb. brevis</i>	B
<i>Lactobacillus</i>	<i>hamsteri</i>	Mitsuoka and Fujisawa, 1988	<i>Lb. delbrueckii</i>	B
<i>Lactobacillus</i>	<i>harbinensis</i>	Miyamoto <i>et al.</i> , 2006	<i>Lb. perolens</i>	B
<i>Lactobacillus</i>	<i>helveticus</i>	(Orla-Jensen, 1919) Bergey <i>et al.</i> , 1925	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>hilgardii</i>	Douglas and a Cruess, 1936	<i>Lb. buchneri</i>	C
<i>Lactobacillus</i>	<i>homohiochii</i>	Kitahara <i>et al.</i> , 1957	<i>Lb. fructivorans</i>	B
<i>Lactobacillus</i>	<i>iners</i>	Falsen <i>et al.</i> , 1999	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>ingluviei</i>	Baele <i>et al.</i> , 2003	<i>Lb. reuteri</i>	C
<i>Lactobacillus</i>	<i>intestinalis</i>	(ex Hemme, 1974) Fujisawa <i>et al.</i> , 1990	<i>Lb. delbrueckii</i>	B
<i>Lactobacillus</i>	<i>jensenii</i>	Gasser <i>et al.</i> , 1970	<i>Lb. delbrueckii</i>	B
<i>Lactobacillus</i>	<i>johnsonii</i>	Fujisawa <i>et al.</i> , 1992	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>kalixensis</i>	Roos <i>et al.</i> , 2005	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>kefiranoformis</i> subsp. <i>kefiranoformis</i>	Fujisawa <i>et al.</i> , 1988	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>kefiranoformis</i> subsp. <i>kefirgranum</i>	(Takizawa <i>et al.</i> , 1994) Vancanneyt <i>et al.</i> , 2004	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>kefiri</i>	Kandler and Kunath, 1983	<i>Lb. buchneri</i>	C
<i>Lactobacillus</i>	<i>kimchii</i>	Yoon <i>et al.</i> , 2000	<i>Lb. alimentarius</i> - <i>Lb. farciminis</i>	B
<i>Lactobacillus</i>	<i>kitasatonis</i>	Mukai <i>et al.</i> , 2003	<i>Lb. delbrueckii</i>	B
<i>Lactobacillus</i>	<i>kunkeei</i>	Edwards <i>et al.</i> , 1998	Single species	C
<i>Lactobacillus</i>	<i>lindneri</i>	Back <i>et al.</i> , 1997	<i>Lb. fructivorans</i>	C
<i>Lactobacillus</i>	<i>malefermentans</i>	Farrow <i>et al.</i> , 1989	Single species	C



Genus	Species	Author	Phylogenetic group ¹	Metabolism ²
<i>Lactobacillus</i>	<i>mali</i>	Carr and Davies, 1970, Kaneuchi <i>et al.</i> , 1988	<i>Lb. salivarius</i>	A
<i>Lactobacillus</i>	<i>manihotivorans</i>	Morlon-Guyet <i>et al.</i> , 1998	Couple3	A
<i>Lactobacillus</i>	<i>mindensis</i>	Ehrmann <i>et al.</i> , 2003	<i>Lb. alimentarius-Lb. farciminis</i>	A
<i>Lactobacillus</i>	<i>mucosae</i>	Roos <i>et al.</i> , 2000	<i>Lb. reuteri</i>	C
<i>Lactobacillus</i>	<i>murinus</i>	Hemme <i>et al.</i> , 1982	<i>Lb. salivarius</i>	B
<i>Lactobacillus</i>	<i>nagelii</i>	Edwards <i>et al.</i> , 2000	<i>Lb. salivarius</i>	A
<i>Lactobacillus</i>	<i>namurensis</i>	Scheirlinck <i>et al.</i> , 2007	<i>Lb. buchneri</i>	C
<i>Lactobacillus</i>	<i>oris</i>	Farrow and Collins, 1988	<i>Lb. reuteri</i>	C
<i>Lactobacillus</i>	<i>panis</i>	Wiese <i>et al.</i> , 1996	<i>Lb. reuteri</i>	C
<i>Lactobacillus</i>	<i>nantensis</i>	Valcheva <i>et al.</i> , 2006	<i>Lb. alimentarius-Lb. farciminis</i>	B
<i>Lactobacillus</i>	<i>pantheris</i>	Liu and Dong, 2002	Single species	A
<i>Lactobacillus</i>	<i>parabrevis</i>	Vancanneyt <i>et al.</i> , 2006	<i>Lb. brevis</i>	C
<i>Lactobacillus</i>	<i>parabuchneri</i>	Farrow <i>et al.</i> , 1989	<i>Lb. buchneri</i>	C
<i>Lactobacillus</i>	<i>paracasei</i> subsp. <i>paracasei</i>	Collins <i>et al.</i> , 1989	<i>Lb. casei</i>	B
<i>Lactobacillus</i>	<i>paracasei</i> subsp. <i>tolerans</i>	Collins <i>et al.</i> , 1989	<i>Lb. casei</i>	B
<i>Lactobacillus</i>	<i>paracollinoides</i>	Suzuki <i>et al.</i> , 2004	<i>Lb. perolens</i>	C
<i>Lactobacillus</i>	<i>parafarraginis</i>	Endo and Okada, 2007	<i>Lb. buchneri</i>	B
<i>Lactobacillus</i>	<i>parakefiri</i>	Takizawa <i>et al.</i> , 1994	<i>Lb. buchneri</i>	C
<i>Lactobacillus</i>	<i>paralimentarius</i>	Cai <i>et al.</i> , 1999	<i>Lb. alimentarius-Lb. farciminis</i>	B
<i>Lactobacillus</i>	<i>paraplantarum</i>	Curk <i>et al.</i> , 1996	<i>Lb. plantarum</i>	B
<i>Lactobacillus</i>	<i>pentosus</i>	Zanoni <i>et al.</i> , 1997	<i>Lb. plantarum</i>	B
<i>Lactobacillus</i>	<i>perolens</i>	Back <i>et al.</i> , 2000	<i>Lb. perolens</i>	B
<i>Lactobacillus</i>	<i>plantarum</i>	(Orla-Jensen, 1919) Bergey <i>et al.</i> , 1923	<i>Lb. plantarum</i>	B
<i>Lactobacillus</i>	<i>platarum</i> subsp. <i>argenteratensis</i>	Bringel <i>et al.</i> , 2005	<i>Lb. plantarum</i>	B
<i>Lactobacillus</i>	<i>pontis</i>	Vogel <i>et al.</i> , 1994	<i>Lb. reuteri</i>	C
<i>Lactobacillus</i>	<i>pscittasi</i>	Lawson <i>et al.</i> , 2001	<i>Lb. delbrueckii</i>	C
<i>Lactobacillus</i>	<i>rennini</i>	Chenoll <i>et al.</i> , 2006	<i>Lb. coryneformis</i>	B
<i>Lactobacillus</i>	<i>reuteri</i>	Kandler <i>et al.</i> , 1982	<i>Lb. reuteri</i>	C
<i>Lactobacillus</i>	<i>rhamnosus</i>	(Hansen, 1968) Collins <i>et al.</i> , 1989	<i>Lb. casei</i>	B
<i>Lactobacillus</i>	<i>rogosae</i>	Holdeman and Moore, 1974	Strain not available	
<i>Lactobacillus</i>	<i>rossiae</i>	Corsetti <i>et al.</i> , 2005	Couple1	C
<i>Lactobacillus</i>	<i>ruminis</i>	Sharpe <i>et al.</i> , 1973	<i>Lb. salivarius</i>	A
<i>Lactobacillus</i>	<i>saerimneri</i>	Pedersen and Roos, 2004	<i>Lb. salivarius</i>	A
<i>Lactobacillus</i>	<i>sakei</i> subsp. <i>carnosus</i>	Torriani <i>et al.</i> , 1996	<i>Lb. sakei</i>	B
<i>Lactobacillus</i>	<i>sakei</i> subsp. <i>sakei</i>	Katagiri <i>et al.</i> , 1934 emend. Klein <i>et al.</i> , 1996	<i>Lb. sakei</i>	B
<i>Lactobacillus</i>	<i>salivarius</i>	Rogosa <i>et al.</i> , 1953, Li <i>et al.</i> , 2006	<i>Lb. salivarius</i>	A
<i>Lactobacillus</i>	<i>sanfranciscensis</i>	Weiss and Schillinger, 1984	<i>Lb. fructivorans</i>	C
<i>Lactobacillus</i>	<i>satsumensis</i>	Endo and Okada, 2005	<i>Lb. salivarius</i>	A
<i>Lactobacillus</i>	<i>secaliphilus</i>	Ehrmann <i>et al.</i> , 2007	<i>Lb. reuteri</i>	B
<i>Lactobacillus</i>	<i>sharpeae</i>	Weiss <i>et al.</i> , 1982	Single species	A
<i>Lactobacillus</i>	<i>siliginis</i>	Aslam <i>et al.</i> , 2006	Couple1	C
<i>Lactobacillus</i>	<i>sobrius</i>	Kostantinov <i>et al.</i> , 2006	<i>Lb. delbrueckii</i>	B
<i>Lactobacillus</i>	<i>spicheri</i>	Meroth <i>et al.</i> , 2004	<i>Lb. buchneri</i>	B
<i>Lactobacillus</i>	<i>suebicus</i>	Kleymans <i>et al.</i> , 1989	Couple2	C
<i>Lactobacillus</i>	<i>tucceti</i>		<i>Lb. alimentarius-Lb. farciminis</i>	A
<i>Lactobacillus</i>	<i>ultunensis</i>	Roos <i>et al.</i> , 2005	<i>Lb. delbrueckii</i>	A
<i>Lactobacillus</i>	<i>vaccinostercus</i>	(Kozaki and Okada, 1983) Okada <i>et al.</i> , 1979	Couple2	C
<i>Lactobacillus</i>	<i>vaginalis</i>	Embley <i>et al.</i> , 1989	<i>Lb. reuteri</i>	C
<i>Lactobacillus</i>	<i>versmoldensis</i>	Kröckel <i>et al.</i> , 2003	<i>Lb. alimentarius-Lb. farciminis</i>	A
<i>Lactobacillus</i>	<i>vini</i>	Rodas <i>et al.</i> , 2006	<i>Lb. salivarius</i>	B
<i>Lactobacillus</i>	<i>vitulinus</i>	Sharpe <i>et al.</i> , 1973		A
<i>Lactobacillus</i>	<i>zeae</i>	Dicks <i>et al.</i> , 1996	<i>Lb. casei</i>	B
<i>Lactobacillus</i>	<i>zymae</i>	Vancanneyt <i>et al.</i> , 2005	<i>Lb. buchneri</i>	C
<i>Paralactobacillus</i>	<i>selangonsis</i>	Leisner <i>et al.</i> , 2000	ND	ND
<i>Pediococcus</i>	<i>acidilactici</i>	Lindner, 1887	ND	ND
<i>Pediococcus</i>	<i>cellicola</i>	Zhang <i>et al.</i> , 2005	ND	ND
<i>Pediococcus</i>	<i>halophilus</i>	Mees, 1934	ND	ND
<i>Pediococcus</i>	<i>inopinatus</i>	Back, 1988	ND	ND
<i>Pediococcus</i>	<i>parvulus</i>	Gunther and White, 1961	ND	ND
<i>Pediococcus</i>	<i>pentosaceus</i>	Mees, 1934	ND	ND
<i>Pediococcus</i>	<i>stilesii</i>	Franz <i>et al.</i> , 2006	ND	ND
<i>Pediococcus</i>	<i>urinaeequi</i>	(ex Mees, 1934) Garvie, 1988	ND	ND

^{1,2} The phylogenetic groups and metabolism of *Lactobacillus* genera are from Felis and Dellaglio, 2007. A: obligately homofermentative; B: facultatively heterofermentative, and C: obligately heterofermentative. ¹*Lactobacillus arizonensis* is a later heterotypic synonym of *Lactobacillus plantarum* (Kostinek *et al.*, 2005).

Appendix B - Current contents of the order *Lactobacillales*, showing all valid species other than *Lactacillaceae* family.

Family	Genus	Species	Author	
<i>Aerococcaceae</i>	<i>Aerococcus</i>	<i>sanguinicola</i>	Lawson <i>et al.</i> , 2001	
	<i>Aerococcus</i>	<i>urinae</i>	Aguirre and Collins, 1992	
	<i>Aerococcus</i>	<i>urinaeequi</i>	(Garvie, 1988) Felis <i>et al.</i> , 2005	
	<i>Aerococcus</i>	<i>urinaehominis</i>	Lawson <i>et al.</i> , 2001	
	<i>Aerococcus</i>	<i>viridans</i>	Williams <i>et al.</i> , 1953	
	<i>Abiotrophia</i>	<i>defectiva</i>	(Bouvet <i>et al.</i> , 1989) Kawamura <i>et al.</i> , 1995	
	<i>Abiotrophia</i>	<i>adiacens</i>	(Bouvet <i>et al.</i> , 1989) Kawamura <i>et al.</i> , 1995	
	<i>Abiotrophia</i>	<i>balaenopterae</i>	Lawson <i>et al.</i> , 1999	
	<i>Abiotrophia</i>	<i>elegans</i>	Roggenkamp <i>et al.</i> , 1999	
	<i>Dolosicoccus</i>	<i>paucivorans</i>	Collins <i>et al.</i> , 1999	
	<i>Eremococcus</i>	<i>coleocora</i>	Collins <i>et al.</i> , 1999	
	<i>Facklamia</i>	<i>hominis</i>	Collins <i>et al.</i> , 1997	
	<i>Facklamia</i>	<i>Ignava</i>	Collins <i>et al.</i> , 1998	
	<i>Facklamia</i>	<i>languida</i>	Lawson <i>et al.</i> , 1999	
	<i>Facklamia</i>	<i>miroungae</i>	Hoyles <i>et al.</i> , 2001	
	<i>Facklamia</i>	<i>sourekii</i>	Collins <i>et al.</i> , 1999	
	<i>Facklamia</i>	<i>tabacinasalis</i>	Collins <i>et al.</i> , 1999	
	<i>Globicatella</i>	<i>sanguinis</i>	Collins <i>et al.</i> , 1995	
	<i>Globicatella</i>	<i>salfidifaciens</i>	Vandamme <i>et al.</i> , 2001	
	<i>Carnobacteriaceae</i>	<i>Ignavigranum</i>	<i>ruoffiae</i>	Collins <i>et al.</i> , 1999
<i>Agitococcus</i>		<i>lubricus</i>	Franzmann and Skermann, 1981	
<i>Alkalibacterium</i>		<i>olivapovliticus</i>	Ntougias and Russel, 2001	
<i>Alkalibacterium</i>		<i>iburiense</i>	Nakajima <i>et al.</i> , 2005	
<i>Alkalibacterium</i>		<i>psychrotolerans</i>	Yumoto <i>et al.</i> , 2004	
<i>Allofustis</i>		<i>seminis</i>	Collins <i>et al.</i> , 2003	
<i>Alloiococcus</i>		<i>otitis</i>	Aguirre and Collins, 1992	
<i>Desemzia</i>		<i>incerta</i>	(Steinhaus, 1941) Stackebrandt <i>et al.</i> , 1999	
<i>Dolosigranulum</i>		<i>pigrum</i>	Aguirre <i>et al.</i> , 1994	
<i>Granulicatella</i>		<i>adiacens</i>	(Bouvet <i>et al.</i> , 1989) Collins and Lawson, 2000	
<i>Granulicatella</i>		<i>balaenopterae</i>	(Lawson <i>et al.</i> , 1999) Collins and Lawson, 2000	
<i>Isobaculum</i>		<i>melis</i>	Collins <i>et al.</i> , 2002	
<i>Lactosphaera</i>		<i>pasteurii</i>	(Schink, 1985) Janssen <i>et al.</i> , 1995	
<i>Marinilactibacillus</i>		<i>psychrotolerans</i>	Ishikawa <i>et al.</i> , 2003	
<i>Trichococcus</i>		<i>collinsii</i>	Liu <i>et al.</i> , 2002	
<i>Trichococcus</i>		<i>flocculiformis</i>	Scheff <i>et al.</i> , 1984	
<i>Trichococcus</i>		<i>pasteurii</i>	(Schink, 1985) Liu <i>et al.</i> , 2002	
<i>Trichococcus</i>		<i>patagoniensis</i>	Pikuta <i>et al.</i> , 2006	
<i>Carnobacterium</i>		<i>alterfunditum</i>	Franzmann <i>et al.</i> , 1993	
<i>Carnobacterium</i>		<i>divergens</i>	(Holzapfel and Gerber, 1984) Collins <i>et al.</i> , 1987	
<i>Carnobacterium</i>		<i>funditum</i>	Franzmann <i>et al.</i> , 1993	
<i>Carnobacterium</i>		<i>gallinarum</i>	Collins <i>et al.</i> , 1987	
<i>Carnobacterium</i>		<i>inhibens</i>	Jöborn <i>et al.</i> , 1999	
<i>Carnobacterium</i>		<i>maltomaticum</i>	(Miller <i>et al.</i> , 1974) Mora <i>et al.</i> , 2003	
<i>Carnobacterium</i>		<i>mobile</i>	Collins <i>et al.</i> , 1987	
<i>Carnobacterium</i>		<i>piscicola</i>	(Hiu <i>et al.</i> , 1984) Collins <i>et al.</i> , 1987	
<i>Carnobacterium</i>		<i>pleistocenum</i>	Pikuta <i>et al.</i> , 2005	
<i>Carnobacterium</i>		<i>viridans</i>	Holley <i>et al.</i> , 2002	
<i>Leuconostocaceae</i>		<i>Leuconostoc</i>	<i>amelibiosum</i>	Schillinger <i>et al.</i> , 1989
		<i>Leuconostoc</i>	<i>argentinum</i>	Dicks <i>et al.</i> , 1993
		<i>Leuconostoc</i>	<i>carnosum</i>	Shaw and Harding, 1989
		<i>Leuconostoc</i>	<i>fallax</i>	Martinez-Murcia and Collins, 1992
	<i>Leuconostoc</i>	<i>ficulneums</i>	Antunes <i>et al.</i> , 2002	
	<i>Leuconostoc</i>	<i>fructosum</i>	Antunes <i>et al.</i> , 2002	
	<i>Leuconostoc</i>	<i>gelidum</i>	Shaw and Harding, 1989	
	<i>Leuconostoc</i>	<i>gasicomitatum</i>	Björkroth <i>et al.</i> , 2001	
	<i>Leuconostoc</i>	<i>inhae</i>	Kim <i>et al.</i> , 2003	
	<i>Leuconostoc</i>	<i>kimchii</i>	Kim <i>et al.</i> , 2000	
	<i>Leuconostoc</i>	<i>mesenteroides</i> subsp. <i>cremoris</i>	(Knudsen and Sørensen, 1929) Garvie, 1983	
	<i>Leuconostoc</i>	<i>mesenteroides</i> subsp. <i>dextranicum</i>	(Beijerinck, 1912) Garvie, 1983	
	<i>Leuconostoc</i>	<i>paramesenteroides</i>	Garvie, 1967	
	<i>Leuconostoc</i>	<i>pseudoficulneum</i>	Chambel <i>et al.</i> , 2006	
	<i>Leuconostoc</i>	<i>pseudomesenteroides</i>	Farrow <i>et al.</i> , 1989	
	<i>Oenococcus</i>	<i>kitaharae</i>	Endo and Okada, 2006	
	<i>Oenococcus</i>	<i>oeni</i>	(Garvie, 1967) Dicks <i>et al.</i> , 1995	
	<i>Weissella</i>	<i>cibaria</i>	Björkroth <i>et al.</i> , 2002	
	<i>Weissella</i>	<i>confusa</i>	(Holzapfel and Knadler, 1969) Collins <i>et al.</i> , 1994	
	<i>Weissella</i>	<i>halotolerans</i>	(Kandler <i>et al.</i> , 1983) Collins <i>et al.</i> , 1994	
	<i>Weissella</i>	<i>hellenica</i>	Collins <i>et al.</i> , 1994	
	<i>Weissella</i>	<i>kandleri</i>	(Holzapfel and van Wyk, 1983) Collins <i>et al.</i> , 1994	
	<i>Weissella</i>	<i>kimchii</i>	Choi <i>et al.</i> , 2002	
	<i>Weissella</i>	<i>koreensis</i>	Lee <i>et al.</i> , 2002	
	<i>Weissella</i>	<i>minor</i>	(Kandler <i>et al.</i> , 1983) Collins <i>et al.</i> , 1994	
	<i>Weissella</i>	<i>paramesenteroides</i>	(Garvie, 1967) Collins <i>et al.</i> , 1994	
	<i>Weissella</i>	<i>solii</i>	Magnusson <i>et al.</i> , 2002	
	<i>Weissella</i>	<i>viridescens</i>	(Niven and Evans, 1957) Collins <i>et al.</i> , 1994	



Family	Genus	Species	Author	
Enterococcaceae	<i>Enterococcus</i>	<i>aquimarinus</i>	Svec <i>et al.</i> , 2005	
	<i>Enterococcus</i>	<i>asini</i>	De Vaux <i>et al.</i> , 1998	
	<i>Enterococcus</i>	<i>avium</i>	(ex Nowlan and Deibel, 1967) Collins <i>et al.</i> , 1984	
	<i>Enterococcus</i>	<i>caccae</i>	Carvalho <i>et al.</i> , 2006	
	<i>Enterococcus</i>	<i>canintestini</i>	Naser <i>et al.</i> , 2005	
	<i>Enterococcus</i>	<i>asini</i>	De Vaux <i>et al.</i> , 1998	
	<i>Enterococcus</i>	<i>avium</i>	(ex Nowlan and Deibel, 1967) Collins <i>et al.</i> , 1984	
	<i>Enterococcus</i>	<i>caccae</i>	Carvalho <i>et al.</i> , 2006	
	<i>Enterococcus</i>	<i>canintestini</i>	Naser <i>et al.</i> , 2005	
	<i>Enterococcus</i>	<i>canis</i>	De Graef <i>et al.</i> , 2003	
	<i>Enterococcus</i>	<i>cecorum</i>	(Devriese <i>et al.</i> , 1983) Williams <i>et al.</i> , 1989	
	<i>Enterococcus</i>	<i>columbae</i>	Devriese <i>et al.</i> , 1993	
	<i>Enterococcus</i>	<i>devriesei</i>	Svec <i>et al.</i> , 2005	
	<i>Enterococcus</i>	<i>dispar</i>	Collins <i>et al.</i> , 1991	
	<i>Enterococcus</i>	<i>durans</i>	(ex Sherman and Wing, 1973) Collins <i>et al.</i> , 1984	
	<i>Enterococcus</i>	<i>faecalis</i>	(Andrewes and Horder, 1906) Schleifer and Kilpper-Bälz, 1984	
	<i>Enterococcus</i>	<i>faecalis</i>	(Andrewes and Horder, 1906) Schleifer and Kilpper-Bälz, 1984	
	<i>Enterococcus</i>	<i>faecium</i>	(Orla-Jensen, 1919) Schleifer and Kilpper-Bälz, 1984	
	<i>Enterococcus</i>	<i>flavescens</i>	Pompei <i>et al.</i> , 1992	
	<i>Enterococcus</i>	<i>gallinarum</i>	(Bridge and Sneath, 1982) Collins <i>et al.</i> , 1984	
	<i>Enterococcus</i>	<i>gilvus</i>	Tyrell <i>et al.</i> , 2002	
	<i>Enterococcus</i>	<i>haemoperoxidus</i>	Svec <i>et al.</i> , 2001	
	<i>Enterococcus</i>	<i>moraviensis</i>	Svec <i>et al.</i> , 2001	
	<i>Enterococcus</i>	<i>mundtii</i>	Collins <i>et al.</i> , 1986	
	<i>Enterococcus</i>	<i>pallens</i>	Tyrell <i>et al.</i> , 2002	
	<i>Enterococcus</i>	<i>phoeniculicola</i>	Law-Brown and Meyers, 2003	
	<i>Enterococcus</i>	<i>hermanniensis</i>	Koort <i>et al.</i> , 2004	
	<i>Enterococcus</i>	<i>hirae</i>	Farrow and Collins, 1985	
	<i>Enterococcus</i>	<i>italicus</i>	Fortina <i>et al.</i> , 2004	
	<i>Enterococcus</i>	<i>malodoratus</i>	(ex Pette, 1955) Collins <i>et al.</i> , 1984	
	<i>Enterococcus</i>	<i>porcinus</i>	Teixerira <i>et al.</i> , 2001	
	<i>Enterococcus</i>	<i>pseudoavium</i>	Collins <i>et al.</i> , 1989	
	<i>Enterococcus</i>	<i>raffinosis</i>	Collins <i>et al.</i> , 1989	
	<i>Enterococcus</i>	<i>saccharolyticus</i>	(Farrow <i>et al.</i> , 1985) Rodrigues and Collins, 1991	
	<i>Enterococcus</i>	<i>seriolicida</i>	Kusuda <i>et al.</i> , 1991	
	<i>Enterococcus</i>	<i>saccharominimus</i>	Vancanneyt <i>et al.</i> , 2004	
	<i>Enterococcus</i>	<i>silesiacus</i>	Svec <i>et al.</i> , 2006	
	<i>Enterococcus</i>	<i>solitarius</i>	Collins <i>et al.</i> , 1989	
	<i>Enterococcus</i>	<i>sulfurous</i>	Martinez-Murcia and Collins, 1991	
	<i>Enterococcus</i>	<i>termitis</i>	Svec <i>et al.</i> , 2006	
	<i>Enterococcus</i>	<i>villorum</i>	Vancanneyt <i>et al.</i> , 2001	
	<i>Atopobacter</i>	<i>phocae</i>	Lawson <i>et al.</i> , 2000	
	<i>Mellissococcus</i>	<i>plutonius</i>	Bailey and Collins, 1983	
	<i>Tetragenococcus</i>	<i>halophilus</i>	(Mees, 1934) Collins <i>et al.</i> , 1993	
	<i>Tetragenococcus</i>	<i>muriatricus</i>	Satomi <i>et al.</i> , 1997	
	<i>Tetragenococcus</i>	<i>koreensis</i>	Lee <i>et al.</i> , 2005	
	<i>Vagococcus</i>	<i>elongatus</i>	Lawson <i>et al.</i> , 2007	
	<i>Vagococcus</i>	<i>fessus</i>	Hoyles <i>et al.</i> , 2000	
	<i>Vagococcus</i>	<i>fluvialis</i>	Collins <i>et al.</i> , 1990	
	<i>Vagococcus</i>	<i>lutrae</i>	Lawson <i>et al.</i> , 1999	
	<i>Vagococcus</i>	<i>salmoninarum</i>	Wallbanks <i>et al.</i> , 1990	
	Streptococcaceae	<i>Streptococcus</i>	<i>adjacens</i>	Bouvet <i>et al.</i> , 1989
		<i>Streptococcus</i>	<i>agalactiae</i>	Lehmann and Neumann, 1896
<i>Streptococcus</i>		<i>alactolyticus</i>	Farrow <i>et al.</i> , 1985	
<i>Streptococcus</i>		<i>alactolyticus</i>	Farrow <i>et al.</i> , 1985	
<i>Streptococcus</i>		<i>anginosus</i>	(Andrews and Horder, 1906) Smith and Sherman, 1938	
<i>Streptococcus</i>		<i>australis</i>	Willcox <i>et al.</i> , 2001	
<i>Streptococcus</i>		<i>bovis</i>	Orla-Jensen, 1919	
<i>Streptococcus</i>		<i>canis</i>	Devriese <i>et al.</i> , 1986	
<i>Streptococcus</i>		<i>caprinus</i>	Brooker <i>et al.</i> , 1996	
<i>Streptococcus</i>		<i>castoreus</i>	Lawson <i>et al.</i> , 2005	
<i>Streptococcus</i>		<i>cecorum</i>	Devriese <i>et al.</i> , 1983	
<i>Streptococcus</i>		<i>constellatus</i>	(Prévot, 1924) Holdeman and Moore, 1974	
<i>Streptococcus</i>		<i>constellatus</i> subsp. <i>constellatus</i>	(Prévot, 1924) Holdeman and Moore, 1974	
<i>Streptococcus</i>		<i>constellatus</i> subsp. <i>pharyngis</i>	Whiley <i>et al.</i> , 1999	
<i>Streptococcus</i>		<i>criceti</i>	Coykendall, 1977	
<i>Streptococcus</i>		<i>cristatus</i>	Handley <i>et al.</i> , 1991	
<i>Streptococcus</i>		<i>defectivus</i>	Bouvet <i>et al.</i> , 1989	
<i>Streptococcus</i>		<i>devriesei</i>	Collins <i>et al.</i> , 2004	
<i>Streptococcus</i>		<i>didelphis</i>	Rurangirwa <i>et al.</i> , 2000	
<i>Streptococcus</i>		<i>difficilis</i>	Eldar <i>et al.</i> , 1995	
<i>Streptococcus</i>		<i>adjacens</i>	Bouvet <i>et al.</i> , 1989	
<i>Streptococcus</i>		<i>durans</i>	(ex Sherman and Wing, 1937) Knight <i>et al.</i> , 1984	
<i>Streptococcus</i>		<i>dysgalactiae</i>	(ex Diernhofer, 1932) Garvie <i>et al.</i> , 1983	
<i>Streptococcus</i>		<i>dysgalactiae</i> subsp. <i>dysgalactiae</i>	(ex Diernhofer, 1932) Garvie <i>et al.</i> , 1983	
<i>Streptococcus</i>		<i>dysgalactiae</i> subsp. <i>equisimilis</i>	Vandamme <i>et al.</i> , 1996	
<i>Streptococcus</i>		<i>entericus</i>	Vela <i>et al.</i> , 2002	
<i>Streptococcus</i>		<i>equi</i>	Sand and Jensen, 1888	
<i>Streptococcus</i>		<i>equi</i> subsp. <i>equi</i>	Sand and Jensen, 1888	
<i>Streptococcus</i>		<i>equi</i> subsp. <i>ruminatorum</i>	Fernández <i>et al.</i> , 2004	



Family	Genus	Species	Author
Streptococcaceae	<i>Streptococcus</i>	<i>equi</i> subsp. <i>zooepidermicus</i>	(ex Frost and Englebrecht, 1936) Farrow and Collins, 1985
	<i>Streptococcus</i>	<i>equinus</i>	Andrewes and Horder, 1906
	<i>Streptococcus</i>	<i>faecalis</i>	Andrewes and Horder, 1906
	<i>Streptococcus</i>	<i>faecium</i>	Orla-Jensen, 1919
	<i>Streptococcus</i>	<i>ferus</i>	(ex Coykendall, 1977) Coykendall, 1983
	<i>Streptococcus</i>	<i>gallinaceus</i>	Collins <i>et al.</i> , 2002
	<i>Streptococcus</i>	<i>gallinarum</i>	Bridge and Sneath, 1982
	<i>Streptococcus</i>	<i>gallolyticus</i>	Osawa <i>et al.</i> , 1996
	<i>Streptococcus</i>	<i>gallolyticus</i> subsp. <i>gallolyticus</i>	Osawa <i>et al.</i> , 1996
	<i>Streptococcus</i>	<i>garvieae</i>	Collins <i>et al.</i> , 1984
	<i>Streptococcus</i>	<i>gordonii</i>	Kilian <i>et al.</i> , 1989
	<i>Streptococcus</i>	<i>halichoeri</i>	Lawson <i>et al.</i> , 2004
	<i>Streptococcus</i>	<i>hansenii</i>	Holdeman and Moore, 1974
	<i>Streptococcus</i>	<i>hyointestinalis</i>	Devriess <i>et al.</i> , 1988
	<i>Streptococcus</i>	<i>hyovaginalis</i>	Devriess <i>et al.</i> , 1997
	<i>Streptococcus</i>	<i>infantarius</i>	Schlegel <i>et al.</i> , 2000
	<i>Streptococcus</i>	<i>infantarius</i> subsp. <i>coli</i>	Schlegel <i>et al.</i> , 2003
	<i>Streptococcus</i>	<i>infantarius</i> subsp. <i>infantarius</i>	Schlegel <i>et al.</i> , 2003
	<i>Streptococcus</i>	<i>infantis</i>	Kawamura <i>et al.</i> , 1998
	<i>Streptococcus</i>	<i>iniae</i>	Pier and Madin, 1976
	<i>Streptococcus</i>	<i>intermedius</i>	Prévot, 1925
	<i>Streptococcus</i>	<i>intestinalis</i>	Robinson <i>et al.</i> , 1988
	<i>Streptococcus</i>	<i>lutetiensis</i>	Poyart <i>et al.</i> , 2002
	<i>Streptococcus</i>	<i>macacae</i>	Beighton <i>et al.</i> , 1984
	<i>Streptococcus</i>	<i>macedonicus</i>	Tsakalidou <i>et al.</i> , 1998
	<i>Streptococcus</i>	<i>marimammalium</i>	Lawson <i>et al.</i> , 2005
	<i>Streptococcus</i>	<i>massiliensis</i>	Glazunova <i>et al.</i> , 2006
	<i>Streptococcus</i>	<i>minor</i>	Vancanneyt <i>et al.</i> , 2004
	<i>Streptococcus</i>	<i>mitis</i>	Andrewes and Horder, 1906
	<i>Streptococcus</i>	<i>morbilloorum</i>	(Prévot, 1933) Holdeman and Moore, 1974
	<i>Streptococcus</i>	<i>mutans</i>	Coykendall, 1977
	<i>Streptococcus</i>	<i>oligofermentans</i>	Tong <i>et al.</i> , 2003
	<i>Streptococcus</i>	<i>oralis</i>	Bridge and Sneath, 1982
	<i>Streptococcus</i>	<i>orisratti</i>	Zhu <i>et al.</i> , 2000
	<i>Streptococcus</i>	<i>ovis</i>	Collins <i>et al.</i> , 2001
	<i>Streptococcus</i>	<i>parasanguinis</i>	Whiley <i>et al.</i> , 1990
	<i>Streptococcus</i>	<i>parauberis</i>	Williams and Collins, 1990
	<i>Streptococcus</i>	<i>parvulus</i>	(Weinberg <i>et al.</i> , 1937) Cato, 1983
	<i>Streptococcus</i>	<i>pasteurianus</i>	Poyart <i>et al.</i> , 2002
	<i>Streptococcus</i>	<i>peroris</i>	Kawamura <i>et al.</i> , 1998
	<i>Streptococcus</i>	<i>plantarum</i>	Collins <i>et al.</i> , 1984
	<i>Streptococcus</i>	<i>pleomorphus</i>	Barnes <i>et al.</i> , 1979
	<i>Streptococcus</i>	<i>pluranimalium</i>	Devriess <i>et al.</i> , 1999
	<i>Streptococcus</i>	<i>pneumoniae</i>	(Klein, 1884) Chester, 1901
	<i>Streptococcus</i>	<i>porcinus</i>	Collins <i>et al.</i> , 1985
	<i>Streptococcus</i>	<i>pseudopneumoniae</i>	Arbique <i>et al.</i> , 2005
	<i>Streptococcus</i>	<i>pseudoporcinus</i>	Bekal <i>et al.</i> , 2007
	<i>Streptococcus</i>	<i>pyogenes</i>	Rosenbach, 1884
	<i>Streptococcus</i>	<i>raffinolactis</i>	Orla-Jensen and Hansen, 1932
	<i>Streptococcus</i>	<i>ratti</i>	Coykendall, 1977
	<i>Streptococcus</i>	<i>saccharolyticus</i>	Farrow <i>et al.</i> , 1985
	<i>Streptococcus</i>	<i>salivarius</i>	Andrewes and Horder, 1906
	<i>Streptococcus</i>	<i>salivarius</i> subsp. <i>salivarius</i>	Andrewes and Horder, 1906
	<i>Streptococcus</i>	<i>salivarius</i> subsp. <i>thermophilus</i>	(Orla-Jensen, 1919) Farrow and Collins, 1984
	<i>Streptococcus</i>	<i>sanguinis</i>	White and Niven, 1946
	<i>Streptococcus</i>	<i>shiloi</i>	Eldar <i>et al.</i> , 1995
	<i>Streptococcus</i>	<i>sinensis</i>	Woo <i>et al.</i> , 2002
	<i>Streptococcus</i>	<i>sobrinus</i>	(ex Coykendall, 1974) Coykendall, 1983
	<i>Streptococcus</i>	<i>suis</i>	(ex Elliot, 1966) Kilper-Bälz and Schleifer 1987
	<i>Streptococcus</i>	<i>thermophilus</i> *	Orla-Jensen, 1919
	<i>Streptococcus</i>	<i>thoraltensis</i>	Devriess <i>et al.</i> , 1997
<i>Streptococcus</i>	<i>uberis</i>	Diernhofer, 1932	
<i>Streptococcus</i>	<i>urinalis</i>	Collins <i>et al.</i> , 2000	
<i>Streptococcus</i>	<i>vestibularis</i>	Whiley and Hardie, 1988	
<i>Streptococcus</i>	<i>waius</i>	Flint <i>et al.</i> , 1999	
<i>Lactococcus</i>	<i>garvieae</i>	(Collins <i>et al.</i> , 1984) Schleifer <i>et al.</i> , 1986	
<i>Lactococcus</i>	<i>lactis</i> subsp. <i>cremoris</i> *	(Orla-Jensen, 1919) Schleifer <i>et al.</i> , 1986	
<i>Lactococcus</i>	<i>lactis</i> subsp. <i>hordniae</i>	(ex Latorre-Guzman <i>et al.</i> , 1977) Schleifer <i>et al.</i> , 1986	
<i>Lactococcus</i>	<i>lactis</i> subsp. <i>lactis</i> *	(Lister, 1873) Schleifer <i>et al.</i> , 1986	
<i>Lactococcus</i>	<i>piscium</i>	Williams <i>et al.</i> , 1990	
<i>Lactococcus</i>	<i>plantarum</i>	(Collins <i>et al.</i> , 1984) Schleifer <i>et al.</i> , 1986	
Incertae sedis	<i>Acetonaerobium</i>	<i>noterae</i>	Sleat <i>et al.</i> , 1985
	<i>Oscillospira</i>	<i>guilliermondii</i>	Chatton and Perad, 1913

* These species are widely used in industry as starter cultures.

Appendix D - Multiple 16S rRNA gene sequence alignment for LAB isolated from natural sorghum fermentations and type strains (Chapter 3). Sequences for isolates from sorghum slurries are indicated in bold.

	10	20	30	40	50	60	70	80	90	100
<i>Lb. casei</i> LMG6904T					aatacatgc	aagtcgaacgagtt	ctgattatt	gaaaggtgct	gcac	cttgatt
<i>Lb. fermentum</i> LMG6902T								ggc	ca	ga
<i>Lb. paracasei</i> LMG1308T								cg	ga	tc
<i>Lb. paraplantarum</i> LMG16673T								ctgg	aa	tt
<i>Lb. pentosus</i> LMG10755T								ac	ctgg	tt
<i>Lb. plantarum</i> LMG9205T								ac	ctgg	tt
<i>Lb. rhamnosus</i> LMG6400T										
4.35								ggc	ggc	ca
3.30								ggc	ca	ga
1.42								ctgg	aa	tt
5.30								ggc	ggc	ca
3.42								ggc	ggc	ca
2.35								ctgg	aa	tt
<i>B. subtilis</i> NCD01769T								g	gaca	ttg
<i>Lb. zeae</i> LMG17315T								tt	g	cg
<i>Lb. oris</i> LMG9848T								g	cac	ggc
<i>Lb. antri</i> LMG22111T								ttgat	cct	ggct
<i>Lb. panis</i> LMG21658T								g	cac	ggc
<i>Lb. frumenti</i> LMG19473T								agagt	ttgat	tat
<i>Lb. vaginalis</i> LMG12891T								g	cac	ggc
<i>Lb. pontis</i> LMG14187T								g	cac	ggc
<i>Lb. reuteri</i> LMG9213T								agagt	ttgat	nt
<i>Lb. coleohominis</i> DSM14060T								g	cac	ggc
<i>Lb. mucosae</i> LMG19534T								agagt	ttgat	cct
<i>Lb. gastricus</i> LMG22113T								g	cac	ggc
<i>Lb. suebicus</i> LMG11408T								g	cac	ggc
<i>Lb. durianis</i> LMG19193T								g	cac	ggc
<i>Lb. vaccinostercus</i> LMG9215T								g	cac	ggc
<i>Lb. rossiae</i> LMG22972T								g	cac	ggc
<i>Lb. plantarum</i> LMG19807T								g	cac	ggc
<i>Lb. arizonensis</i> LMG19807T*								g	cac	ggc
<i>Lb. ingluviei</i> LMG20380T								ctg	ctc	agg
	110	120	130	140	150	160	170	180	190	200
<i>Lb. casei</i> LMG6904T										
<i>Lb. fermentum</i> LMG6902T										
<i>Lb. paracasei</i> LMG1308T										
<i>Lb. paraplantarum</i> LMG16673T										
<i>Lb. pentosus</i> LMG10755T										
<i>Lb. plantarum</i> LMG9205T										
<i>Lb. rhamnosus</i> LMG6400T										
4.35										
3.30										
1.42										
5.30										
3.42										
2.35										
<i>B. subtilis</i> NCD01769T										
<i>Lb. zeae</i> LMG17315T										
<i>Lb. oris</i> LMG9848T										
<i>Lb. antri</i> LMG22111T										
<i>Lb. panis</i> LMG21658T										
<i>Lb. frumenti</i> LMG19473T										
<i>Lb. vaginalis</i> LMG12891T										
<i>Lb. pontis</i> LMG14187T										
<i>Lb. reuteri</i> LMG9213T										
<i>Lb. coleohominis</i> DSM14060T										
<i>Lb. mucosae</i> LMG19534T										
<i>Lb. gastricus</i> LMG22113T										
<i>Lb. suebicus</i> LMG11408T										
<i>Lb. durianis</i> LMG19193T										
<i>Lb. vaccinostercus</i> LMG9215T										
<i>Lb. rossiae</i> LMG22972T										
<i>Lb. plantarum</i> LMG19807T										
<i>Lb. arizonensis</i> LMG19807T*										
<i>Lb. ingluviei</i> LMG20380T										
	210	220	230	240	250	260	270	280	290	300
<i>Lb. casei</i> LMG6904T										
<i>Lb. fermentum</i> LMG6902T										
<i>Lb. paracasei</i> LMG1308T										
<i>Lb. paraplantarum</i> LMG16673T										
<i>Lb. pentosus</i> LMG10755T										
<i>Lb. plantarum</i> LMG9205T										
<i>Lb. rhamnosus</i> LMG6400T										
4.35										
3.30										
1.42										
5.30										
3.42										
2.35										
<i>B. subtilis</i> NCD01769T										
<i>Lb. zeae</i> LMG17315T										
<i>Lb. oris</i> LMG9848T										
<i>Lb. antri</i> LMG22111T										
<i>Lb. panis</i> LMG21658T										
<i>Lb. frumenti</i> LMG19473T										
<i>Lb. vaginalis</i> LMG12891T										
<i>Lb. pontis</i> LMG14187T										
<i>Lb. reuteri</i> LMG9213T										
<i>Lb. coleohominis</i> DSM14060T										
<i>Lb. mucosae</i> LMG19534T										
<i>Lb. gastricus</i> LMG22113T										
<i>Lb. suebicus</i> LMG11408T										
<i>Lb. durianis</i> LMG19193T										
<i>Lb. vaccinostercus</i> LMG9215T										
<i>Lb. rossiae</i> LMG22972T										
<i>Lb. plantarum</i> LMG19807T										
<i>Lb. arizonensis</i> LMG19807T*										
<i>Lb. ingluviei</i> LMG20380T										



Lb. casei LMG6904T
Lb. fermentum LMG6902T
Lb. paracasei LMG1308T
Lb. paraplantarum LMG16673T
Lb. pentosus LMG10755T
Lb. plantarum LMG9205T
Lb. rhamnosus LMG6400T
4.35
3.30
1.42
5.30
3.42
2.35
B. subtilis NCD01769T
Lb. zeae LMG17315T
Lb. oris LMG9848T
Lb. antri LMG22111T
Lb. panis LMG21658T
Lb. frumenti LMG19473T
Lb. vaginalis LMG12891T
Lb. pontis LMG14187T
Lb. reuteri LMG9213T
Lb. coleohominis DSM14060T
Lb. mucosae LMG19534T
Lb. gastricus LMG22113T
Lb. suebicus LMG11408T
Lb. durianis LMG19193T
Lb. vaccिनosterculus LMG9215T
Lb. rossiae LMG22972T
Lb. plantarum LMG19807T
Lb. arizonensis LMG19807T*
Lb. ingluviei LMG20380T

Lb. casei LMG6904T
Lb. fermentum LMG6902T
Lb. paracasei LMG1308T
Lb. paraplantarum LMG16673T
Lb. pentosus LMG10755T
Lb. plantarum LMG9205T
Lb. rhamnosus LMG6400T
4.35
3.30
1.42
5.30
3.42
2.35
B. subtilis NCD01769T
Lb. zeae LMG17315T
Lb. oris LMG9848T
Lb. antri LMG22111T
Lb. panis LMG21658T
Lb. frumenti LMG19473T
Lb. vaginalis LMG12891T
Lb. pontis LMG14187T
Lb. reuteri LMG9213T
Lb. coleohominis DSM14060T
Lb. mucosae LMG19534T
Lb. gastricus LMG22113T
Lb. suebicus LMG11408T
Lb. durianis LMG19193T
Lb. vaccिनosterculus LMG9215T
Lb. rossiae LMG22972T
Lb. plantarum LMG19807T
Lb. arizonensis LMG19807T*
Lb. ingluviei LMG20380T

Lb. casei LMG6904T
Lb. fermentum LMG6902T
Lb. paracasei LMG1308T
Lb. paraplantarum LMG16673T
Lb. pentosus LMG10755T
Lb. plantarum LMG9205T
Lb. rhamnosus LMG6400T
4.35
3.30
1.42
5.30
3.42
2.35
B. subtilis NCD01769T
Lb. zeae LMG17315T
Lb. oris LMG9848T
Lb. antri LMG22111T
Lb. panis LMG21658T
Lb. frumenti LMG19473T
Lb. vaginalis LMG12891T
Lb. pontis LMG14187T
Lb. reuteri LMG9213T
Lb. coleohominis DSM14060T
Lb. mucosae LMG19534T
Lb. gastricus LMG22113T
Lb. suebicus LMG11408T
Lb. durianis LMG19193T
Lb. vaccिनosterculus LMG9215T
Lb. rossiae LMG22972T
Lb. plantarum LMG19807T
Lb. arizonensis LMG19807T*
Lb. ingluviei LMG20380T

```

610          660          670          680          690          700
.....|.....|.....|.....|.....|.....|.....|.....|.....|.....|.....|
ttggcgtaaagcgagcgcagggcgg-ttttttaagtctgatgtgaaagcc-ctcggttaaccgagnhagtgatcggaactgggaaacttgagtgcag
.....|.....|.....|.....|.....|.....|.....|.....|.....|.....|.....|
Lb. casei LMG6904T      .a..t.....-..c.....-t.....ga.....at.....
Lb. fermentum LMG6902T .a..t.....-..c.....-t.....ga.....at.....
Lb. paracasei LMG1308T .a..t.....-..c.....-t.....ga.....at.....
Lb. paraplantarum LMG16673T .a..t.....-..c.....-t.....ga.....at.....
Lb. pentosus LMG10755T .a..t.....-..c.....-t.....ga.....at.....
Lb. plantarum LMG9205T .a..t.....-..c.....-t.....ga.....at.....
Lb. rhamnosus LMG6400T .a..t.....-..c.....-t.....ga.....at.....
4.35 .a..t.....-..c.....-t.....ga.....at.....
3.30 .a..t.....-..c.....-t.....ga.....at.....
1.42 .a..t.....-..c.....-t.....ga.....at.....
5.30 .a..t.....-..c.....-t.....ga.....at.....
3.42 .a..t.....-..c.....-t.....ga.....at.....
2.35 .a..t.....-..c.....-t.....ga.....at.....
B. subtilis NCD01769T .g.ct.....-..c.....-c.....g..g.gt...t.....g.....
Lb. zeae LMG17315T .a..t.....-..c.....-t.....ga.....at.....
Lb. oris LMG9848T .gc..g.....-t.....a.....c.....c.....cg.....
Lb. antri LMG22111T .gc..g.....-t.....a.....c.....c.....cg.....
Lb. panis LMG21658T .gc..g.....-t.....a.....c.....c.....cg.....
Lb. frumenti LMG19473T .gc..g.....-t.....a.....c.....c.....cg.....
Lb. vaginalis LMG12891T .a..-gc..g.....-t.....a.....g.....c.....cg.....
Lb. pontis LMG14187T .ac.....-t.....a.....t.....g.....
Lb. reuteri LMG9213T .gc..g.....-t.....a.....c.....c.....cg.....
Lb. coleohominis DSM14060T .gc..g.....-t.....ga.....g.....at.....a.....
Lb. mucosae LMG19534T .ga.....-t.t.....a.....t.c.g.....
Lb. gastricus LMG22113T .a..t.....-..c.....-t.....c.....a.....g.....a.....
Lb. suebicus LMG11408T .a.....-t.....a.....g.....
Lb. durianis LMG19193T .a.....-t.....a.....g.....
Lb. vaccिनosterculus LMG9215T .a.....-t.....a.....g.....
Lb. rossiae LMG22972T .g..t.....-a.....-t.....ga.....g.....at.....
Lb. plantarum LMG19807T .c.....-ncn..n.....nnn-t.....c.....a.....
Lb. arizonensis LMG19807T* .a.....-t.....c.....a.....
Lb. ingluviei LMG20380T .c.....-t.....a.....atg.....

710          720          730          740          750          760          770          780          790          800
.....|.....|.....|.....|.....|.....|.....|.....|.....|.....|.....|
aagaggacagtggaaactccatgtgtagcgggtgaaatgcgtag-atataggagaacaccagtgccgaaggcggcngtctgg-tctgtaactgacgctga
.....|.....|.....|.....|.....|.....|.....|.....|.....|.....|.....|
Lb. casei LMG6904T      .gt.....g.....-t.....at.....g.....ac.....c.....
Lb. fermentum LMG6902T .gt.....g.....-t.....at.....g.....ac.....c.....
Lb. paracasei LMG1308T .gt.....g.....-t.....at.....g.....ac.....c.....
Lb. paraplantarum LMG16673T .gt.....g.....-t.....at.....g.....ac.....c.....
Lb. pentosus LMG10755T .gt.....g.....-t.....at.....g.....ac.....c.....
Lb. plantarum LMG9205T .gt.....g.....-t.....at.....g.....ac.....c.....
Lb. rhamnosus LMG6400T .gt.....g.....-t.....at.....g.....ac.....c.....
4.35 .gt.....g.....-t.....at.....g.....ac.....c.....
3.30 .gt.....g.....-t.....at.....g.....ac.....c.....
1.42 .gt.....g.....-t.....at.....g.....ac.....c.....
5.30 .gt.....g.....-t.....at.....g.....ac.....c.....
3.42 .gt.....g.....-t.....at.....g.....ac.....c.....
2.35 .gt.....g.....-t.....at.....g.....ac.....c.....
B. subtilis NCD01769T .g.....t.....c.....n.....-g..g...g.....a.c.....
Lb. zeae LMG17315T .g.....-t.....c.....
Lb. oris LMG9848T .g.....-t.....c.....
Lb. antri LMG22111T .g.....-t.....c.....
Lb. panis LMG21658T .g.....-t.....c.....
Lb. frumenti LMG19473T .g.....-t.....c.....
Lb. vaginalis LMG12891T .g.....-t.....c.....
Lb. pontis LMG14187T .g.....-t.....c.....
Lb. reuteri LMG9213T .g.....-t.....c.....
Lb. coleohominis DSM14060T .g.....-t.....a.ct.c.....
Lb. mucosae LMG19534T .g.....-t.....a.....c.....
Lb. gastricus LMG22113T .g.....-t.....c.....c.....
Lb. suebicus LMG11408T .g.....-t.....c.....
Lb. durianis LMG19193T .g.....-t.....c.....
Lb. vaccिनosterculus LMG9215T .gt.....g.....-t.....ac.a.....
Lb. rossiae LMG22972T .gt.....g.....-t.....ac.a.....
Lb. plantarum LMG19807T .gt.....g.....-t.....n..n..n.....
Lb. arizonensis LMG19807T* .gt.....g.....-t.....n.....n.....
Lb. ingluviei LMG20380T .gt.....g.....-t.....c.....

810          820          830          840          850          860          870          880          890          900
.....|.....|.....|.....|.....|.....|.....|.....|.....|.....|.....|
ggctcgaagcatgggtagcgaacaggattagataccctggtagtccatgccgtaaacgatgaatgctagggtgttg-gagggtttccgcccttcagtgcc
.....|.....|.....|.....|.....|.....|.....|.....|.....|.....|.....|
Lb. casei LMG6904T      .a.....g.....-t.....
Lb. fermentum LMG6902T .a.....g.....-t.....
Lb. paracasei LMG1308T .a.....g.....-t.....
Lb. paraplantarum LMG16673T .a.....g.....-t.....
Lb. pentosus LMG10755T .a.....g.....-t.....
Lb. plantarum LMG9205T .a.....g.....-t.....
Lb. rhamnosus LMG6400T .a.....g.....-t.....
4.35 .a.....g.....-t.....
3.30 .a.....g.....-t.....
1.42 .a.....g.....-t.....
5.30 .a.....g.....-t.....
3.42 .a.....g.....-t.....
2.35 .a.....g.....-t.....
B. subtilis NCD01769T .ag.....g.....g.....c.....g.....a.....a.g.....c.t.....t.....
Lb. zeae LMG17315T .a.....g.....-t.....
Lb. oris LMG9848T .a.....g.....-t.....
Lb. antri LMG22111T .a.....g.....-t.....
Lb. panis LMG21658T .a.....g.....-t.....
Lb. frumenti LMG19473T .a.....g.....-t.....
Lb. vaginalis LMG12891T .a.....g.....-t.....
Lb. pontis LMG14187T .a.....g.....-t.....
Lb. reuteri LMG9213T .a.....g.....-t.....
Lb. coleohominis DSM14060T .a.....g.....-t.....
Lb. mucosae LMG19534T .a.....g.....-t.....
Lb. gastricus LMG22113T .a.....g.....-t.....
Lb. suebicus LMG11408T .a.....g.....-t.....
Lb. durianis LMG19193T .a.....g.....-t.....
Lb. vaccिनosterculus LMG9215T .a.....g.....-t.....
Lb. rossiae LMG22972T .a.....g.....-t.....
Lb. plantarum LMG19807T .a.....g.....-t.....
Lb. arizonensis LMG19807T* .a.....g.....-t.....
Lb. ingluviei LMG20380T .a.....g.....-t.....

```



910 920 930 940 950 960 970 980 990 1000

Lb. casei LMG6904T
Lb. fermentum LMG6902T
Lb. paracasei LMG1308T
Lb. paraplantarum LMG16673T
Lb. pentosus LMG10755T
Lb. plantarum LMG9205T
Lb. rhamnosus LMG6400T
4.35
3.30
1.42
5.30
3.42
2.35
B. subtilis NCD01769T
Lb. zeae LMG17315T
Lb. oris LMG9848T
Lb. antri LMG22111T
Lb. panis LMG21658T
Lb. frumenti LMG19473T
Lb. vaginalis LMG12891T
Lb. pontis LMG14187T
Lb. reuteri LMG9213T
Lb. coleohominis DSM14060T
Lb. mucosae LMG19534T
Lb. gastricus LMG22113T
Lb. suebicus LMG11408T
Lb. durianis LMG19193T
Lb. vaccिनosterculus LMG9215T
Lb. rossiae LMG22972T
Lb. plantarum LMG19807T
Lb. arizonensis LMG19807T*
Lb. ingluviei LMG20380T

1010 1020 1030 1040 1050 1060 1070 1080 1090 1100

Lb. casei LMG6904T
Lb. fermentum LMG6902T
Lb. paracasei LMG1308T
Lb. paraplantarum LMG16673T
Lb. pentosus LMG10755T
Lb. plantarum LMG9205T
Lb. rhamnosus LMG6400T
4.35
3.30
1.42
5.30
3.42
2.35
B. subtilis NCD01769T
Lb. zeae LMG17315T
Lb. oris LMG9848T
Lb. antri LMG22111T
Lb. panis LMG21658T
Lb. frumenti LMG19473T
Lb. vaginalis LMG12891T
Lb. pontis LMG14187T
Lb. reuteri LMG9213T
Lb. coleohominis DSM14060T
Lb. mucosae LMG19534T
Lb. gastricus LMG22113T
Lb. suebicus LMG11408T
Lb. durianis LMG19193T
Lb. vaccिनosterculus LMG9215T
Lb. rossiae LMG22972T
Lb. plantarum LMG19807T
Lb. arizonensis LMG19807T*
Lb. ingluviei LMG20380T

1110 1120 1130 1140 1150 1160 1170 1180 1190 1200

Lb. casei LMG6904T
Lb. fermentum LMG6902T
Lb. paracasei LMG1308T
Lb. paraplantarum LMG16673T
Lb. pentosus LMG10755T
Lb. plantarum LMG9205T
Lb. rhamnosus LMG6400T
4.35
3.30
1.42
5.30
3.42
2.35
B. subtilis NCD01769T
Lb. zeae LMG17315T
Lb. oris LMG9848T
Lb. antri LMG22111T
Lb. panis LMG21658T
Lb. frumenti LMG19473T
Lb. vaginalis LMG12891T
Lb. pontis LMG14187T
Lb. reuteri LMG9213T
Lb. coleohominis DSM14060T
Lb. mucosae LMG19534T
Lb. gastricus LMG22113T
Lb. suebicus LMG11408T
Lb. durianis LMG19193T
Lb. vaccिनosterculus LMG9215T
Lb. rossiae LMG22972T
Lb. plantarum LMG9205T
Lb. arizonensis LMG19807T*
Lb. ingluviei LMG20380T

Appendix E - Multiple 16S rRNA gene sequence alignment for *Lactobacillus* and *Weissella* species isolated from natural sorghum fermentations and type strains (Chapter 4). Sequences for isolates from sorghum slurries are indicated in bold.

	10	20	30	40	50	60	70	80	90	100	
P11	aaatacatgc	---aagt	cgaacgaact	-----ctgg	tattgat	tggtgct	tgcat	catgat	---ttacat	tt---ga	gtgagtc
P1N	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
P31	-----	gt	-----ctgataatttatacgg
L235N	-----	gt	-----ctgataatttatacgg
P3	-----	gt	-----ctgataatttatacgg
M5444Nc	gca.gtcctt	gtggtt	.aac.ga	ttgaa.atcagat	.gacga	gg.ca	.gc.aag
M544Mc	gca.gtcctt	gtggtt	.aac.ga	ttgaa.atcagat	.gacga	gg.ca	.gc.aag
105c	gca.gtcctt	gtggtt	.aac.ga	ttgaa.atcagat	.gacga	gg.ca	.gc.aag
27291	-----	t	-----cgt	g-gcc	.aat.ga	tgac	tgat.gatt	.gg-	-cgctacg
27292	-----	t	-----cgt	g-gcc	.aat.ga	tgac	tgat.gatt	.gg-	-cgctacg
27293	-----	t	-----cgt	g-gcc	.aat.ga	tgac	tgat.gatt	.gg-	-cgctacg
27294	-----	t	-----cgt	g-gcc	.aat.ga	tgac	tgat.gatt	.gg-	-cgctacg
P2	-----	gc	g-gcc	.aac	gaaatgaccaga	atggacg	.gg-	cccc
<i>Lb. antri</i> LMG22111T	-----	gc	g-gcc	.aac	gaaatgaccaga	atggacg	.gg-	cccc
<i>Lb. casei</i> LMG6904T	-----	gt	-----ctgataatttatacgg
<i>Lb. coleohominis</i> DSM14060T	-----	gt	-----ctgataatttatacgg
<i>Lb. fermentum</i> LMG6902T	-----	t	-----cgt	g-gcc	.aat.ga	tgac	tgat.gatt	.ggtc	-----ctacg
<i>Lb. gastricus</i> LMG22113T	-----	cgt	g-gcc	.aac.ga	tgac.agat	gacg	.ggct	acg
<i>Lb. mucosae</i> LMG19534T	-----	cgt	g-gcc	.aac.ga	tgac.agat	gacg	.ggct	acg
<i>Lb. oris</i> LMG9848T	-----	gc	g-gcc	.aac	gaaatgacctgatt	gacg	.gg-	cccc
<i>Lb. panis</i> LMG21658T	-----	gc	g-gcc	.aac	gaaatgacctgatt	gacg	.gg-	cccc
<i>Lb. pentosus</i> LMG10755T	-----	gt	-----ctgataatttatacgg
<i>Lb. rhamnosus</i> LMG6400T	-----	tc	g-gcc	.aac	ga.tgac	tgat.gacga	gg	-----caccg
<i>Lb. reuteri</i> LMG9213T	-----	gc	g-gcc	.aac	ga.atgacctgaat	gacg	.gg	-----accc
<i>Lb. vaginalis</i> LMG12891T	-----	gt	-----ctgataatttatacgg
<i>Lb. zeae</i> LMG17315T	-----	gt	-----ctgataatttatacgg
<i>Lb. arizonensis</i> LMG19807Tc	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. frumenti</i> LMG19473T	-----	gc	g-gcc	.aac	ga.atgacctgatt	gacga	gg	-----accc
<i>Lb. ingluviei</i> LMG20380T	-----	cgt	g-gcc	.aac	ga.tga	tcaggat	gacg	.gg	-----accc
<i>Lb. paracasei</i> LMG1308T	-----	gt	-----ctc	t.gacc	ga	-----ca	-----c	-----a	-----g
<i>Lb. paraplantarum</i> DSM10667T	-----	n	-----n	-----n	-----n	-----n	-----n	-----n	-----n	-----n	-----n
<i>Lb. plantarum</i> LMG19807T	-----	n	-----n	-----n	-----n	-----n	-----n	-----n	-----n	-----n	-----n
<i>Lb. plantarum</i> LMG9205T	-----	n	-----n	-----n	-----n	-----n	-----n	-----n	-----n	-----n	-----n
<i>Lb. rossiae</i> LMG22972T	-----	ac	t-cggg	.tt.ga	tgacc	gat.gacgagg	-----ag
<i>Lb. suebicus</i> LMG11408T	-----	tc	t-c	cggt	.tc.ga	tgacc	taat.gacgagg	-----a
<i>Lb. vaccिनosterсus</i> LMG9215T	-----	tc	t-c	cggt	.tt.ga	tgaac	tgat.gataaacg-a
<i>Lb. pontis</i> LMG14187T	-----	yc	.aac	ga.atgacctgatt	gacga	gg	-----accc	-----gg
<i>W. cibaria</i> LMG17699T	-----	ctt	gtggtt	.aac	ga	ttgaa.atcagat	.gacga	gg.ca	.gc.aag
<i>W. confusa</i> JCM1093T	-----	ctt	gtggtt	.aac	ga	ttgaa.atcagat	.gacga	gg.ca	.gc.aag
<i>W. halotolerans</i> NRIC1627T	-----	cct	gtcgtt	.aac	ga	ttaga.atca	.ac	gacg	aga.cta	ac.ag
<i>W. hellenica</i> NCFB2973T	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>W. kandleri</i> NRIC1628T	-----	cgtggtt	gaaa	gagatga	aat	.a	-----tcaaat	gcca	.ca
<i>W. kimchii</i> DSM14295T	-----	ctt	gtggtt	.aac	gaaatgaatcagat	.gacga	gg.ca	.gc.aagg
<i>W. koreensis</i> S-5623T	-----	ct	gtggtt	gaaa	ga.atgaa	aattcagat	gatt	.ca	.gc
<i>W. minor</i> NRIC1625T	-----	ctt	gtggtt	.aac	ga	atgaa.atcggatt	gaaga	ga	.ca	.gc
<i>W. paramesenteroides</i> NRIC154T	-----	ctt	gt-c	cttaan	.aac	ga	ttgan.actgatt	gat	-----tt	.c
<i>W. soli</i> LMG20113T	-----	ctt	gtggtt	.aac	ga	atagcca	at	gatt	.a	.caa
<i>W. thailandensis</i> FS61-1T	-----	ctt	gt-ntt	aat	.aac	ga	atgaa.atntgatt	gat	-----tt	.c
<i>W. viridescens</i> NRIC1536T	-----	ctt	gtggtc	.aac	ga	ttgaa.atcagat	.gacga	gg.ca	.gc.aag
<i>Lb. durianis</i> LMG19193T	-----	tc	t-c	cggt	.tt.ga	tgaac	tgat.gataaacg-a
<i>B. subtilis</i> NCD01769T	-----	gg	.ag	-----	-----	-----	-----	-----	-----	-----

	110	120	130	140	150	160	170	180	190	200	
P11	acacgtggg	aaacctg	ccca-gaa	gcggggg	ataacac	cttggaa	acagat	gctaata	accgcat	aaaca	actggac
P1N	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
P31	-----	t	-----t	-----t	-----t	-----t	-----atc	aa.a	-----t	-----tt	-----gc
L235N	-----	t	-----t	-----t	-----t	-----t	-----atc	aa.a	-----t	-----tt	-----gc
P3	-----	t	-----t	-----t	-----t	-----t	-----atc	aa.a	-----t	-----tt	-----gc
M5444Na	tc-ttattttagcatgctacaaaa
M544Ma	tc-ttattttagcatgctacaaaa
105a	tc-ttattttagcatgctacaaaa
27291a	tc-ttactttttaa	aacca
27292a	tc-ttactttttaa	aacca
27293a	tc-ttactttttaa	aacca
27294a	tc-ttactttttaa	aacca
P2a	tc-ttactttttaa	aacca
<i>Lb. antri</i> LMG22111Tc	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. casei</i> LMG6904Tc	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. coleohominis</i> DSM14060Ta	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. fermentum</i> LMG6902Ta	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. gastricus</i> LMG22113Ta	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. mucosae</i> LMG19534Ta	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. oris</i> LMG9848Tcc	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. panis</i> LMG21658Tc	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. pentosus</i> LMG10755Tt	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. rhamnosus</i> LMG6400Ta	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. reuteri</i> LMG9213Ta	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. vaginalis</i> LMG12891Tc	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. zeae</i> LMG17315Tt	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. arizonensis</i> LMG19807Tc	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. frumenti</i> LMG19473Tc	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. ingluviei</i> LMG20380Ta	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. paracasei</i> LMG1308Tt	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. paraplantarum</i> DSM10667Tc	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. plantarum</i> LMG19807Tc	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. plantarum</i> LMG9205Tg	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. rossiae</i> LMG22972Tt	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. suebicus</i> LMG11408Tt	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. vaccिनosterсus</i> LMG9215Tc	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>Lb. pontis</i> LMG14187Tt	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>W. cibaria</i> LMG17699Ta	tc-ttattttagcatgctacaaaa
<i>W. confusa</i> JCM1093Ta	tc-ttattttagcatgctacaaaa
<i>W. halotolerans</i> NRIC1627Ta	tc-ttattttagcatgctacaaaa
<i>W. hellenica</i> NCFB2973Tt	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>W. kandleri</i> NRIC1628Ta	tc-ttattttagcatgctacaaaa
<i>W. kimchii</i> DSM14295Ta	tc-ttattttagcatgctacaaaa
<i>W. koreensis</i> S-5623Ta	tc-ttattttagcatgctacaaaa
<i>W. minor</i> NRIC1625Ta	tc-ttattttagcatgctacaaaa
<i>W. paramesenteroides</i> NRIC154Tt	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>W. soli</i> LMG20113Ta	tc-ttattttagcatgctacaaaa
<i>W. thailandensis</i> FS61-1Tt	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<i>W. viridescens</i> NRIC1536Ta	tc-ttattttagcatgctacaaaa



P11
P1N
P31
I235N
P3
M5444N
M544M
105
27291
27292
27293
27294
P2
Lb. antri LMG22111T
Lb. casei LMG6904T
Lb. coleohominis DSM14060T
Lb. fermentum LMG6902T
Lb. gastricus LMG22113T
Lb. mucosae LMG19534T
Lb. oris LMG9848T
Lb. panis LMG21658T
Lb. pentosus LMG10755T
Lb. rhamnosus LMG6400T
Lb. reuteri LMG9213T
Lb. vaginalis LMG12891T
Lb. zeae LMG17315T
Lb. arizonensis LMG19807T
Lb. frumenti LMG19473T
Lb. ingluviei LMG20380T
Lb. paracasei LMG1308T
Lb. paraplantarum DSM10667T
Lb. plantarum LMG19807T
Lb. plantarum LMG9205T
Lb. rossiae LMG22972T
Lb. suebicus LMG11408T
Lb. vaccinostercus LMG9215T
Lb. pontis LMG14187T
W. cibaria LMG17699T
W. confusa JCM1093T
W. halotolerans NRIC1627T
W. hellenica NCFB2973T
W. kandleri NRIC1628T
W. kimchii DSM14295T
W. koreensis S-5623T
W. minor NRIC1625T
W. paramesenteroides NRIC154T
W. soli LMG20113T
W. thailandensis FS61-1T
W. viridescens NRIC1536T
Lb. durianis LMG19193T
B. subtilis NCD01769T

P11
P1N
P31
I235N
P3
M5444N
M544M
105
27291
27292
27293
27294
P2
Lb. antri LMG22111T
Lb. casei LMG6904T
Lb. coleohominis DSM14060T
Lb. fermentum LMG6902T
Lb. gastricus LMG22113T
Lb. mucosae LMG19534T
Lb. oris LMG9848T
Lb. panis LMG21658T
Lb. pentosus LMG10755T
Lb. rhamnosus LMG6400T
Lb. reuteri LMG9213T
Lb. vaginalis LMG12891T
Lb. zeae LMG17315T
Lb. arizonensis LMG19807T
Lb. frumenti LMG19473T
Lb. ingluviei LMG20380T
Lb. paracasei LMG1308T
Lb. paraplantarum DSM10667T
Lb. plantarum LMG19807T
Lb. plantarum LMG9205T
Lb. rossiae LMG22972T
Lb. suebicus LMG11408T
Lb. vaccinostercus LMG9215T
Lb. pontis LMG14187T
W. cibaria LMG17699T
W. confusa JCM1093T
W. halotolerans NRIC1627T
W. hellenica NCFB2973T
W. kandleri NRIC1628T
W. kimchii DSM14295T
W. koreensis S-5623T
W. minor NRIC1625T
W. paramesenteroides NRIC154T
W. soli LMG20113T
W. thailandensis FS61-1T
W. viridescens NRIC1536T
Lb. durianis LMG19193T
B. subtilis NCD01769T



P11
P1N
P31
L235N
P3
M5444N
M544M
105
27291
27292
27293
27294
P2
Lb. antri LMG22111T
Lb. casei LMG6904T
Lb. coleohominis DSM14060T
Lb. fermentum LMG6902T
Lb. gastricus LMG22113T
Lb. mucosae LMG19534T
Lb. oris LMG9848T
Lb. panis LMG21658T
Lb. pentosus LMG10755T
Lb. rhamnosus LMG6400T
Lb. reuteri LMG9213T
Lb. vaginalis LMG12891T
Lb. zeae LMG17315T
Lb. arizonensis LMG19807T
Lb. frumenti LMG19473T
Lb. ingluviei LMG20380T
Lb. paracasei LMG1308T
Lb. paraplantarum DSM10667T
Lb. plantarum LMG19807T
Lb. plantarum LMG9205T
Lb. rossiae LMG22972T
Lb. suebicus LMG11408T
Lb. vaccinosfercus LMG9215T
Lb. pontis LMG14187T
W. cibaria LMG17699T
W. confusa JCM1093T
W. halotolerans NRIC1627T
W. hellenica NCFB2973T
W. kandleri NRIC1628T
W. kimchii DSM14295T
W. koreensis S-5623T
W. minor NRIC1625T
W. paramesenteroides NRIC154T
W. soli LMG20113T
W. thailandensis FS61-1T
W. viridescens NRIC1536T
Lb. durianis LMG19193T
B. subtilis NCD01769T

P11
P1N
P31
L235N
P3
M5444N
M544M
105
27291
27292
27293
27294
P2
Lb. antri LMG22111T
Lb. casei LMG6904T
Lb. coleohominis DSM14060T
Lb. fermentum LMG6902T
Lb. gastricus LMG22113T
Lb. mucosae LMG19534T
Lb. oris LMG9848T
Lb. panis LMG21658T
Lb. pentosus LMG10755T
Lb. rhamnosus LMG6400T
Lb. reuteri LMG9213T
Lb. vaginalis LMG12891T
Lb. zeae LMG17315T
Lb. arizonensis LMG19807T
Lb. frumenti LMG19473T
Lb. ingluviei LMG20380T
Lb. paracasei LMG1308T
Lb. paraplantarum DSM10667T
Lb. plantarum LMG19807T
Lb. plantarum LMG9205T
Lb. rossiae LMG22972T
Lb. suebicus LMG11408T
Lb. vaccinosfercus LMG9215T
Lb. pontis LMG14187T
W. cibaria LMG17699T
W. confusa JCM1093T
W. halotolerans NRIC1627T
W. hellenica NCFB2973T
W. kandleri NRIC1628T
W. kimchii DSM14295T
W. koreensis S-5623T
W. minor NRIC1625T
W. paramesenteroides NRIC154T
W. soli LMG20113T
W. thailandensis FS61-1T
W. viridescens NRIC1536T
Lb. durianis LMG19193T
B. subtilis NCD01769T



P11
P1N
P31
I235N
P3
M5444N
M544M
105
27291
27292
27293
27294
P2
Lb. antri LMG22111T
Lb. casei LMG6904T
Lb. coleohominis DSM14060T
Lb. fermentum LMG6902T
Lb. gastricus LMG22113T
Lb. mucosae LMG19534T
Lb. oris LMG9848T
Lb. panis LMG21658T
Lb. pentosus LMG10755T
Lb. rhamnosus LMG6400T
Lb. reuteri LMG9213T
Lb. vaginalis LMG12891T
Lb. zeae LMG17315T
Lb. arizonensis LMG19807T
Lb. frumenti LMG19473T
Lb. ingluviei LMG20380T
Lb. paracasei LMG1308T
Lb. paraplantarum DSM10667T
Lb. plantarum LMG19807T
Lb. plantarum LMG9205T
Lb. rossiae LMG22972T
Lb. suebicus LMG11408T
Lb. vaccinostercus LMG9215T
Lb. pontis LMG14187T
W. cibaria LMG17699T
W. confusa JCM1093T
W. halotolerans NRIC1627T
W. hellenica NCFB2973T
W. kandleri NRIC1628T
W. kimchii DSM14295T
W. koreensis S-5623T
W. minor NRIC1625T
W. paramesenteroides NRIC154T
W. soli LMG20113T
W. thailandensis FS61-1T
W. viridescens NRIC1536T
Lb. durianis LMG19193T
B. subtilis NCD01769T

P11
P1N
P31
I235N
P3
M5444N
M544M
105
27291
27292
27293
27294
P2
Lb. antri LMG22111T
Lb. casei LMG6904T
Lb. coleohominis DSM14060T
Lb. fermentum LMG6902T
Lb. gastricus LMG22113T
Lb. mucosae LMG19534T
Lb. oris LMG9848T
Lb. panis LMG21658T
Lb. pentosus LMG10755T
Lb. rhamnosus LMG6400T
Lb. reuteri LMG9213T
Lb. vaginalis LMG12891T
Lb. zeae LMG17315T
Lb. arizonensis LMG19807T
Lb. frumenti LMG19473T
Lb. ingluviei LMG20380T
Lb. paracasei LMG1308T
Lb. paraplantarum DSM10667T
Lb. plantarum LMG19807T
Lb. plantarum LMG9205T
Lb. rossiae LMG22972T
Lb. suebicus LMG11408T
Lb. vaccinostercus LMG9215T
Lb. pontis LMG14187T
W. cibaria LMG17699T
W. confusa JCM1093T
W. halotolerans NRIC1627T
W. hellenica NCFB2973T
W. kandleri NRIC1628T
W. kimchii DSM14295T
W. koreensis S-5623T
W. minor NRIC1625T
W. paramesenteroides NRIC154T
W. soli LMG20113T
W. thailandensis FS61-1T
W. viridescens NRIC1536T
Lb. durianis LMG19193T
B. subtilis NCD01769T

Appendix F - Multiple *pheS* gene sequence alignment for *Lactobacillus* species isolated from spontaneous sorghum fermentations and type strains (Chapter 4). Sequences for isolates from fermented sorghum slurries are indicated in bold.

	10	20	30	40	50	60	70	80	90	100
<i>Lb. casei</i> LMG6904T	aaactttttatcaacaatgaattgctaagtcggtcagcagacgagcccgatgcaggcagcaaccatggaaaagcatgatttcagcaaggaccgctgaa									
<i>Lb. paracasei</i> LMG1308T	.g.....c.t.c.....g.a.c.....c.g.....a.t.a.....g.g.a.....c.c.t.c.....									
<i>Lb. paraplantarum</i> LMG16673T	.t..g.c.c.c.t.c...a..cg.at..t.a.ta...a...tctg.gac.ac.g.gt.at...t...c.tcg.g.g...t....									
<i>Lb. pentosus</i> LMG10755T	.t.c.c.c.c.t.c...g..cg...ct...a.g.a..ttctg.gac.ac...tt.ac.t...t...tca.g.c.at.a..									
<i>Lb. plantarum</i> LMG6907T	.c.g.c.c.c.t.c...a..cg...c.a.ca.g.....tctg.tgat...c.g.gt.ac.t...t.c.....ttct.....									
<i>Lb. arizonensis</i> LMG19807T	.c.g.c.c.c.t.c...a..cg...c.a.ca.g.....tctg.tgat...c.g.gt.ac.t...t.c.....ttct.....									
<i>Lb. plantarum</i> LMG9205T	.c.g.c.c.c.t.c...a..cg.at.gc...a.g.....tctg.tgat...c.g.gt.ac.t...t.c.....ttct.....									
<i>Lb. rhamnosus</i> LMG6400Tt.c.....g.....t.c.a.c.....g.g.....g.a.....t.ct.g.c.....									
<i>Lb. zeae</i> LMG17315Tt.g.....t.g.....g.....g.g.....t.g.....t.g.....t....									
<i>E. faecalis</i> LMG7937T	.t.g.....ttcag...ga.t.t.ta.g.t.ttca.ag.t.a.t...a.....a.c.....ttca...tg.at.acg									
P3Mt.c.....g.....g.....c.a.c.....g.g.g.....g.a.....c.t.ct.g.c.....									
P31t.c.....g.....g.....c.a.c.....g.g.g.....g.a.....c.t.ct.g.c.....									
P3Mt.c.....g.....g.....c.a.c.....g.g.g.....g.a.....c.t.ct.g.c.....									
P1N	.c.g.c.c.c.t.c...a..cg...c.a.ca.g.....tctg.tgat...c.g.gt.ac.t...t.c.....ttct.....									
P11	.c.g.c.c.c.t.c...a..cg...c.a.ca.g.....tctg.tgat...c.g.gt.ac.t...t.c.....ttct.....									
	110	120	130	140	150	160	170	180	190	200
<i>Lb. casei</i> LMG6904T	gatgatcagccgggtgtggtgtatcgtcggatgatgaagatgccacgcacagccatcagtttcatacaaatggaggcttagctcattgataagcacatt									
<i>Lb. paracasei</i> LMG1308T	a.....t.....t.g.....t.a.t.....t.t.t.....c.g.....a.ac.c.....c.....t.a									
<i>Lb. paraplantarum</i> LMG16673T	.g.ct.gtca.t.ccgc.t...g.....acc.t...g.c.ttca...a.c.....t.a.a.g...g.g.....t...									
<i>Lb. pentosus</i> LMG10755T	ag.c.tca.a.ccgc.t...g.t.cacg.t...g.c.ttc...a.c.....g.t.a.gc...ag.c.....a.t.c									
<i>Lb. plantarum</i> LMG6907T	.g.ct.gtca.t.ccgc.t...g.t.acg.t...a.c.ttc...a.....t.a.g.....g.g.c.....t...									
<i>Lb. arizonensis</i> LMG19807T	.g.ct.gtca.t.ccgc.t...g.t.acg.t...a.c.ttc...a.....t.a.g.....g.g.c.....t...									
<i>Lb. plantarum</i> LMG9205T	.g.ct.gtca.a.ccgc.c.t...g.t.acg.t...c.ttca...a.c.....t.a.a.g...g.a.c.....t...									
<i>Lb. rhamnosus</i> LMG6400T	a.....t.....c.....g.t.....a.t.....c.....c.....g.....a.tc.g.g.c.c.....t...									
<i>Lb. zeae</i> LMG17315Tt.....t.....g.....t.....a.t.....c.....c.....g.....a.tc.g.g.c.c.....t...									
<i>E. faecalis</i> LMG7937T	a.....ttct.t.gaaa.a.tc...t...aca.t...g.a.t.t.t.c.a.c.....g.t.a.c.t.tg.c...aa.tg.c									
P3M	a.....t.a.c.....t.....g.t.....t.....a.t.....c.....c.....g.....a.c.g.g.c.c.....t...									
P31	a.....t.a.c.....t.....g.t.....t.....a.t.....c.....c.....g.....a.c.g.g.c.c.....t...									
P3M	a.....t.a.c.....t.....g.t.....t.....a.t.....c.....c.....g.....a.c.g.g.c.c.....t...									
P1N	.g.ct.gtca.t.ccgc.t...g.t.acg.t...a.c.ttc...a.....t.a.g.....g.g.c.....t...									
P11	.g.ct.gtca.t.ccgc.t...g.t.acg.t...a.c.ttc...a.....t.a.g.....g.g.c.....t...									
	210	220	230	240	250	260	270	280	290	300
<i>Lb. casei</i> LMG6904T	accatggctgatctcaagggaaacgctgctgoccatgtgtcagcaagcttttggcgctgatcgaacaattcggttgctcggagttatttccattcacgg									
<i>Lb. paracasei</i> LMG1308Ta.....c.t.t.g.....c.a.....taaa...g.....c.....g.a.....t.....									
<i>Lb. paraplantarum</i> LMG16673Ta.....t.g.....c.c.t.a.cct.g.tgcc.aac.t.g.....aa.c.attcgatg.c.c.a.g.a.c.t.c.c.....									
<i>Lb. pentosus</i> LMG10755Tg.....ct.g.a.t.ct.a.ttag.agcca.aacgt.g.....t.acc.gtttgatg.a.c...g.a.c.t.c.c.....									
<i>Lb. plantarum</i> LMG6907Tg.....t.g.....c.ct.aa.ctgg.tgcc.a.actt.g.....a.c.attcgatg...c.a.g.a.c.t.c.....									
<i>Lb. arizonensis</i> LMG19807Tg.....t.g.....c.ct.aa.ctgg.tgcc.a.actt.g.....a.c.attcgatg...c.a.g.a.c.t.c.....									
<i>Lb. plantarum</i> LMG9205Tg.....c.ct.g.....t.ta.a.cctag.tgcc.a.ac.t.a.c.....acc.gtttgatg.c.c.c...g.a.c.t.c.....									
<i>Lb. rhamnosus</i> LMG6400Ta.....g.....t.a.c.....t.g.....g.....g.g.....g.g.....c.c.....c.....g.t.a									
<i>Lb. zeae</i> LMG17315Ta.....t.ct.a.g.....t.g.....g.....c.....c.....g.....c.....c.....c.....t.a									
<i>E. faecalis</i> LMG7937Ta.g...t.a.a.g...t.agaa.t...atga.aa.aa.g...t.aa...t.a...t.a...t.c...c.t.t.a									
P3Ma.....t.a.g.....t.a.c.....t.g.....c.g.g.....c.c.....c.....c.....g.t.a									
P31a.....t.a.g.....t.a.c.....t.g.....c.g.g.....c.c.....c.....c.....g.t.a									
P3Ma.....t.a.g.....t.a.c.....t.g.....c.g.g.....c.c.....c.....c.....g.t.a									
P1Ng.....t.g.....c.ct.aa.ctgg.tgcc.a.actt.g.....a.c.attcgatg...c.a.g.a.c.t.c.....									
P11g.....t.g.....c.ct.aa.ctgg.tgcc.a.actt.g.....a.c.attcgatg...c.a.g.a.c.t.c.....									
	310	320	330	340	350	360	370	380	390	400
<i>Lb. casei</i> LMG6904T	aaccgtctgttgaagtctgatgttctcgtctcgttgcggcggaaggctgcccggtttgcaagatataccgggttgattgaagtctcgggtgccggcat									
<i>Lb. paracasei</i> LMG1308T	.g.a.c.....t.....t.....t.t.....t.a.t.....t.....									
<i>Lb. paraplantarum</i> LMG16673Ta.c.....ct....ca.t....taa...aat.....tg.aa.c.....c.g.t.t...c.....gt.g.....									
<i>Lb. pentosus</i> LMG10755Tc.....ct....ga.....aac.taa.....									
<i>Lb. plantarum</i> LMG6907Ta.c.a.....ct....aa.t....taa...aat.....tg.aa.c.t...c.a.g.t...c.....a.g.....									
<i>Lb. arizonensis</i> LMG19807Ta.c.a.....ct....aa.t....taa...aat.....tg.aa.c.t...c.a.g.t...c.....a.g.....									
<i>Lb. plantarum</i> LMG9205Ta.c.a.....ct....aa.t....taa...aat.....tg.aa.c.t...c.a.g.t...c.....gt.g.....									
<i>Lb. rhamnosus</i> LMG6400Tt.....a.....c.....c.....c.....t.....t.....t.....t.....t.....t.....									
<i>Lb. zeae</i> LMG17315Tt.....g.....t.....g.....g.....g.....g.....g.....g.....g.....g.....									
<i>E. faecalis</i> LMG7937Tt.a.c.....a.....ag...t.taaa.t...gca.t.taac.c...ac...a.t...a.tt.a...g.g...									
P3Ma.c.g.....g.c.....c.t.t.....t.....t.....t.....c.....t.....t.....									
P31a.c.g.....g.c.....c.t.t.....t.....t.....t.....c.....t.....t.....									
P3Ma.c.g.....g.c.....c.t.t.....t.....t.....t.....c.....t.....t.....									
P1Na.c.a.....ct....aa.t....taa...aat.....tg.aa.c.t...c.a.g.t...c.....a.g.....									
P11a.c.a.....ct....aa.t....taa...aat.....tg.aa.c.t...c.a.g.t...c.....a.g.....									
	410	420	430	440						
<i>Lb. casei</i> LMG6904T	ggtgcatccgaatgttctgcgggcagccaagattgatgccgatgtct									
<i>Lb. paracasei</i> LMG1308Tc.....gt.a.....g.c.....t...c.t...									
<i>Lb. paraplantarum</i> LMG16673Tt.c.c.c.c.at.agaaatgt.tggc.....c.a.a.aa.									
<i>Lb. pentosus</i> LMG10755Tt.c.cc.c.gt.agaaatgt.tggc.....c.a.a.aa.									
<i>Lb. plantarum</i> LMG6907Tt.c.cc.c.gt.agaaatgt.tggc.....c.a.a.aa.									
<i>Lb. arizonensis</i> LMG19807Tt.c.cc.c.gt.agaaatgt.tggc.....c.a.a.aa.									
<i>Lb. plantarum</i> LMG9205Tt.c.cc.c.gt.agaaatgt.tggc.....c.a.a.aa.									
<i>Lb. rhamnosus</i> LMG6400Tt.....t.....g.t.t...c.....a									
<i>Lb. zeae</i> LMG17315Tt.....t.....g.....t.....t.....t.....									
<i>E. faecalis</i> LMG7937Tt.....ag...gt.a.aaatgt.agg.....c.aaca.ag									
P3M	-----									
P31	-----									
P3M	-----									
P1N	-----									
P11	-----									



	310		360	370	380	390	400			
P123r1	taggggaatcttcggcaagtggacgaaaagtctgaccgagcaacgccgctgagtgaaagaaggttttcggatcgtaaaactctgttggtagagaagaacgt									
P123r2									
P123r3									
E548Nt.....aa									
E548Mt.....aa									
E5412Mt.....aa									
M027Nt.....aa									
M027Mt.....aa									
M0331Mt.....aa									
M0331Nt.....aa									
107Nt.....aa									
106t.....aa									
1052t.....aa									
M030Mt.....aa									
1053t.....aa									
E. faecium DSM20477Tt.....aa									
E. durans DSM20633Tt.....aa									
E. hirae ATCC8043Tt.....aa									
E. ratti ATCC700914Tt.....aa									
E. villosorum LMG12287Tt.....aa									
E. mundtii ATCC43186Tt.....aa									
E. haemoperoxidus CCM4851Tt.....aa									
E. moraviensis CCM4856Tt.....aa									
L. lactis ATCC19435Tt.....aa									
L. lactis NCD02181Tt.....aa									
L. lactis ATCC19257Tt.....aa									
L. garviae LMG8893Tgg.-c.-cc.-n.....t.....aa									
L. piscium CCUG32732Tt.....aa									
L. plantarum NCD01869Tt.....aa									
L. raffinolactis NCD0617Tn.....n.....t.....aa									
B. subtilis NCD01769Tc.....g.....t.....g.....aa									
E. faecalis ATCC19433Tt.....aa									
	410	420	430	440	450	460	470	480	490	500
P123r1	tg-gt-gagagtggaagctcattcaagtgacggttaactaccagaa-aggagcggtaactacgtgccagcagccggttaacgtaggtcccagcgcgt									
P123r2									
P123r3									
E548N	g.-ac.tt..aactgaacg-..ccc.....t..a.....c.-cc.....gg.a.....									
E548M	g.-ac.tt..aactgaacg-..ccc.....t..a.....c.-cc.....gg.a.....									
E5412M	g.-ac.tt..aactgaacg-..ccc.....t..a.....c.-cc.....gg.a.....									
M027N	g.-ac.tt..aactgttca-c.cct.....t..a.....c.-cc.....gg.a.....									
M027M	g.-ac.tt..aactgttca-c.cct.....t..a.....c.-cc.....gg.a.....									
M0331M	g.-ac.tt..aactgttca-c.cct.....t..a.....c.-cc.....gg.a.....									
M0331N	g.-ac.tt..aactgttca-c.cct.....t..a.....c.-cc.....gg.a.....									
107N	g.-ac.tt..aactgttca-c.cct.....t..a.....c.-cc.....gg.a.....									
106	g.-ac.tt..aactgttca-c.cct.....t..a.....c.-cc.....gg.a.....									
1052	g.-ac.tt..aactgttca-c.cct.....t..a.....c.-cc.....gg.a.....									
M030M	g.-ac.tt..aactgttca-c.cct.....t..a.....c.-cc.....gg.a.....									
1053	g.-ac.tt..aactgttca-c.cct.....t..a.....c.-cc.....gg.a.....									
E. faecium DSM20477T	g.-a.....aactgttca-..cct.....t..a.....c.-cc.....gg.a.....									
E. durans DSM20633T	g.-a.....aactgttca-..cct.....t..a.....c.-cc.....gg.a.....									
E. hirae ATCC8043T	g.-a.....aactgttca-..cct.....t..a.....c.-cc.....gg.a.....									
E. ratti ATCC700914T	g.-a.....aactgttca-..cct.....t..a.....c.-cc.....n.gg.a.....									
E. villosorum LMG12287T	g.-a.....aactgttca-..cct.....t..a.....c.-cc.....gg.a.....									
E. mundtii ATCC43186T	g.-a.....aactgttca-..cct.....t..a.....c.-cc.....gg.a.....									
E. haemoperoxidus CCM4851T	a.tag.....aactgtct-..acct.....t..a.....c.-cc.....gg.a.....									
E. moraviensis CCM4856T	gt-ag.....aactgtct-..acct.....t..a.....c.-cc.....gg.a.....									
L. lactis ATCC19435Tt.....n.....									
L. lactis NCD02181Tt.....n.....									
L. lactis ATCC19257Tt.....n.....									
L. garviae LMG8893T	..-agt.....at.ac-.t.....t..na.....a.....									
L. piscium CCUG32732T	..-tgt.....at.ac-.a.....t..a.....n.....									
L. plantarum NCD01869T	ca-.....t.....ctg.....t..a.....n.....									
L. raffinolactis NCD0617T	..-cat.....at.at-g.....t..a.....n.....n.....n.....									
B. subtilis NCD01769T	gt-accgttcgaat.g.gcgg-..acct.....c..a.....c.-cc.....gg.a.....									
E. faecalis ATCC19433T	g.-ac.tt..aactgaacg-..ccc.....t..a.....c.-cc.....gg.a.....									
	510	520	530	540	550	560	570	580	590	600
P123r1	tgtccggatttattgggc-gtaaagcagagcgcaggtggtttattaaagtctggtgtaaaagcagtggtcaaccattgtatg-cattggaaactggtaga									
P123r2									
P123r3									
E548Nt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
E548Mt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
E5412Mt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
M027Nt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
M027Mt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
M0331Mt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
M0331Nt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
107Nt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
106t.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
1052t.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
M030Mt-tc.....c..c.....g..a..g...c.ccc..t.....ggg.agg.t.....g..									
1053t.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
E. faecium DSM20477Tt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
E. durans DSM20633Tt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
E. hirae ATCC8043Tt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
E. ratti ATCC700914Tt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
E. villosorum LMG12287Tt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
E. mundtii ATCC43186Tt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
E. haemoperoxidus CCM4851Tt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
E. moraviensis CCM4856Tt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
L. lactis ATCC19435Tt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
L. lactis NCD02181Tt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
L. lactis ATCC19257Tt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									
L. garviae LMG8893Tn.....c.....a.....a.....g.....g.....g.....									
L. piscium CCUG32732Tn.....c.....a.....a.....g.....g.....g.....									
L. plantarum NCD01869Tn.....c.....a.....a.....g.....g.....g.....									
L. raffinolactis NCD0617T	n.....n.....n.....a.....a.....n.....g.....t.....									
B. subtilis NCD01769T	..n.....a.....-a.....g.ct.....c.....c.....a..g...c.ccc.....ggg.agg.t.....gga.									
E. faecalis ATCC19433Tt.....c.....c.....a..g...c.ccc.....ggg.agg.t.....g..									



```

610          660          670          680          690          700
P123r1      cttgagtgcaggagaggaggtgaa-ttccatgtgtagc-ggtgaaatgcgt---agatatat-ggaggaaacaccggtggcgaagcggctctctggcc
P123r2
P123r3
E548N      .a.....-.....-.....-.....g.....a.....g.....t.....
E548M      .a.....-.....-.....-.....g.....a.....g.....t.....
E5412M     .a.....-.....g.....-.....-.....a.....a.....g.....t.....
M027N     .a.....-.....t.....-.....-.....a.....a.....g.....t.....
M027M     .a.....-.....t.....-.....-.....a.....g.....g.....t.....
M0331M    .a.....-.....t.....-.....-.....a.....g.....g.....t.....
M0331N    .a.....-.....t.....-.....-.....a.....g.....g.....t.....
107N      .a.....-.....t.....-.....-.....a.....g.....g.....t.....
106       .a.....-.....t.....-.....-.....a.....g.....g.....t.....
1052      .a.....-.....t.....-.....-.....a.....g.....g.....t.....
M030M     .a.....-.....t.c.....g.t.....-ttggat.....a.....g.....t.....
1053      .a.....-.....t.....-.....-.....a.....g.....g.....t.....
E. faecium DSM20477T .a.....-.....-.....-.....a.....g.....g.....t.....
E. durans DSM20633T .a.....-.....-.....-.....a.....g.....g.....t.....
E. hirae ATCC8043T .a.....-.....-.....-.....a.....g.....g.....t.....
E. ratti ATCC700914T .a.....-.....-.....-.....a.....g.....g.....t.....
E. villorum LMG12287T .a.....-.....-.....-.....a.....g.....g.....t.....
E. mundtii ATCC43186T .a.....-.....-.....-.....a.....g.....g.....t.....
E. haemoperoxidus CCM4851T .a.....-.....atg.t...tg-a.....a.....g.....a.....t.....
E. moraviensis CCM4856T .a.....-.....-.....-.....a.....g.....a.....t.....
L. lactis ATCC19435T .a.....-.....-.....-.....a.....g.....g.....t.....
L. lactis NCD02181T .a.....-.....-.....-.....a.....g.....g.....t.....
L. lactis ATCC19257T .a.....-.....-.....-.....a.....g.....g.....t.....
L. garviae LMG8893T .a.....-.....-.....-.....a.....g.....g.....t.....
L. piscium CCUG32732T .t.....-.....-.....-.....a.....g.....g.....t.....
L. plantarum NCD01869T .t.....-.....-.....-.....a.....g.....g.....t.....
L. raffinolactis NCD0617T .t.....-.....-.....-.....a.....g.....g.....t.....
B. subtilis NCD01769T .a.....-.....sub.c...n.....-g.g.....a.....g.....a.....t.....
E. faecalis ATCC19433T .a.....-.....-.....-.....a.....g.....g.....t.....

710          720          730          740          750          760          770          780          790          800
P123r1      tgtaaactgacactgaggctcnaagcgtggggagcaaacaggattagataccctggtagtcca-cgccgtaaacgatgagtgctagatgta-gggagcta
P123r2
P123r3
E548N      .g.....-.....-.....-.....ag...tg...ag.g.t.....
E548M      .g.....-.....-.....-.....ag...tg...ag.g.t.....
E5412M     .g.....-.....-.....-.....ag...t...ag.g.t.....
M027N     .g.....-.....-.....c.....ag...t...ag.g.t.....
M027M     .g.....-.....-.....c.....ag...t...ag.g.t.....
M0331M    .g.....-.....-.....c.....ag...t...ag.g.t.....
M0331N    .g.....-.....-.....c.....ag...t...ag.g.t.....
107N      .g.....-.....-.....c.....ag...t...ag.g.t.....
106       .g.....-.....-.....c.....ag...t...ag.g.t.....
1052      .g.....-.....-.....c.....ag...t...ag.g.t.....
M030M     .g.....-.....-.....c.....ag...tg...ag.g.t.....
1053      .g.....-.....-.....c.....ag...t...ag.g.t.....
E. faecium DSM20477T .g.....-.....-.....-.....ag...t...ag.g.t.....
E. durans DSM20633T .g.....-.....-.....-.....ag...t...ag.g.t.....
E. hirae ATCC8043T .g.....-.....-.....-.....ag...t...ag.g.t.....
E. ratti ATCC700914T .g.....-.....-.....-.....ag...t...ag.g.t.....
E. villorum LMG12287T .g.....-.....-.....-.....ag...t...ag.g.t.....
E. mundtii ATCC43186T .g.....-.....-.....-.....ag...t...ag.g.t.....
E. haemoperoxidus CCM4851T .g.....-.....-.....-.....ag...t...ag.g.t.....
E. moraviensis CCM4856T .g.....-.....-.....m.....ag...t...ag.g.t.....
L. lactis ATCC19435T .g.....-.....-.....-.....ag...t...ag.g.t.....
L. lactis NCD02181T .g.....-.....-.....-.....ag...t...ag.g.t.....
L. lactis ATCC19257T .g.....-.....-.....-.....ag...t...ag.g.t.....
L. garviae LMG8893T .g.....-.....-.....n.....c.....t.....t.g....
L. piscium CCUG32732T .t.....-.....-.....-.....n.....t...t-t-g....
L. plantarum NCD01869T .n.....n.....-.....t...n.....-.....n.....n.....n.....t...t-t-g....
L. raffinolactis NCD0617T .c.....-.....-.....n.....-.....n.....-.....n.....t...t-t-g....
B. subtilis NCD01769T .g.....-.....ag.....g.....-.....ag...t-a...g.g.t.....
E. faecalis ATCC19433T .g.....-.....-.....-.....ag...t...ag.g.t.....

810          820          830          840          850          860          870          880          890          900
P123r1      t-aagttctctgtatcgcagctaacgcgaataaagcactccgcctggggagtagcagccgcaaggttgaaactcaaaggaattgacgggggcongca-acaagc
P123r2
P123r3
E548N      .ccgccct.a.gct...a...t.....
E548M      .ccgccct.a.gct...a...t.....
E5412M     .ccgccct.a.gct...a...t.....
M027N     .ccgccct.a.gct...t.....
M027M     .ccgccct.a.gct...t.....
M0331M    .ccgccct.a.gct...t.....
M0331N    .ccgccct.a.gct...t.....
107N      .ccgccct.a.gct...t.....
106       .ccgccct.a.gct...t.....
1052      .ccgccct.a.gct...t.....
M030M     .ccgccct.a.gct...t.....
1053      .ccgccct.a.gct...t.....
E. faecium DSM20477T .ccgccct.a.gct...t.....
E. durans DSM20633T .ccgccct.a.gct...t.....
E. hirae ATCC8043T .ccgccct.a.gct...t.....
E. ratti ATCC700914T .ccgccct.a.gct...t.....
E. villorum LMG12287T .ccgccct.a.gct...t.....
E. mundtii ATCC43186T .ccgccct.a.gct...t.....
E. haemoperoxidus CCM4851T .ccgccct.a.gct...t.....
E. moraviensis CCM4856T .ccgccct.a.gct...t.....
L. lactis ATCC19435T .ccgccct.a.gct...t.....
L. lactis NCD02181T .ccgccct.a.gct...t.....
L. lactis ATCC19257T .ccgccct.a.gct...t.....
L. garviae LMG8893T .ccgccct.a.gct...t.....
L. piscium CCUG32732T .cc.cc.aa.ga...a...t.....
L. plantarum NCD01869T .cc.cc.nna.ga...a...t.....
L. raffinolactis NCD0617T .cc.cc.na.ga...n...n.....t.....
B. subtilis NCD01769T .ccgccct.a.gct...t.n.....gt...ac.....
E. faecalis ATCC19433T .ccgccct.a.gct...a...t.....

```



910 960 970 980 990 1000

ggtggagcatgtggttta-attogaagcaacgcgaa-gaaccttaccaggtcttg-acatactcgtgctattcctagataggaagt-tccttcgggac

P123r1
P123r2
P123r3
E548N
E548M
E5412M
M027N
M027M
M0331M
M0331N
107N
106
1052
M030M
1053
E. faecium DSM20477T
E. durans DSM20633T
E. hirae ATCC8043T
E. ratti ATCC700914T
E. villorum LMG12287T
E. mundtii ATCC43186T
E. haemoperoxidus CCM4851T
E. moraviensis CCM4856T
L. lactis ATCC19435T
L. lactis NCD02181T
L. lactis ATCC19257T
L. garviae LMG8893T
L. piscium CCUG32732T
L. plantarum NCD01869T
L. raffinolactis NCD0617T
B. subtilis NCD01769T
E. faecalis ATCC19433T

910 1010 1020 1030 1040 1050 1060 1070 1080 1090 1100

acgggatcacagtggtgcatggtgtcgtcagctcgtgctgagatggtgggtaagtcccgaacgagcgcgaacccctattgtagttgccatcatta

P123r1
P123r2
P123r3
E548N
E548M
E5412M
M027N
M027M
M0331M
M0331N
107N
106
1052
M030M
1053
E. faecium DSM20477T
E. durans DSM20633T
E. hirae ATCC8043T
E. ratti ATCC700914T
E. villorum LMG12287T
E. mundtii ATCC43186T
E. haemoperoxidus CCM4851T
E. moraviensis CCM4856T
L. lactis ATCC19435T
L. lactis NCD02181T
L. lactis ATCC19257T
L. garviae LMG8893T
L. piscium CCUG32732T
L. plantarum NCD01869T
L. raffinolactis NCD0617T
B. subtilis NCD01769T
E. faecalis ATCC19433T

1110 1120 1130 1140 1150 1160 1170 1180 1190 1200

agttgggc-actctaagcagagactgccggtgataaacgggagg-aaggtggggatgacgtcaaatcatcatgcccttatgacctgggctacacacgtgct

P123r1
P123r2
P123r3
E548N
E548M
E5412M
M027N
M027M
M0331M
M0331N
107N
106
1052
M030M
1053
E. faecium DSM20477T
E. durans DSM20633T
E. hirae ATCC8043T
E. ratti ATCC700914T
E. villorum LMG12287T
E. mundtii ATCC43186T
E. haemoperoxidus CCM4851T
E. moraviensis CCM4856T
L. lactis ATCC19435T
L. lactis NCD02181T
L. lactis ATCC19257T
L. garviae LMG8893T
L. piscium CCUG32732T
L. plantarum NCD01869T
L. raffinolactis NCD0617T
B. subtilis NCD01769T
E. faecalis ATCC19433T



1210 1260 1270 1280 1290 1300
 P123r1
 P123r2
 P123r3
 E548N
 E548M
 E5412M
 M027N
 M027M
 M0331M
 M0331N
 M07N
 M06
 M052
 M030M
 M053
 E. faecium DSM20477T
 E. durans DSM20633T
 E. hirae ATCC8043T
 E. ratti ATCC700914T
 E. villorum LMG12287T
 E. mundtii ATCC43186T
 E. haemoperoxidus CCM4851T
 E. moraviensis CCM4856T
 L. lactis ATCC19435T
 L. lactis NCD02181T
 L. lactis ATCC19257T
 L. garviae LMG8893T
 L. piscium CCUG32732T
 L. plantarum NCD01869T
 L. raffinolactis NCD0617T
 B. subtilis NCD01769T
 E. faecalis ATCC19433T

1310 1320 1330 1340 1350 1360 1370 1380 1390 1400
 P123r1
 P123r2
 P123r3
 E548N
 E548M
 E5412M
 M027N
 M027M
 M0331M
 M0331N
 M07N
 M06
 M052
 M030M
 M053
 E. faecium DSM20477T
 E. durans DSM20633T
 E. hirae ATCC8043T
 E. ratti ATCC700914T
 E. villorum LMG12287T
 E. mundtii ATCC43186T
 E. haemoperoxidus CCM4851T
 E. moraviensis CCM4856T
 L. lactis ATCC19435T
 L. lactis NCD02181T
 L. lactis ATCC19257T
 L. garviae LMG8893T
 L. piscium CCUG32732T
 L. plantarum NCD01869T
 L. raffinolactis NCD0617T
 B. subtilis NCD01769T
 E. faecalis ATCC19433T

1410
 P123r1
 P123r2
 P123r3
 E548N
 E548M
 E5412M
 M027N
 M027M
 M0331M
 M0331N
 M07N
 M06
 M052
 M030M
 M053
 E. faecium DSM20477T
 E. durans DSM20633T
 E. hirae ATCC8043T
 E. ratti ATCC700914T
 E. villorum LMG12287T
 E. mundtii ATCC43186T
 E. haemoperoxidus CCM4851T
 E. moraviensis CCM4856T
 L. lactis ATCC19435T
 L. lactis NCD02181T
 L. lactis ATCC19257T
 L. garviae LMG8893T
 L. piscium CCUG32732T
 L. plantarum NCD01869T
 L. raffinolactis NCD0617T
 B. subtilis NCD01769T
 E. faecalis ATCC19433T

Appendix H - References

- Abo-Elnaga, I.G. and Kandler, O.** (1965). Zur taxonomie der gattung *Lactobacillus* Beijerinck. I. Das subgenus Streptobacterium Orla-Jensen. Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene. Abteilung II, **119**, 1-36. Approved Lists of Bacterial Names. International Journal of Systematic Bacteriology, 1980, **30**, 225-420.
- Aguirre, M. and Collins, M.D.** (1992). Phylogenetic analysis of *Alloiococcus otitis* gen. nov., sp. nov., an organism from human middle ear fluid. International Journal of Systematic Bacteriology, 1992, **42**, 79-83.
- Aguirre, M. and Collins, M.D.** (1992). Phylogenetic analysis of some Aerococcus-like organisms for urinary tract infections: description of *Aerococcus urinae* sp. nov. Journal of General Microbiology, **138**, 401-405.
- Aguirre, M., Morrison, D., Cookson, B.D., Gay, F.W. and Collins, M.D.** (1993). Phenotypic and phylogenetic characterization of some *Gemella*-like organisms from human infections: description of *Dolosigranulum pigrum* gen. nov., sp. nov. Journal of Applied Bacteriology, **75**, 608-612. Validation List N° 49. International Journal of Systematic Bacteriology, **44**, 1994, 370-371.
- Andrewes, F.W. and Horder, T.J.** (1906). A study of the streptococci pathogenic for man. Lancet, **2**, 708-713.
- Antunes, A., Rainey, F.A., Nobre, M.F., Schumann, P., Ferreira, A.M., Ramos, A., Santos, H. and Da Costa, M.S.** (2002). *Leuconostoc ficulneum* sp. nov., a novel lactic acid bacterium isolated from a ripe fig, and reclassification of *Lactobacillus fructosus* as *Leuconostoc fructosum* comb. nov. International Journal of Systematic and Evolutionary Microbiology, 2002, **52**, 647-655.
- Arbique, J.C., Poyart, C., Trieu-Cuot, P., Quesne, G., Carvalho, M da G.S., Steigerwalt, A.G., Morey, R.E., Jackson, D., Davidson, R.J. and Facklam, R.R.** (2004). Accuracy of phenotypic and genotypic testing for identification of *Streptococcus pneumoniae* and description of *Streptococcus pseudopneumoniae* sp. nov. Journal of Clinical Microbiology, **42**, 4686-4696. Validation List N° 101. International Journal of Systematic and Evolutionary Microbiology, 2005, **55**, 1-2.
- Aslam, Z., Im, W.T., Ten, L.M., Lee, M.J., Kim, K.H. and Lee, S.T.** (2006). *Lactobacillus siliginis* sp. nov., isolated from wheat sourdough in South Korea. International Journal of Systematic and Evolutionary Microbiology, **56**, 2209-2213.
- Back, W.** (1978). Elevation of *Pediococcus cerevisiae* subsp. *dextrinicus* (Coster and White) comb. nov. International Journal of Systematic Bacteriology, **28**, 523-527.
- Back, W.** (1978). Zur Taxonomie der gattung *Pediococcus*. Brauwissenschaft, **31**, 237-250, 312-320, 336-343. Validation List N° 25. International Journal of Systematic Bacteriology, 1988, **38**, 220-222.
- Back, W., Bohak, I., Ehrmann, M., Ludwig, W. and Schleifer, K.H.** (1996). Revival of the species *Lactobacillus lindneri* and the design of a species specific oligonucleotide probe. Applied Microbiology, **19**, 322-325. Validation List N° 61. International Journal of Systematic Bacteriology, 1997, **47**, 601-602.
- Back, W., Bohak, I., Ehrmann, M., Ludwig, W., Pot, B., Kersters, K. and Schleifer, K.H.** (1999). *Lactobacillus perolens* sp. nov., a soft drink spoilage bacterium. Systematic and Applied Microbiology, **22**, 354-359. International Journal of Systematic and Evolutionary Microbiology, 2000, **50**, 3-4.
- Baele, M., Vancanneyt, M., Devriese, L.A., Lefebvre, K., Swings, J. and Haesebrouck, F.** (2003). *Lactobacillus ingluviei* sp. nov., isolated from the intestinal tract of pigeons. International Journal of Systematic and Evolutionary Microbiology, **53**, 133-136.
- Bailey, L. and Collins, M.D.** (1982). Reclassification of "*Streptococcus pluton*" (White) in a new genus *Melissococcus* as *Melissococcus pluton* nom. rev.; comb. nov. Journal of Applied Bacteriology, **53**, 215-217. Validation List N° 11. International Journal of Systematic Bacteriology, 1983, **33**, 672-674.
- Barnes, E. M., Impey, C.S., Stevens, B.J.H. and Peel, J.L.** (1977). *Streptococcus pleomorphus* sp. nov.: an anaerobic streptococcus isolated from the caeca of birds. Journal of General Microbiology, **102**, 45-53. Validation List N° 2. International Journal of Systematic Bacteriology, 1979, **29**, 79-80.
- Beck, R., Weiss, N. and Winter, J.** (1988). *Lactobacillus graminis* sp. nov., a new species of facultatively heterofermentative lactobacilli surviving at low pH in grass silage. Systematic and Applied Microbiology, **10**, 279-283. Validation List N° 28. International Journal of Systematic Bacteriology, 1989, **39**, 93-94.

- Beighton, D., Hayday, H., Russell, R.K.B. and Whitey, K.A.** (1984). *Streptococcus macacae* sp. nov., from dental plaque of monkeys (*Macaca fascicularis*). International Journal of Systematic Bacteriology, **34**, 332-335.
- Beijerinck, M.W.** (1901). Sur les ferments lactiques de l'industrie. Archives Néerlandaises des Sciences Exactes et Naturelles (Section 2), **6**, 212-243.
- Beijerinck, M.W.**(1912). Mutation bei Mikroben. Folia Mikrobiologiya (Delft), **1**, 4-100.
- Bekal, S., Gaudreau, C., Luarence, R. A., Simoneau, E. and Raynal, L.** (2006). *Streptococcus pseudoporcinus* sp. nov., a novel species isolated from the genitourinary tract of women. Journal of Clinical Microbiology, **44**, 2584-2586.
- Bergey, D.H., Breed, R.S., Hammer, B.W., Huntoon, F.M., Murray, E.G.D. and Harrison, F.C.** (Eds.). (1934). Bergey's Manual of Determinative Bacteriology, 4th Ed., The Williams and Wilkins Co, Baltimore, pp. 1-664.
- Bergey, D.H., Harrison, F.C., Breed, R.S., Hammer, B.W. and Huntoon, F.M.** (1923). Bergey's Manual of Determinative Bacteriology, 1st ed., The Williams and Wilkins Co, Baltimore, pp. 1- 442.
- Bergey, D.H., Harrison, F.C., Breed, R.S., Hammer, B.W. and Huntoon, F.M.** (Eds.). (1925). Bergey's Manual of Determinative Bacteriology, 2nd ed., The Williams and Wilkins Co, Baltimore, pp. 1-462.
- Björkroth, K.J., Geisen, R., Schillinger, U., Weiss, N., De Vos, P., Holzapfel, W.H., Korkeala, H.J. and Vandamme, P.** (2000). Characterization of *Leuconostoc gasicomitatum* sp. nov., associated with spoiled raw tomato-marinated broiler meat strips packaged under modified-atmosphere conditions. Applied and Environmental Microbiology, **66**, 3764-3772. Validation List N° 79. International Journal of Systematic and Evolutionary Microbiology, 2001, **51**, 263-265.
- Björkroth, K.J., Schillinger, U., Geisen, R., Weiss, N., Hoste, B., Holzapfel, W.H., Korkeala, H. J. And Vandamme, P.** (2002). Taxonomic study of *Weissella confusa* and description of *Weissella cibaria* sp. nov., detected in food and clinical samples. International Journal of Systematic and Evolutionary Microbiology, **52**, 141-148.
- Bohak, I., Back, W., Richter, L., Ehrmann, M., Ludwig, W. and Schleifer, K.H.** (1998). *Lactobacillus amyolyticus* sp. nov., isolated from beer malt and beer wort. Systematic and Applied Microbiology, **21**, 360-364. Validation List N° 68. International Journal of Systematic Bacteriology, 1999, **49**, 1-3.
- Bouvet, A., Grimont, F. and Grimont, P.A.D.** (1989). *Streptococcus defectivus* sp. nov. and *Streptococcus adjacens* sp. nov., nutritionally variant streptococci from human clinical specimens. International Journal of Systematic Bacteriology, **39**, 290-294.
- Bridge, P.D. and Sneath, P.H.A.** (1982). *Streptococcus gallinarum* sp. nov. and *Streptococcus oralis* sp. nov. International Journal of Systematic Bacteriology, **32**, 410-415.
- Bringel, F., Castioni, A., Olukoya, D.K., Felis, G.E., Torriani, S. and Delaglio, F.** (2005). *Lactobacillus plantarum* subsp. *argenteratensis* subsp. nov., isolated from vegetable matrices. International Journal of Systematic and Evolutionary Microbiology, **55**, 1629- 1634.
- Brooker, J.D., O'Donovan, L.A., Skene, I., Clarke, K., Blackall, L. and Muslera, P.** (1994). *Streptococcus caprinus* sp. nov., a tannin-resistant ruminal bacterium from feral goats. Letters in Applied Microbiology, **18**, 313-318.
- Brygoo, E.R. and Aladame, N.** (1953). Étude d'une espèce nouvelle anaérobie stricte du genre Eubacterium : *E. crispatum* n. sp. Annales de l'Institut Pasteur (Paris), **84**, 640-641.
- Cai, Y., Okada, H., Mori, H., Benno, Y. and Nakase, T.** (1999). *Lactobacillus paralimentarius* sp. nov., isolated from sourdough. International Journal of Systematic Bacteriology, **49**, 1451-1455.
- Carr, G. and Davies, P.A.** (1972). The ecology and classification of *Lactobacillus collinoides* nov., spec.: A bacterium commonly found in fermenting apple juice. Journal of Applied Bacteriology, **35**, 463-471.
- Carr, J. and Davies, P.A.** (1970). Homofermentative lactobacilli of ciders including *Lactobacillus mali* nov. spec. Journal of Applied Bacteriology, **33**, 768-774.
- Carvalho, M.G.S., Shewmaker, P.L., Steigerwalt, A.G., Morey, R.E., Sampson, A.J., Joyce, K., Barrett, T. J., Teixeira, L.M. and Facklam, R.R.** (2006). *Enterococcus caccae* sp. nov., isolated from human stools. International Journal of Systematic and Evolutionary Microbiology, **56**, 1505-1508.

- Cato, E.P.** (1983). Transfer of *Peptostreptococcus parvulus* (Weinberg, Nativelle, and Prévot) Smith 1957. *Streptococcus parvulus* (Weinberg, Nativelle, and Prévot 1937) comb. nov., emend. International Journal of Systematic Bacteriology, **33**, 82-84.
- Chambel, L., Chelo, I.M., Zé-zé, L., Pedro, L.G., Santos, M.A. and Tenreiro, R.** (2006). *Leuconostoc pseudoficulneum* sp. nov., isolated from a ripe fig. International Journal of Systematic and Evolutionary Microbiology, **56**, 1375-1381.
- Charlton, D.B., Nelson, M.E. and Werkman, C.H.** (1934). Physiology of *Lactobacillus fructivorans* sp. nov. isolated from spoiled salad dressing. Iowa State Journal of Science, **9**, 1-11.
- Chatton, E. and Perard, C.** (1913). Schizophytes du caecum du cobaye. I. *Oscillospira guilliermondi* n.g., n.sp. Comptes Rendus des Séances de la Société de Biologie (Paris), **74**, 1159-1162
- Chenoll, E., Carmen Macian, M. and Aznar, R.** (2006). *Lactobacillus tucseti* sp. nov., a new lactic acid bacterium isolated from sausage. Systematic and Applied Microbiology, **29**, 389-395.
- Chenoll, E., Macian, M. C. and Aznar, R.** (2006). *Lactobacillus rennini* sp. nov., isolated rennin and associated with cheese spoilage. International Journal of Systematic and Evolutionary Microbiology, **56**, 449-452.
- Chester, F.D.** (1901). A manual of Determinative Bacteriology, The MacMillan Co., New York, 1901, pp. 1-401.
- Choi, H.J., Cheigh, C.I., Kim, S.B., Lee, D.W., Choi, S.W., Park, J.M. and Pyun, Y.R.** (2002). *Weissella kimchii* sp. nov., a novel lactic acid bacterium from *kimchi*. International Journal of Systematic and Evolutionary Microbiology, **52**, 507-511.
- Claussen, N.H.** (1903). Études sur les bactéries dites sarcines et sur les maladies qu'elles provoquent dans la bière. Compte Rendu des Travaux du Laboratoire de Carlsberg, 1903, **6**, 64-83.
- Collins, M.D. and Lawson, P.** (2000). The genus *Abiotrophia* (Kawamura *et al.*) is not monophyletic: proposal of *Granulicatella* gen. nov., *Granulicatella adiacens* comb. nov., *Granulicatella elegans* comb. nov. and *Granulicatella balaenopterae* comb. nov. International Journal of Systematic and Evolutionary Microbiology, **50**, 365-369.
- Collins, M.D., Aguirre, M., Facklam, R.R., Shallcross, J. and Williams, A.M.** (1992). *Globicatella sanguinis* gen. nov., sp. nov., a new gram-positive catalase-negative bacterium from human sources. Journal of Applied Bacteriology, **73**, 433-437. Validation List N° 53. International Journal of Systematic Bacteriology, 1995, **45**, 418-419.
- Collins, M.D., Asch, C., Farrow, J.A.E., Wallbanks, S. and Williams, A.M.** (1990). 16S ribosomal ribonucleic acid sequence analyses of lactococci and related taxa. Description of *Vagococcus fluvialis* gen. nov., sp. nov. Journal of Applied Bacteriology, **67**, 453-460.
- Collins, M.D., Facklam, R.R., Farrow, J.A.E. and Williamson, R.** (1989). *Enterococcus raffinosus* sp. nov., *Enterococcus solitarius* sp. nov. and *Enterococcus pseudoavium* sp. nov. FEMS Microbiology Letters, **57**, 283-288.
- Collins, M.D., Falsen, E., Lemozy, J., Åkervall, E., Sjöden, B. and Lawson, P.A.** (1997). Phenotypic and phylogenetic characterization of some *Globicatella*-like organisms from human sources: description of *Facklamia hominis* gen. nov., sp. nov. International Journal of Systematic Bacteriology, **47**, 880-882.
- Collins, M.D., Farrow, J.A.E. and Jones, D.** (1986). *Enterococcus mundtii* sp. nov. International Journal of Systematic Bacteriology, **36**, 8-12.
- Collins, M.D., Farrow, J.A.E., Katic, V. and Kandler, O.** (1985). Taxonomic studies on streptococci of serological groups E, P, U and V: description of *Streptococcus porcinius* sp. nov. Systematic and Applied Microbiology, **5**, 402-413.
- Collins, M.D., Farrow, J.A.E., Phillips, B.A., Fergus, S. and Jones, D.** (1987). Classification of *Lactobacillus divergens*, *Lactobacillus piscicola*, and some catalase-negative, asporogenous, rod-shaped bacteria from poultry in a new genus, *Carnobacterium*. International Journal Systematic Bacteriology, **37**, 310-316.
- Collins, M.D., Farrow, J.A.E., Phillips, B.A., Fergus, S. and Jones, D.** (1987). Classification of *Lactobacillus divergens*, *Lactobacillus piscicola*, and some catalase-negative, asporogenous, rod-shaped bacteria from poultry in a new genus, *Carnobacterium*. International Journal of Systematic Bacteriology, **37**, 310-316.

- Collins, M.D., Higgins, R., Messier, S., Fortin, M., Hutson, R.A., Lawson, P.A. and Falsen, E.** (2003). *Allofustis seminis* gen. nov., sp. nov., a novel Gram-positive, catalase-negative, rod-shaped bacterium from pig semen. *International Journal of Systematic and Evolutionary Microbiology*, **53**, 811-814.
- Collins, M.D., Hutson, R.A., Falsen, E. and Sjöden, B.** (1999). *Facklamia sourekii* sp. nov., isolated from human sources. *International Journal of Systematic Bacteriology*, **49**, 635-638.
- Collins, M.D., Hutson, R.A., Falsen, E. and Sjöden, B.** (1999). *Facklamia tabacinasalis* sp. nov., from powdered tobacco. *Journal of Clinical Microbiology*, **49**, 1247-1250.
- Collins, M.D., Hutson, R.A., Falsen, E., Inganäs, E. and Bisgaard, M.** (2002). *Streptococcus gallinaceus* sp. nov., from chickens. *International Journal of Systematic and Evolutionary Microbiology*, **52**, 1161-1164.
- Collins, M.D., Hutson, R.A., Falsen, E., Nikolaitchouk, N., Laclaire, L. and Facklam, R.R.** (2000). An unusual *Streptococcus* from human urine, *Streptococcus urinalis* sp. nov. *International Journal of Systematic and Evolutionary Microbiology*, **50**, 1173-1178.
- Collins, M.D., Hutson, R.A., Foster, G., Falsen, E. and Weiss, N.** (2002). *Isobaculum melis* gen. nov., sp. nov., a *Carnobacterium*-like organism isolated from the intestine of a badger. *International Journal of Systematic and Evolutionary Microbiology*, **52**, 207-210.
- Collins, M.D., Hutson, R.A., Hoyles, L., Falsen, E., Nikolaitchouk, N. and Foster, G.** (2001). *Streptococcus ovis* sp. nov., isolated from sheep. *International Journal of Systematic and Evolutionary Microbiology*, **51**, 1147-1150.
- Collins, M.D., Jones, D., Farrow, J.A.E., Kilpper-Bälz, R. and Schleifer, K.H.** (1984). *Enterococcus avium* nom. rev., comb. nov.; *E. casseliflavus* nom. rev., comb. nov.; *E. durans* nom. rev., comb. nov.; *E. gallinarum* comb. nov.; and *E. malodoratus* sp. nov. *International Journal of Systematic Bacteriology*, **34**, 220-223.
- Collins, M.D., Lawson, P.A., Monasterio, R., Falsen, E., Sjöden, B. and Facklam, R.R.** (1998). *Facklamia ignava* sp. nov., isolated from human clinical specimens. *Journal of Clinical Microbiology*, **36**, 2146-2148.
- Collins, M.D., Lawson, P.A., Monasterio, R., Falsen, E., Sjöden, B. and Facklam, R.R.** (1999). *Ignavigranum ruoffiae* sp. nov., isolated from human clinical specimens. *International Journal of Systematic Bacteriology*, **49**, 97-101.
- Collins, M.D., Lundström, T., Welinder-Olsson, C., Hansson, I., Wattle, O., Hutson, R.A. and Falsen, E.** (2004). *Streptococcus devriesei* sp. nov., from equine teeth. *Systematic and Applied Microbiology*, **27**, 146-150.
- Collins, M.D., Phillips, B.A. and Zanoni, P.** (1989). Deoxyribonucleic acid homology studies of *Lactobacillus casei*, *Lactobacillus paracasei* sp. nov., subsp. *paracasei* and subsp. *tolerans*, and *Lactobacillus rhamnosus* sp. nov., comb. nov. *International Journal of Systematic Bacteriology*, **39**, 105-108.
- Collins, M.D., Rodrigues, U.M., Pigott, N.E. and Facklam, R.R.** (1991). *Enterococcus dispar* sp. nov. a new *Enterococcus* species from human sources. *Letters in Applied Microbiology*, **12**, 95-98.
- Collins, M.D., Rodriguez Jovita, M., Hutson, R.A., Falsen, E., Sjöden, B. and Facklam, R.R.** (1999). *Dolosicoccus paucivorans* gen. nov., sp. nov., isolated from human blood. *International Journal of Systematic Bacteriology*, **49**, 1439-1442.
- Collins, M.D., Rodriguez Jovita, M., Lawson, P.A., Falsen, E. and Foster, G.** (1999). Characterization of a novel Gram-positive, catalase-negative coccus from horses: description of *Eremococcus coleocola* gen. nov., sp. nov. *International Journal of Systematic Bacteriology*, **49**, 1381-1385.
- Collins, M.D., Samelis, J., Metaxopoulos, J. and Wallbanks, S.** (1993). Taxonomic studies on some *Leuconostoc*-like organisms from fermented sausages: description of a new genus *Weissella* for the *Leuconostoc paramesenteroides* group of species. *Journal of Applied Bacteriology*, **75**, 595-603. Validation List N° 49. *International Journal Systematic Bacteriology*, 1994, **44**, 370 -371.
- Collins, M.D., Williams, A.M. and Wallbanks, S.** (1993). The phylogeny of *Aerococcus* and *Pediococcus* as determined by 16S rRNA sequence analysis: description of *Tetragenococcus halophilus* gen. nov. *FEMS Microbiology Letters*, **70**, 255-262.
- Corsetti, A., Settani, L., Van Sinderen, D., Felis, G.E., Dellaglio, F. and Gobbbetti, M.** (2005). *Lactobacillus rossiae* sp. nov., isolated from wheat sourdough. *International Journal of Systematic and Evolutionary Microbiology*, **55**, 35- 40.

- Coster, E. and White, H.R.** (1964). Further studies of the genus *Pediococcus*. *Journal of General Microbiology*, **37**, 15-31.
- Coykendall, A.L.** (1974). Four types of *Streptococcus mutans* based on their genetic, antigenic and biochemical characteristics. *Journal of General Microbiology*, **83**, 327-338.
- Coykendall, A.L.** (1977). Proposal to elevate the subspecies of *Streptococcus mutans* to species status, based on their molecular composition. *International Journal of Systematic Bacteriology*, **27**, 26-30.
- Coykendall, A.L.** (1983). *Streptococcus sobrinus* nom. rev. and *Streptococcus ferus* nom. rev.: habitat of these and other mutans streptococci. *International Journal of Systematic Bacteriology*, **33**, 883-885.
- Curk, M.C., Hubert, J.C. and Bringel, F.** (1996). *Lactobacillus paraplantarum* sp. nov., a new species related to *Lactobacillus plantarum*. *International Journal of Systematic Bacteriology*, **46**, 595-598.
- De Graef, E.M., Devriese, L.A., Vancanneyt, M., Baele, M., Collins, M.D., Lefebvre, K., Swings, J. and Haesebrouck, F.** (2003). Description of *Enterococcus canis* sp. nov. from dogs and reclassification of *Enterococcus porcinus* Teixeira *et al.* 2001 as a later synonym of *Enterococcus villorum* Vancanneyt *et al.* 2001. *International Journal of Systematic and Evolutionary Microbiology*, **53**, 1069-1074.
- De Vaux, A., Laguerre, G., Diviés, C. and Prévost, H.** (1998). *Enterococcus asini* sp. nov. isolated from the caecum of donkeys (*Equus asinus*). *International Journal of Systematic Bacteriology*, **48**, 383-387.
- Dellaglio, F., Felis, G.E., Castioni, A., Torriani, S. and Germond, J.E.** (2005). *Lactobacillus delbrueckii* subsp. *indicus* subsp. nov., isolated from Indian dairy products. *International Journal of Systematic and Evolutionary Microbiology*, **55**, 401-404.
- Dellaglio, F., Torriani, S. and Felis, G.E.** (2004). Reclassification of *Lactobacillus cellobiosus* Rogosa *et al.* 1953 as a later synonym of *Lactobacillus fermentum* Beijerinck 1901. *International Journal of Systematic and Evolutionary Microbiology*, **54**, 809-812.
- Dent, V.E. and Williams, R.A.D.** (1982). *Lactobacillus animalis* sp. nov., a new species of *Lactobacillus* from the alimentary canal of animals. *Zentralbl. Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene, Abteilung 1, Originalarbeiten*, **C3**, 377-386. Validation List N° 10. *International Journal of Systematic Bacteriology*, 1983, **33**, 438-440.
- Devriese, L.A., Ceysens, K., Rodrigues, U.M. and Collins, M.D.** (1990). *Enterococcus columbae*, a species from pigeon intestines. *FEMS Microbiology Letters*, **71**, 247-252. Validation N° 44. *International Journal of Systematic Bacteriology*, 1993, **43**, 188-189.
- Devriese, L.A., Dutta, G.N., Farrow, J.A.E., Van de Kerckhove, A. and Phillips, B.A.** (1983). *Streptococcus cecorum*, a new species isolated from chickens. *International Journal of Systematic Bacteriology*, **33**, 772-776.
- Devriese, L.A., Kilpper-Bälz, R. and Schleiffer, K.H.** (1988). *Streptococcus hyointestinalis* sp. nov. from the gut of swine. *International Journal of Systematic Bacteriology*, **38**, 440-441.
- Devriese, L.A., Pot, B., Vandamme, P., Kersters, K., Collins, M.D., Alvarez, N., Haesebrouck, F. and Hommez, J.** (1997). *Streptococcus hyovaginalis* sp. nov. and *Streptococcus thoralensis* sp. nov., from the genital tract of sows. *International Journal of Systematic Bacteriology*, **47**, 1073-1077.
- Devriese, L.A., Vandamme, P., Collins, M.D., Alvarez, N., Pot, B., Hommez, J., Butaye, P. and Haesebrouck, F.** (1999). *Streptococcus pluranimalium* sp. nov., from cattle and other animals. *International Journal of Systematic Bacteriology*, **49**, 1221-1226.
- Devriese, L.A., Hommez, J., Kilpper-Bälz, R. and Schleiffer, K.H.** (1986). *Streptococcus canis* sp. nov.: a species of group G streptococci from animals. *International Journal of Systematic Bacteriology*, **36**, 422-425.
- Devriese, L.A., Pot, B., Vandamme, P., Kersters, K., Collins, M.D., Alvarez, N., Haesebrouck, F. and Hommez, J.** (1997). *Streptococcus hyovaginalis* sp. nov. and *Streptococcus thoralensis* sp. nov., from the genital tract of sows. *International Journal of Systematic Bacteriology*, **47**, 1073-1077.
- Dicks, L.M.T., Dellaglio, F. and Collins, M.D.** (1995). Proposal to reclassify *Leuconostoc oenos* as *Oenococcus oeni* [corrig.] gen. nov., comb. nov. *International Journal of Systematic Bacteriology*, **45**, 395-397.
- Dicks, L.M.T., Du Plessis, E.M., Dellaglio, F. and Lauer, E.** (1996). Reclassification of *Lactobacillus casei* subsp. *casei* ATCC 393 and *Lactobacillus rhamnosus* ATCC 15820 as *Lactobacillus zaeae* nom. rev., designation of ATCC 334

as the neotype of *L. casei* subsp. *casei*, and rejection of the name *Lactobacillus paracasei*. International Journal of Systematic Bacteriology, **46**, 337- 340.

Dicks, L.M.T., Fantuzzi, L., Gonzalez, F.C., Du Toit, M. and Dellaglio, F. (1993). *Leuconostoc argentinum* sp. nov., isolated from argentine raw milk. International Journal of Systematic Bacteriology, **43**, 347-351.

Dicks, L.M.T., Silvester, M., Lawson, P.A. and Collins, M.D. (2000). *Lactobacillus fornicalis* sp. nov. , isolated from the posterior fornix of the human vagina. International Journal of Systematic and Evolutionary Microbiology, **50**, 1253-1258.

Diernhofer, K. (1932). Asculinbouillon als Hilfsmittel für die differenzierung von euter-und milchstreptokokken bei massenuntersuchungen. Milchwirtschaftliche Forschungen, **13**, 368-374.

Douglas, H.C. and Cruess, W.V. (1936). A lactobacillus from California wine: *Lactobacillus hilgardii*. Food Research, **1**, 113-119.

Edwards, C.G., Collins, M.D., Lawson, P.A. and Rodriguez, A.V. (2000). *Lactobacillus nagelii* sp. nov., an organism isolated from a partially fermented wine. International Journal of Systematic and Evolutionary Microbiology, **50**, 699-702.

Edwards, C.G., Haag, K.M., Collins, M.D., Hutson, R.A. and Huang, Y.C. (1998). *Lactobacillus kunkeei* sp. nov.: a spoilage microorganism associated with grape juice fermentations. Journal of Applied Microbiology, **84**, 698-702.

Eggerth, A.H. (1935). The gram-positive non-spore-bearing anaerobic bacilli of human feces. Journal of Bacteriology, **30**, 277-299.

Ehrmann, M.A., Brand, M., Stolz, P., Vogel, R.F. and Korakli, M. (2007). *Lactobacillus secaliphilus* sp. nov., isolated from type II sourdough fermentation. International Journal of Systematic and Evolutionary Microbiology, **57**, 745-750.

Ehrmann, M.A., Müller, M.R.A. and Vogel, R.F. (2003). Molecular analysis of sourdough reveals *Lactobacillus mindensis* sp. nov. International Journal of Systematic and Evolutionary Microbiology, **53**, 7-13.

Eldar, A., Bejerano, Y. and Bercovier, H. (1994): *Streptococcus shiloi* and *Streptococcus difficile*: two new streptococcal species causing a meningoencephalitis in fish. Current Microbiology, **28**, 139-143. Validation List N° 52. International Journal of Systematic Bacteriology, 1995, **45**, 197-198.

Eldar, A., Frelief, P.F., Assenta, L., Varner, P.W., Lawhon, S. and Bercovier, H. (1995). *Streptococcus shiloi*, the name for an agent causing septicemic infection in fish, is a junior synonym of *Streptococcus iniae*. International Journal of Systematic Bacteriology, **45**, 840-842.

Elliott, S.D. (1966). Streptococcal infection in young pigs. I. An immunochemical study of the causative agent (PM *Streptococcus*). Journal of Hygiene (Lond), **64**, 205-212.

Embley, T.M., Faquir, N., Bossart, W. and Collins, M.D. (1989). *Lactobacillus vaginalis* sp. nov. from the human vagina. International Journal of Systematic Bacteriology, **39**, 368-370.

Endo, A. and Okada, S. (2005). *Lactobacillus satsumensis* sp nov., made from fermented rice and other starchy materials. International Journal of Systematic and Evolutionary Microbiology, **55**, 83-85.

Endo, A. and Okada, S. (2006). *Oenococcus kitaharae*, of *Kitaharae* sp. nov., a non-acidophilic and non-malolactic-fermenting oenococcus isolated from a composting distilled *shochu* residue. International Journal of Systematic and Evolutionary Microbiology, **56**, 2345-2348.

Endo, A. and Okada, S. (2007). *Lactobacillus composti* sp. nov., a lactic acid bacterium isolated from a compost of distilled *shochu* residue. International Journal of Systematic and Evolutionary Microbiology, **57**, 870-872.

Endo, A. and Okada, S. (2007). *Lactobacillus farraginis* sp. nov. and *Lactobacillus parafarraginis* sp. nov., heterofermentative lactobacilli isolated from a compost of distilled *shochu* residue. International Journal of Systematic and Evolutionary Microbiology, **57**, 708-712.

Ennahar, S. and Cai, Y. (2005). Biochemical and genetic evidence for the transfer of *Enterococcus solitarius* Collins *et al.* 1989 to the genus *Tetragenococcus* as *Tetragenococcus solitarius* comb. nov. International Journal of Systematic and Evolutionary Microbiology, **55**, 589-592.

- Entani, E., Masai, H. and Suzuki, K.I.** (1986). *Lactobacillus acetotolerans*, a new species from fermented vinegar broth. *International Journal of Systematic Bacteriology*, **36**, 544-549.
- Falsen, E., Pascual, C., Sjöden, B., Ohlén, M. and Collins, M.D.** (1999). Phenotypic and phylogenetic characterization of a novel *Lactobacillus* species from human sources: description of *Lactobacillus iners* sp. nov. *International Journal of Systematic Bacteriology*, **49**, 217-221.
- Farrow, J.A.E. and Collins, M.D.** (1984). DNA base composition, DNA-DNA homology and long-chain fatty acid studies on *Streptococcus thermophilus* and *Streptococcus salivarius*. *Journal of General Microbiology*, **130**, 357-362.
- Farrow, J.A.E. and Collins, M.D.** (1985). *Enterococcus hirae*, a new species that includes amino acid assay strain NCDO 1258 and strains causing growth depression in young chickens. *International Journal of Systematic Bacteriology*, **35**, 73-75.
- Farrow, J.A.E. and Collins, M.D.** (1985). Taxonomic studies on streptococci of serological groups C, G and L and possibly related taxa. *Systematic and Applied Microbiology*, **5**, 483-493.
- Farrow, J.A.E. and Collins, M.D.** (1988). *Lactobacillus oris* sp. nov. from the human oral cavity. *International Journal of Systematic Bacteriology*, **38**, 116-118.
- Farrow, J.A.E., Facklam, R.R. and Collins, M.D.** (1989). Nucleic acid homologies of some vancomycin-resistant leuconostocs and description of *Leuconostoc citreum* sp. nov. and *Leuconostoc pseudomesenteroides* sp. nov. *International Journal of Systematic Bacteriology*, **39**, 279-283.
- Farrow, J.A.E., Kruze, J., Phillips, B.A., Bramley, A.J. and Collins, M.D.** (1984). Taxonomic studies on *Streptococcus bovis* and *Streptococcus equinus*: description of *Streptococcus alactolyticus* sp. nov. and *Streptococcus saccharolyticus* sp. nov. *Systematic and Applied Microbiology*, **5**, 467-482. Validation List N° 17. *International Journal of Systematic Bacteriology*, 1985, **35**, 223-225.
- Farrow, J.A.E., Phillips, B.A. and Collins, M.D.** (1988). Nucleic acid studies on some heterofermentative lactobacilli: description of *Lactobacillus malefermentans* sp. nov. and *Lactobacillus parabuchneri* sp. nov. *FEMS Microbiology Letters*, **55**, 163-168. Validation List N° 30. *International Journal of Systematic Bacteriology*, **39**, 371.
- Felis, G.E., Torriani, S. and Dellaglio, F.** (2005). Reclassification of *Pediococcus urinaeequi* (ex Mees, 1934) Garvie 1988 as *Aerococcus urinaeequi* comb. nov. *International Journal of Systematic and Evolutionary Microbiology*, **55**, 1325-1327.
- Felis, G.E. and Dellaglio, F.** (2007). Taxonomy of *Lactobacilli* and *Bifidobacteria*. *Current Issues in Intestinal Microbiology*, **8**, 44-61.
- Fernández, E., Blume, V., Garrido, P., Collins, M.D., Mateos, A., Domínguez, L. and Fernández-Garayzábal, J.F.** (2004). *Streptococcus equi* subsp. *ruminatorum* subsp. nov., isolated from mastitis in small ruminants. *International Journal of Systematic and Evolutionary Microbiology*, **54**, 2291-2296.
- Flint, S.H., Ward, L.J.H. and Brooks, J.D.** (1999). *Streptococcus waius* sp. nov., a thermophilic streptococcus from a biofilm. *International Journal of Systematic Bacteriology*, **49**, 759-767.
- Fortina, M.G., Ricci, G., Mora, D. and Manachini, P.L.** (2004). Molecular analysis of artisanal Italian cheeses reveals *Enterococcus italicus* sp. nov. *International Journal of Systematic and Evolutionary Microbiology*, **54**, 1717-1721.
- Franz, C.M.A.P., Vancanneyt, M., Vandemeulebroecke, K., De Wachter, M., Cleenwerck, I., Hoste, B., Schillinger, U., Holzappel, W.H. and Swings, J.** (2006). *Pediococcus stilesii* sp. nov., isolated from maize grains. *International Journal of Systematic and Evolutionary Microbiology*, **56**, 329-333.
- Franzmann, P.D. and Skerman, V.B.D.** (1981). *Agitococcus lubricus* gen. nov., sp. nov., a lipolytic, twitching coccus from freshwater. *International Journal of Systematic Bacteriology*, **31**, 177-183.
- Franzmann, P.D., Höpel, P., Weiss, N. and Tindall, B.J.** (1991). Psychrotrophic, lactic acid-producing bacteria from anoxic waters in Ace Lake, Antarctica: *Carnobacterium funditum* sp. nov. and *Carnobacterium alterfunditum* sp. Nov. *Arch. Microbiol.*, **156**, 255-262. Validation List N° 44. *International Journal of Systematic Bacteriology*, 1993, **43**, 188-189.
- Frost, W.D., and M.A. Engelbrecht.** (1940). *The streptococci*. Willdorf Book Co., Madison, Wis.

- Fujisawa, T., Adachi, S., Toba, T., Arimura, K. and Mitsuoka, T.** (1988). *Lactobacillus kefiranofaciens* sp. nov. isolated from kefir grains. *International Journal of Systematic Bacteriology*, **38**, 12-14.
- Fujisawa, T., Benno, Y., Yaeshima, T. and Mitsuoka, T.** (1992). Taxonomic study of the *Lactobacillus acidophilus* group, with recognition of *Lactobacillus gallinarum* sp. nov. and *Lactobacillus johnsonii* sp. nov., and synonymy of *Lactobacillus acidophilus* group A3 (Johnson *et al.* 1980) with the type strain of *Lactobacillus amylovorus* (Nakamura 1981). *International Journal of Systematic Bacteriology*, **42**, 487-491.
- Fujisawa, T., Itoh, K., Benno, Y. and Mitsuoka, T.** (1990). *Lactobacillus intestinalis* (ex Hemme 1974) sp. nov., nom. rev., isolated from the intestines of mice and rats. *International Journal of Systematic Bacteriology*, **40**, 302-304.
- Fujisawa, T., Shirasaka, S., Watabe, J. and Mitsuoka, T.** (1984). *Lactobacillus aviarius* sp. nov.: a new species isolated from the intestines of chickens. *Systematic and Applied Microbiology*, **5**, 414-420. Validation List N° 17. *International Journal of Systematic Bacteriology*, 1985, **35**, 223-225.
- Fujisawa, T., Shirasaka, S., Watabe, J. and Mitsuoka, T.** (1984). *Lactobacillus aviarius* sp. nov.: a new species isolated from the intestines of chickens. *Systematic and Applied Microbiology*, **5**, 414-420. Associate editor, IJSB: Validation list n° 20. *International Journal of Systematic Bacteriology*, 1986, **36**, 354-356.
- Garvie, E.I.** (1967). The growth factor and amino acid requirements of species of the genus *Leuconostoc* including *Leuconostoc paramesenteroides* sp. nov. and *Leuconostoc oenos*. *Journal of General Microbiology*, 1967, **48**, 439-447.
- Garvie, E.I.** (1983). *Leuconostoc mesenteroides* subsp. *cremoris* (Knudsen and Sørensen) comb. nov. and *Leuconostoc mesenteroides* subsp. *dextranicum* (Beijerinck) comb. nov. *International Journal of Systematic Bacteriology*, **33**, 118-119.
- Garvie, E.I.** (1986). Genus *Pediococcus* Claussen 1903, 68 AL. In: P. H. A. Sneath, N. S. Mair, M. E. Sharpe and J. G. Holt (Eds), *Bergey's Manual of Systematic Bacteriology*, first edition, **2**, The Williams and Wilkins Co., Baltimore. pp. 1075-1079. Validation List N° 25. *International Journal of Systematic Bacteriology*, 1988, **38**, 220-222.
- Garvie, E.I., Farrow, J.A.E. and Bramley, A.J.** (1983). *Streptococcus dysgalactiae* (Diernhofer) nom. Rev. *International Journal of Systematic Bacteriology*, **33**, 404-405.
- Gasser, F., Mandel, M. and Rogosa, M.** (1970). *Lactobacillus jensenii* sp. nov., a new representative of the subgenus *Thermobacterium*. *Journal of General and Applied Microbiology*, **62**, 219-222.
- Glazunova, O.O., Raoult, D. and Roux, V.** (2006). *Streptococcus massiliensis* sp. nov., isolated from a patient blood culture. *International Journal of Systematic and Evolutionary Microbiology*, **56**, 1127-1131.
- Gunther, H.L. and White, H.R.** (1961): The cultural and physiological characters of the pediococci. *Journal of General Microbiology*, **26**, 185-197.
- Handley, P., Coykendall, A., Beighton, D., Hardie, J.M. and Whiley, R.A.** (1991). *Streptococcus crista* sp. nov., a viridans streptococcus with tufted fibrils, isolated from the human oral cavity and throat. *International Journal of Systematic Bacteriology*, **41**, 543-547.
- Hansen, P.A.** (1968). Type strains of *Lactobacillus* species. American Type Culture Collection, Rockville, Md.
- Hansen, P.A. and Lessel, E.F.** (1971). *Lactobacillus casei* (Orla Jensen) comb. nov. *International Journal of Systematic Bacteriology*, **21**, 69-71.
- Hansen, P.A. and Mocquot, G.** (1970). *Lactobacillus acidophilus* (Moro) comb. nov. *International Journal of Systematic Bacteriology*, **20**, 325-327.
- Hemme, D., Raibaud, P., Ducluzeau, R., Galpin, J.V., Sicard, P. and Van Heijenoort, J.** (1980). *Lactobacillus murinus* n. sp., une nouvelle espèce de la flore dominante autochtone du tube digestif du rat et de la souris. *Annals of Microbiology*, **131A**, 297-308. Validation List N° 9. *International Journal of Systematic Bacteriology*, 1982, **32**, 384-385.
- Henneberg, W.** (1903). Zur Kenntnis der Milchsäurebakterien der Brennereimaische,
- Hiu, S.F., Holt, R.A., Sriranganathan, N., Seidler, R.J. and Freyer, J.L.** (1984). *Lactobacillus piscicola*, a new species from salmonid fish. *International Journal of Systematic Bacteriology*, **34**, 393-400.

- Holdeman, L.V. and Moore, W.E.C.** (1974). New genus, *Coprococcus*, twelve new species, and emended descriptions of four previously described species of bacteria from human feces. *International Journal of Systematic Bacteriology*, **24**, 260-277.
- Holley, R.A., Guan, T.Y., Peirson, M. and Yost, C.K.** (2002). *Carnobacterium viridans* sp. nov., an alkaliphilic, facultative anaerobe isolated from refrigerated, vacuum-packed bologna sausage. *International Journal of Systematic and Evolutionary Microbiology*, **52**, 1881-1885.
- Holzappel, W.H. and Gerber, E.S.** (1983). *Lactobacillus divergens* sp. nov., a new heterofermentative *Lactobacillus* species producing L(+)-lactate. *Systematic and Applied Microbiology*, **4**, 522-534.
- Holzappel, W.H. and Van Wyk, E.P.** (1982): *Lactobacillus kandleri* sp. nov., a new species of the subgenus *Betabacterium*, with glycine in the peptidoglycan. *Zentralbl. Bakteriol. Parasitenkd. Infektionskr. Hyg. Abt. 1 Orig.*, **C3**, 495-502. Validation List N° 10. *International Journal of Systematic Bacteriology*, 1983, **33**, 438-440.
- Holzappel, W.H., and O. Kandler.** (1969). Zur Taxonomie der Gattung *Lactobacillus* Beijerinck. VI: *Lactobacillus coprophilus* subsp. *confusus* nov. subsp., eine neue Unterart der Untergattung *Betabacterium* *Zentralbl. Bakteriol. Parasitenkd. Infektionskr. Hyg.* 123 657-666.
- Hoyles, L., Foster, G., Falsen, E., Thomson, L.F. and Collins, M.D.** (2001). *Facklamia miroungae* sp. nov., from a juvenile southern elephant seal (*Mirounga leonina*). *International Journal of Systematic and Evolutionary Microbiology*, **51**, 1401-1403.
- Hoyles, L., Lawson, P.A., Foster, G., Falsen, E., Ohlén, M., Grainger, J.M. and Collins, M.D.** (2000). *Vagococcus fessus* sp. nov., isolated from a seal and a harbour porpoise. *International Journal of Systematic and Evolutionary Microbiology*, **50**, 1151-1154.
- Ishikawa, M., Nakajima, K., Yanagi, M., Yamamoto, Y. and Yamasato, K.** (2003). *Marinilactibacillus psychrotolerans* gen. nov., sp. nov., a halophilic and alkaliphilic marine lactic acid bacterium isolated from marine organisms in temperate and subtropical areas of Japan. *International Journal of Systematic and Evolutionary Microbiology*, 2003, **53**, 711-720.
- Janssen, P.H., Evers, S., Rainey, F.A., Weiss, N., Ludwig, W., Harfoot, C.G. and Schink, B.** (1995). *Lactosphaera* gen. nov., a new genus of lactic acid bacteria, and transfer of *Ruminococcus pasteurii* Schink 1984 to *Lactosphaera pasteurii* comb. nov. *International Journal of Systematic Bacteriology*, **45**, 565-571.
- Jöborn, A., Dorsch, M., Olsson, J.C., Westerdahl, A. and Kjelleberg, S.** (1999). *Carnobacterium inhibens* sp. nov., isolated from the intestine of Atlantic salmon (*Salmo salar*). *International Journal of Systematic Bacteriology*, **49**, 1891-1898.
- Kandler, O. and Kunath, P.** (1983). *Lactobacillus kefir* sp. nov., a component of the microflora of kefir. *Systematic and Applied Microbiology*, **4**, 286-294.
- Kandler, O., Schillinger, U. and Weiss N.** (1983): *Lactobacillus halotolerans* sp. nov. nom. rev. and *Lactobacillus minor* sp. nov., nom. rev. *Systematic and Applied Microbiology*, **4**, 280-285.
- Kandler, O., Schillinger, U. and Weiss, N.** (1983). *Lactobacillus bifermentans* sp. nov., nom. rev., an organism forming CO₂ and H₂ from lactic acid. *Systematic and Applied Microbiology*, **4**, 408-412.
- Kandler, O., Stetter, K.O. and Köhl, R.** (1980). *Lactobacillus reuteri* sp. nov., a new species of heterofermentative lactobacilli. *Zentralbl. Mikrobiol. Parasitenkd. Infektionskr. Hyg. Abt. 1 Orig.*, 1980, C1 264-269. Validation list N° 8. *International Journal of Systematic Bacteriology*, 1982, **32**, 266-268.
- Kaneuchi, C., Seki, M. and Komagata, K.** (1988). Taxonomic study of *Lactobacillus mali* Carr and Davis 1970 and related strains: validation of *Lactobacillus mali* Carr and Davis 1970 over *Lactobacillus yamanashiensis* Nonomura 1983. *International Journal of Systematic Bacteriology*, **38**, 269-272.
- Katagiri, H., Kitahara, K. and Fukami, K.** (1934). The characteristics of the lactic acid bacteria isolated from moto, yeast mashes for sake manufacture. IV. Classification of the lactic acid bacteria. *Bulletin of the Agricultural Chemical Society of Japan*, **10**, 156-157.
- Kato, Y., Sakala, R.M., Hayashidani, H., Kiuchi, A., Kaneuchi, C., and Ogawa, M.** (2000). *Lactobacillus algidus* sp. nov., a psychrophilic lactic acid bacterium isolated from vacuum-packaged refrigerated beef. *International Journal of Systematic and Evolutionary Microbiology*, **50**, 1143-1149.

- Kawamura, Y., Hou, X.G., Sultana, F., Liu, S., Yamamoto, H. and Ezaki, T.** (1995). Transfer of *Streptococcus adjacens* and *Streptococcus defectivus* to *Abiotrophia* gen. nov. as *Abiotrophia adiacens* comb. nov. and *Abiotrophia defectiva* comb. nov., respectively. *International Journal of Systematic Bacteriology*, **45**, 798-803.
- Kawamura, Y., Hou, X.G., Todome, Y., Sultana, F., Hirose, K., Shu, S.E., Ezaki, T. and Ohkuni, H.** (1998). *Streptococcus peroris* sp. nov. and *Streptococcus infantis* sp. nov., new members of the *Streptococcus mitis* group, isolated from human clinical samples. *International Journal of Systematic Bacteriology*, **48**, 921-927.
- Kilian, M., Mikkelsen, L. and Henriksen, J.** (1989). Taxonomic study of viridans streptococci: description of *Streptococcus gordonii* sp. nov. and emended descriptions of *Streptococcus sanguis* (White and Niven, 1946), *Streptococcus oralis* (Bridge and Sneath, 1982), and *Streptococcus mitis* (Andrewes and Horder, 1906). *International Journal of Systematic Bacteriology*, **39**, 471-484.
- Kim, B., Lee, J., Jang, J. and Han, H.** (2003). *Leuconostoc inhae* sp. nov., a lactic acid bacterium isolated from *kimchi*. *International Journal of Systematic and Evolutionary Microbiology*, **53**, 1123-1126.
- Kim, J., Chun, J. and Han, H.U.** (2000). *Leuconostoc kimchii* sp. nov., a new species from *kimchi*. *International Journal of Systematic and Evolutionary Microbiology*, **50**, 1915-1919.
- Kitahara, K., Kaneko, T. and Goto, O.** (1957). Taxonomic studies on the hiochi-bacteria, specific saprophytes of sake. II. Identification and classification of hiochi-bacteria. *Journal of General and Applied Microbiology*, **3**, 111-120.
- Klein, E.** (1884). Micro-organisms and diseases. *Practitioner*, **32**, 321-352.
- Klein, G., Dicks, L.M.T., Pack, A., Hack, B., Zimmermann, K., Dellaglio, F. and Reuter, G.** (1996). Emended descriptions of *Lactobacillus sake* (Katagiri, Kitahara, and Fukami) and *Lactobacillus curvatus* (Abo-Elnaga and Kandler): numerical classification revealed by protein fingerprinting and identification based on biochemical patterns and DNA-DNA hybridizations. *International Journal of Systematic Bacteriology*, **46**, 367-376.
- Kleymans, U., Heinzl, H. and Hammes, W.P.** (1989). *Lactobacillus suebicus* sp. nov., an obligately heterofermentative *Lactobacillus* species isolated from fruit mashes. *Systematic and Applied Microbiology*, **11**, 267 - 271.
- Klipper-Bälz, R. and Schleiffer, K.H.** (1987). *Streptococcus suis* sp. nov., nom. rev. *International Journal of Systematic Bacteriology*, **37**, 160-162.
- Knight, R.G., Schlaes, D.M. and Messineo, L.** (1984). Deoxyribonucleic acid relatedness among major human enterococci. *International Journal of Systematic Bacteriology*, **34**, 327-331.
- Knudsen, S. and Sørensen, A.** (1929): Beiträge zur Bakteriologie der Säurewecker. *Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene. Abteilung II*, **79**, 75-85.
- Konstantinov, S.R., Poznanski, E., Fuentes, S., Akkermans, A.D.L., Smidt, H. and De Vos, W.M.** (2006). *Lactobacillus sobrius* sp. nov., abundant in the intestine of weaning piglets. *International Journal of Systematic and Evolutionary Microbiology*, **56**, 29-32.
- Koort, J., Coenye, T., Vandamme, P., Sukura, A. and Björkroth, J.** (2004). *Enterococcus hermannienseis* sp. nov., from modified-atmosphere-packaged broiler and canine tonsils. *International Journal of Systematic and Evolutionary Microbiology*, **54**, 1823-1827.
- Koort, J., Murros, A., Coenye, T., Eerola, S., Vandamme, P., Sukura, A. and Björkroth, J.** (2005). *Lactobacillus oligofermentans* sp. nov., associated with spoilage of modified-atmosphere-packaged poultry products. *Applied and Environmental Microbiology*, **71**, 4400-4406.
- Kozaki, M. and Okada, S.** (1983). *Lactobacillus vaccinostercus* sp. nov. In Validation of the Publication of New Names and New Combinations Previously Effectively Published Outside the IJSB, List no. 10. *International Journal of Systematic Bacteriology*, **33**, 438-440.
- Kröckel, L., Schillinger, U., Franz, C. M.A.P., Bantleon, A. and Ludwig, W.** (2003). *Lactobacillus versmoldensis* sp. nov., isolated from raw fermented sausage. *International Journal of Systematic and Evolutionary Microbiology*, **56**, 513 -517.
- Krooneman, J., Faber, F., Alderkamp, A.C., Oude Elferink, S.J.H.W., Driehuis, F., Cleenwerck, I., Swings, J., Gottschal, J.C. and Vancanneyt, M.** (2002). *Lactobacillus diolivorans* sp. nov., a 1,2-propanediol-degrading

bacterium isolated from aerobically stable maize silage. *International Journal of Systematic and Evolutionary Microbiology*, **52**, 639-646.

Krumholz, L.R. and Bryant, M.P. (1986) *Syntrophococcus sucromutans* sp. nov. gen. nov. uses carbohydrates as electron donors and formate, methoxybenzoids or *Methanobrevibacter* as electron acceptor systems. *Archives of Microbiology*, 1986, **143**, 313-318.

Kusuda, R., Kawai, K., Salati, F., Banner, C.R. and Fryer, J.L. (1991). *Enterococcus seriolicida* sp. nov., a fish pathogen. *International Journal of Systematic Bacteriology*, **41**, 406-409.

Latorre-Guzman, B.A., Kado, C.I., and Kunkee, R. (1977). *Lactobacillus hordniae*, a new species from the leafhopper (*Hordnia circellata*). *International Journal of Systematic Bacteriology*, **27**, 362-370.

Lauer, E. and Kandler, O. (1980). *Lactobacillus gasseri* sp. nov., a new species of the subgenus *Thermobacterium*. *Zentralbl. Bakteriol. Parasitenkd. Infektionskr. Hyg. Abt. 1. Orig. Reihe C*, **1**, 75-78.

Law-Brown, J. and Meyers, P.R. (2003). *Enterococcus phoeniculicola* sp. nov., a novel member of the enterococci isolated from the uropygial gland of the Red-billed Woodhoopoe, *Phoeniculus purpurens*. *International Journal of Systematic and Evolutionary Microbiology*, **53**, 683-685.

Lawson, P.A., Collins, M.D., Falsen, E., Sjöden, B. and Facklam, R.R. (1999). *Facklamia languida* sp. nov., isolated from human clinical specimens. *Journal of Clinical Microbiology*, **37**, 1161-1164.

Lawson, P.A., Falsen, E., Cotta, M.A. and Whitehead, T.R. (2007). *Vagococcus elongatus* sp. nov., isolated from a swine-manure storage pit. *International Journal of Systematic and Evolutionary Microbiology*, **57**, 751-754.

Lawson, P.A., Falsen, E., Ohlén, M. and Collins, M.D. (2001). *Aerococcus urinaehominis* sp. nov., isolated from human urine. *International Journal of Systematic and Evolutionary Microbiology*, **51**, 683-686.

Lawson, P.A., Falsen, E., Truberg-Jensen, K. and Collins, M.D. (2001). *Aerococcus sanguicola* sp. nov., isolated from a human clinical source. *International Journal of Systematic and Evolutionary Microbiology*, **51**, 475- 479.

Lawson, P.A., Foster, G., Falsen, E. and Collins, M.D. (2005). *Streptococcus marimammalium* sp. nov., isolated from seals. *International Journal of Systematic and Evolutionary Microbiology*, **55**, 271-274.

Lawson, P.A., Foster, G., Falsen, E., Davison, N. and Collins, M.D. (2004). *Streptococcus halichoeri* sp. nov., isolated from grey seals (*Halichoerus gryous*). *International Journal of Systematic and Evolutionary Microbiology*, **54**, 1753-1756.

Lawson, P.A., Foster, G., Falsen, E., Markopoulos, S.J. and Collins, M.D. (2005). *Streptococcus castoreus* sp. nov., isolated from a beaver (*Castor fiber*). *International Journal of Systematic and Evolutionary Microbiology*, **55**, 843-846.

Lawson, P.A., Foster, G., Falsen, E., Ohlén, M. and Collins, M.D. (1999). *Vagococcus lutrae* sp. nov., isolated from the common otter (*Lutra lutra*). *International Journal of Systematic Bacteriology*, **49**, 1251-1254.

Lawson, P.A., Foster, G., Falsen, E., Sjöden, B. and Collins, M.D. (1999). *Abiotrophia balaenopterae* sp. nov., isolated from the minke whale (*Balaenoptera acutorostrata*). *International Journal of Systematic Bacteriology*, **49**, 503-506.

Lawson, P.A., Foster, G., Falsen, E., Sjöden, B. and Collins, M.D. (1999). *Abiotrophia balaenopterae* sp. nov., isolated from the minke whale (*Balaenoptera acutorostrata*). *International Journal of Systematic Bacteriology*, **49**, 503-506.

Lawson, P.A., Wachter, C., Hansson, I., Falsen, E. and Collins, M.D. (2001). *Lactobacillus psittaci* sp. nov., isolated from a hyacinth macaw (*Anodorhynchus hyacinthinus*). *International Journal of Systematic and Evolutionary Microbiology*, **51**, 967-970.

Lee, J.S., Lee, K.C., Ahn, J.S., Mheen, T.I., Pyun, Y.R. and Park, Y.H. (2002). *Weissella koreensis* sp. nov., isolated from *kimchi*. *International Journal of Systematic and Evolutionary Microbiology*, **52**, 1257-1261.

Lee, M., Kim, M.K., Vnancanneyt, M., Swings, J., Kim, S.H., Kang, M.S. and Lee, S.T. (2005). *Tetragenococcus koreensis* sp. nov., a novel rhamnolipid-producing bacterium. *International Journal of Systematic and Evolutionary Microbiology*, **55**, 1409-1413.

- Lehmann, K.B. and Neumann, R.** (1896). Atlas und Grundriss der Bakteriologie und Lehrbuch der speziellen bakteriologischen Diagnostik, 1st Edition, J.F. Lehmann, München. In: Skerman, V. B. D., McGowan, V. and Sneath, P. H. A. (Eds), Approved Lists of Bacterial Names. International Journal of Systematic Bacteriology, 1980, **30**, 225-420.
- Leichmann, G.** (1896). Über die im Brennereiprozess bei der Bereitung der Kunsthefe auftretende spontane Milchsäuregärung. Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene. Abteilung II, **2**, 281-285.
- Leisner, J.J., Vancanneyt, M., Goris, J., Christensen, H. and Rusul, G.** (2000). Description of *Paralactobacillus selangolensis* gen. nov., sp.nov., a new lactic acid bacterium isolated from *chili bo*, a Malaysian food ingredient. International Journal of Systematic and Evolutionary Microbiology, **50**, 19-24.
- Li, Y., Raftis, E., Canchaya, C., Fitzgerald, G.F., Van Sinderen, D. and O'Toole, P.W.** (2006). Polyphasic analysis indicates that *Lactobacillus salivarius* subsp. *salivarius* and *Lactobacillus salivarius* subsp. *salicinius* do not merit separate subspecies status. International Journal of Systematic and Evolutionary Microbiology, **56**, 2397-2403.
- Lindner, P.** (1887). Über ein neues in Malzmaischen vorkommendes, milchsäurebildendes Ferment. Wochenschrift für Brauerei, **4**, 437-440. Approved Lists of Bacterial Names. International Journal. of. Systematic Bacteriology, 1980, **30**, 225-420.
- Lister, J.** (1873). A further contribution to the natural history of bacteria and the germ theory of fermentative changes. Quart. Microbiol. Sci., **13**, 380-408.
- Liu, B. and Dong, X.** (2002). *Lactobacillus pantheris* sp. nov., isolated from faeces of a jaguar. International Journal of Systematic and Evolutionary Microbiology, **52**, 1745 – 1748.
- Liu, J.R., Tanner, R.S., Schumann, P., Weiss, N., McKenzie, C.A., Janssen, P.H., Seviour, E.M., Lawson, P. A., Allen, T.D. and Seviour, R.J.** (2002). Emended description of the genus *Trichococcus*, description of *Trichococcus collinsii* sp. nov., and reclassification of *Lactosphaera pasteurii* as *Trichococcus pasteurii* comb. nov. and of *Ruminococcus palustris* as *Trichococcus palustris* comb. nov. in the low-G + C Gram-positive bacteria. International Journal of Systematic and Evolutionary Microbiology, **52**, 1113-1126.
- Liu, L., Zhang, B., Tong, H. and Dong, X.** (2006) *Pediococcus ethanolidurans* sp. nov., isolated from the walls of a distilled-spirit-fermenting cellar. Interantional Journal of. Systematic and. Evolutionary Microbiology, **56**, 2405-2408.
- Magnusson, J., Jonsson, H., Schnürer, J. and Roos, S.** (2002). *Weissella soli* sp. nov., a lactic acid bacterium isolated from soli. International Journal of Systematic and Evolutionary Microbiology, **52**, 831-834.
- Martinez-Murcia, A.J. and Collins, M.D.** (1991). A phylogenetic analysis of an atypical leuconostoc: description of *Leuconostoc fallax* sp. nov. FEMS Microbiology Letters, **82**, 55-60. Validation List N° 40. International Journal of Systematic Bacteriology, 1992, **42**, 191- 192.
- Martinez-Murcia, A.J. and Collins, M.D.** (1991). *Enterococcus sulfurous*, a new yellow-pigmented *Enterococcus* species. FEMS Microbiology Letters, **80**, 69-74.
- Mees, R.H.** (1934): Onderzoekingen over de Biersarcina. Thesis. Technical University Delft, Holland. pp. 1-110.
- Meroth, C.B., Hammes, W.P. and Hertel, C.** (2004). Characterization of the microbiota of rice sourdoughs and description of *Lactobacillus spicheri* sp. nov. Systematic and Applied Microbiology, **27**, 151-159.
- Miller, A., III, Morgan, M.E. and Libbey, L.M.** (1974). *Lactobacillus maltaromicus*, a new species producing a malty aroma. International Journal of Systematic Bacteriology, **24**, 346-354.
- Mitsuoka, T. and Fujisawa, T.** (1987). *Lactobacillus hamsteri*, a new species from the intestine of hamsters. Proc. Jpn. Acad. Ser. B Phys. Biol. Sci. Validation List N° 25. International Journal of Systematic Bacteriology, **38**, 220-222. Validation List N° 25. International Journal of Systematic Bacteriology, 1988, **38**, 220-222.
- Miyamoto, M., Seto, Y., Hao, D.H., Teshima, T., Sun, Y.B., Kabuki, T., Yao, L.B. and Nakajima, H.** (2006). *Lactobacillus harbinensis* sp. nov., consisted of strains isolated from traditional fermented vegetables ‘Suan cai’ in Harbin, Northeastern China and *Lactobacillus perolens* DSM 12745. Systematic and Applied Microbiology, **28**, 688-694.
- Moore, W.E.C. and Holdeman, L.V.** (1970). *Propionibacterium*, *Arachnia*, *Actinomyces*, *Lactobacillus* and *Bifidobacterium*. In: Outline of Clinical Methods in Anaerobic Bacteriology. Cato *et al.* (Eds.). Virginia Polytechnic Institute, Anaerobe Laboratory, Blacksburg, Virginia, pp. 15-22.

- Mora, D., Scarpellini, M., Franzetti, L., Colombo, S. and Galli, A.** (2003). Reclassification of *Lactobacillus maltaromaticum* comb. Nov. International Journal of Systematic and Evolutionary Microbiology, **53**, 675-768.
- Morlon-Guyot, J., Guyot, J.P., Pot, B., Jacobe de Haut, I. and Rimbault, M.** (1998). *Lactobacillus manihotivorans* sp. nov., a new starch-hydrolysing lactic acid bacterium isolated during cassava sour starch fermentation. International Journal of Systematic Bacteriology, **48**, 1101-1109.
- Moro E.** (1900). Über den *Bacillus acidophilus* n. spec. Ein Beitrag zur Kenntnis der normalen Darmbakterien des Säuglings (*Bacillus acidophilus* n. spec. A contribution to the knowledge of the normal intestinal bacteria of infants.) Jahrbuch für Kinderheilkunde, **52**, 38–55.
- Morotomi, M., Yuki, N., Kado, Y., Kushiro, A., Shimazaki, T., Watanabe, K. and Yuyama, T.** (2002). *Lactobacillus equi* sp. nov., a predominant intestinal *Lactobacillus* species of the horse isolated from faeces of healthy horses. International Journal of Systematic and Evolutionary Microbiology, **52**, 211-214.
- Mukai, T., Arihara, K., Ikeda, A., Nomura, K., Suzuki, F. and Ohiri, H.** (2003). *Lactobacillus kitasotonis* sp. nov., from chicken intestine. International Journal of Systematic and Evolutionary Microbiology, **53**, 2055-2059.
- Müller, M.R.A., Ehrmann, M.A. and Vogel, R.F.** (2000). *Lactobacillus frumenti* sp. nov., a new species of lactic acid bacterium isolated from rye-bran fermentations with a long fermentation period. International Journal of Systematic and Evolutionary Microbiology, **50**, 2127-2133.
- Nakajima, K., Hirota, K., Nodasaka, Y. and Yumoto, I.** (2005). *Alkalibacterium iburiense* sp. nov., an obligate alkaliphile that reduces an indigo dye. International Journal of Systematic and Evolutionary Microbiology, **55**, 1525-1530.
- Nakamura, L.K.** (1981). *Lactobacillus amylovorus*, a new starch-hydrolyzing species from cattle waste-corn fermentations. International Journal of Systematic Bacteriology, **31**, 56-63.
- Nakamura, L.K. and Crowell, C.D.** (1979). *Lactobacillus amylophilus*, a new starch-hydrolyzing species from swine waste-corn fermentation. Dev. Ind. Microbiol., **20**, 532-540. Validation List N° 6. International Journal of Systematic Bacteriology, 1981, **31**, 215-218.
- Naser, S.M., Vancanneyt, M., De Graef, E., Devriese, L.A., Snauwaert, C., Lefebvre, K., Hoste, B., Svec, P., Decostere, A., Haesebrouck, F. and Swings, J.** (2005). *Enterococcus canintestini* sp. nov., from faecal samples of healthy dogs. International Journal of Systematic and Evolutionary Microbiology, **55**, 2177-2182.
- Naser, S.M., Vancanneyt, M., Snauwaert, C., Vrancken, G., Hoste, B., De Vuyst, L. and Swings, J.** (2006). Reclassification of *Lactobacillus amylophilus* LMG 11400 and NRRL B-4435 as *Lactobacillus amylotrophicus* sp. nov. International Journal of Systematic and Evolutionary Microbiology., **56**, 2523-2527.
- Nikolaitchouk, N., Wachter, C., Falsen, E., Andersch, B., Collins, M.D. and Lawson, P.A.** (2001). *Lactobacillus coleohominis* sp. nov., isolated from human sources. International Journal of Systematic and Evolutionary Microbiology, **51**, 2081-2085.
- Niven Jr., C.F. and Evans, J.B.** (1957). *Lactobacillus viridescens* nov. spec., a heterofermentative species that produces a green discoloration of cured meat pigments. *Journal of Bacteriology* **73**, pp. 758–759.
- Nowlan, S.S. and Deibel, R.H.** (1967). Group Q Streptococci II. Nutritional characteristics and growth relationship to thymine, folate, and folinate. *Journal of Bacteriology*; **94** (2), 297–299.
- Ntougias, S. and Russell, N.J.** (2001). *Alkalibacterium olivoapovliticus* gen. nov., sp. nov., a new obligately alkaliphilic bacterium isolated from edible-olive wash-waters. International Journal of Systematic and Evolutionary Microbiology, **51**, 1161-1170.
- Okada, S., Suzuki, Y. and Kozaki, M.** (1979). A new heterofermentative *Lactobacillus* species with meso-diaminopimelic acid in peptidoglycan, *Lactobacillus vaccinostrercus* Kozaki and Okada sp. nov. *J. Gen. Appl. Microbiol.*, **25**, 215-221. Validation List N° 10. International Journal Systematic Bacteriology, 1983, **33**, 438-440.
- Orla-Jensen, A.D. and Hansen, P.A.** (1932). The bacteriological flora of spontaneously soured milk and of commercial starters for butter making. *Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene. Abteilung II*, **86**, 6-29.
- Orla-Jensen, S.** (1916). *Mælkeri-Bakteriologi*. Schøberske Forlag, Copenhagen.

Orla-Jensen, S. (Ed.). (1919). The lactic acid bacteria. Høst and Son, Copenhagen. pp. 1-196.

Osawa, R., Fujisawa, T. and Pukall, R. (2006). *Lactobacillus apodemi* sp. nov., a tannase-producing species isolated from wild mouse faeces. International Journal of Systematic and Evolutionary Microbiology, **56**, 1693-1696.

Osawa, R., Fujisawa, T. and Sly, L.L. (1995). *Streptococcus gallolyticus* sp. nov; gallate degrading organisms formerly assigned to *Streptococcus bovis*. Systematic and Applied Microbiology, **18**, 74-78. Validation List N° 56, International Journal of Systematic Bacteriology, 1996, **46**, 362-363.

Pedersen, C. and Roos, S. (2004). *Lactobacillus saemneri* sp. nov., isolated from pig faeces. International Journal of Systematic and Evolutionary Microbiology, **54**, 1365-1368.

Pier, G.B. and Madin, S.H. (1976). *Streptococcus iniae* sp. nov., a beta hemolytic streptococcus isolated from an Amazon freshwater dolphin, *Inia geoffrensis*. International J

Prévot, A.R. (1925). Les streptocoques anaérobies. Annales de l'Institut Pasteur (Paris), **39**, 417-447.

Pikuta, E.V., Hoover, R.B., Bej, A.K., Marsic, D., Whitman, W.B., Krader, P.E. and Tang, J. (2006). *Trichococcus patagoniensis* sp. nov., a facultative anaerobe that grows at -5 °C, isolated from penguin guano in Chilean Patagonia. International Journal of Systematic and Evolutionary Microbiology, **56**, 2055-2062.

Pikuta, E.V., Marsic, D., Bej, A., Tang, J., Krader, P. and Hoover, R.B. (2005). *Carnobacterium pleistocenium* sp. nov., a novel psychotolerant, facultative anaerobe isolated from permafrost of the Fox Tunnel in Alaska. International Journal of Systematic and Evolutionary Microbiology, **55**, 473-478.

Pompei, R., Berlutti, F., Thaller, M.C., Ingianni, A., Cortis, G. and Dainelli, B. (1992). *Enterococcus flavescens* sp. nov., a new species of enterococci of clinical origin. International Journal of Systematic Bacteriology, **42**, 365-369.

Poyart, C., Quesne, G. and Trieu, Cuot, P. (2002). Taxonomic dissection of *Streptococcus bovis* group by analysis of manganese-dependent superoxide dismutase gene (*sodA*) sequences: reclassification of '*Streptococcus infantarius* subsp. coli' as *Streptococcus lutetiensis* sp. nov. and *Streptococcus bovis* biotype 11.2 as *Streptococcus pasteurianus* sp. nov. International Journal of Systematic and Evolutionary Microbiology, **52**, 1247-1255.

Prévot A.R. (1933). Etudes de systematique bactérienne. I. Lois generales. II. Cocci anaerobies. Ann Soc Nat Zool Biol Anim., **15**, 23-26.

Prevot, A.R. (1924). Comptes Rendus des Seances de la Societe de Biologie (Paris), **91**, 426-428; Holdeman, L. V. and W. E. C. Moore. 1974. International Journal of Systematic Bacteriology **24**, 260-277.

Reuter, G. (1970). Laktobazillen und eng verwandte Mikroorganismen in Fleisch und Fleischwaren. 2. Mitteilung: Die Charakterisierung der isolierten Laktobazillenstämme Fleischwirtschaft, **50**, 954-962.

Reuter, G. (1983). *Lactobacillus alimentarius* sp. nov., nom. rev. and *Lactobacillus farciminis* sp. nov. , nom. rev. Systematic and Applied Microbiology, **4**, 277-279.

Robinson, I.M., Stromley, J.M., Varel, V.H. and Cato, E.P. (1988). *Streptococcus intestinalis*, a new species from the colons of pigs. International Journal of Systematic Bacteriology, **38**, 245-248.

Rodas, A.M., Chenoll, E., Macián, M.C., Ferrer, S., Pardo, I. and Aznar, R. (2006). *Lactobacillus vini* sp. nov., a wine lactic acid bacterium homofermentative for pentoses. International Journal of Systematic and Evolutionary Microbiology, **56**, 513-517.

Rodrigues, U. and Collins, M.D. (1990). Phylogenetic analysis of *Streptococcus saccharolyticus* based on 16S rRNA sequencing. FEMS Microbiology Letters, **71**, 231-234. Validation List N° 36, International Journal of Systematic Bacteriology, 1991, **41**, 178-179.

Roggenkamp, A., Abele-Horn, M., Trebesius, K.H., Tretter, U., Autenrieth, I. B. and Heesemann, J. (1998). *Abiotrophia elegans* sp. nov., a possible pathogen in patients with culture-negative endocarditis. Journal of Clinical Microbiology, **36** (1), 100-104. Validation List N° 68. International Journal of. Systematic. Bacteriology, 1999, **49**, 1-3.

Rogosa, M., Wiseman, R.F., Mitchell, J.A. and Disraely, M.N. (1953). Species differentiation of oral lactobacilli from man including descriptions of *Lactobacillus salivarius* nov. spec. and *Lactobacillus cellobiosus* nov. spec. Journal of Bacteriology, **65**, 681-699.

- Roos, S., Engstrand, L. and Jonsson, H.** (2005). *Lactobacillus gastricus* sp. nov., *Lactobacillus antri* sp. nov., *Lactobacillus kalixensis* sp. nov., and *Lactobacillus ultunensis* sp. nov., isolated from human stomach mucosa. *International Journal of Systematic and Evolutionary Microbiology*, **55**, 77-82.
- Roos, S., Karner, F., Axelsson, L. and Jonsson, H.** (2000). *Lactobacillus mucosae* sp. nov., a new species with in vitro mucus-binding activity isolated from pig intestine. *International Journal of Systematic and Evolutionary Microbiology*, **50**, 251-258.
- Rosenbach, F.J.** (1884). *Microorganismen bei den wund-infections-krankheiten des menschen*. J. F. Bergmann, Wiesbaden, pp. 1-122.
- Rurangirwa, F.R., Teitzel, C.A., Cui, J., French, D.M., McDonough, P.L. and Besser, T.** (2000). *Streptococcus didelphis* sp. nov., a streptococcus with marked catalase activity isolated from opossums (*Didelphis virginiana*) with suppurative dermatitis and liver fibrosis. *International Journal of Systematic and Evolutionary Microbiology*, **50**, 759-765.
- Sakala, R.M., Kato, Y., Hayashidani, H., Murakami, M., Kaneuschi, C. and Ogawa, M.** (2002). *Lactobacillus fuchuensis* sp. nov., isolated from vacuum-packaged refrigerated beef. *International Journal of Systematic and Evolutionary Microbiology*, **52**, 1151-1154.
- Sand, G. and Jensen, C.O.** (1888). Die aetilogie der druise. *Deutsche Zeitschrift fur Tiermedizin und Vergleichende Pathologie*, **13**, 437-464.
- Satomi, M., Kimura, B., Mizoi, M., Sato, T. and Fujii, T.** (1997). *Tetragenococcus muriaticus* sp. Nov., a new moderately halophilic lactic acid bacterium isolated from fermented squid liver sauce. *International Journal of Systematic Bacteriology*, **47**, 832-836.
- Scheff, G., Salcher, O. and Lingens, F.** (1984). *Trichococcus flocculiformis* gen. nov. sp. nov. A new gram-positive filamentous bacterium isolated from bulking sludge. *Applied Microbiology Biotechnology*, **19**, 114-119.
- Scheirlinck, I., Van Der Meulen, R., Van Schoor, A., Cleenwerck, I., Huys, G., Vandamme, P., De Vuyst, L. and Vancanneyt, M.** (2007). *Lactobacillus namurensis* sp. nov., isolated from a traditional Belgian sourdough. *International Journal of Systematic and Evolutionary Microbiology*, **57**, 223-227.
- Schillinger, U., Holzappel, W. and Kandler, O.** (1989). Nucleic acid hybridization studies on *Leuconostoc* and heterofermentative lactobacilli and description of *Leuconostoc amelibiosum* sp. nov. *Systematic and Applied Microbiology*, **12**, 48-55.
- Schink, B.** (1985). *Ruminococcus pasteurii* sp. nov. In: Validation of the Publication of New Names and New Combinations Previously Effectively Published Outside the IJSB, List no. 18. *International Journal of Systematic Bacteriology*, **35**, 375-376.
- Schlegel, L., Grimont, F., Ageron, E., Grimont, P.A.D. and Bouvet, A.** (2003). Reappraisal of the taxonomy of the *Streptococcus bovis*/*Streptococcus equinus* complex and related species: description of *Streptococcus gallolyticus* subsp. *gallolyticus* subsp. nov., *S. gallolyticus* subsp. *macedonicus* subsp. nov. and *S. gallolyticus* subsp. *pasteurianus* subsp. nov. *International Journal of Systematic and Evolutionary Microbiology*, **53**, 631-645.
- Schlegel, L., Grimont, F., Collins, M.D., Régnault, B., Grimont, P.A.D. and Bouvet, A.** (2000). *Streptococcus infantarius* sp. nov., *Streptococcus infantarius* subsp. *infantarius* subsp. nov. and *Streptococcus infantarius* subsp. *coli* subsp. nov., isolated from humans and food. *International Journal of Systematic and Evolutionary Microbiology*, **50**, 1425-1434.
- Schleifer, K.H. and Kilpper-Bälz, R.** (1984). Transfer of *Streptococcus faecalis* and *Streptococcus faecium* to the genus enterococcus nom. rev. as *Enterococcus faecalis* comb. nov. and *Enterococcus faecium* comb. nov. *International Journal of Systematic Bacteriology*, **34**, 31-34.
- Schleifer, K.H., Ehrmann, M., Krusch, U. and Neve, H.** (1991). Revival of the species *Streptococcus thermophilus* (ex Orla-Jensen, 1919) nom. rev. *Systematic Applied Microbiology*, 1991, **14**, 386-388. Validation List N° 54. *International Journal Systematic Bacteriology*, 1995, **45**, 619-620.
- Schleifer, K.H., Kraus, J., Dvorak, C., Kilpper-Bälz, R., Collins, M.D. and Fischer, W.** (1985). Transfer of *Streptococcus lactis* and related streptococci to the genus *Lactococcus* gen. nov. *Systematic and Applied Microbiology*, **6**, 183-195. Validation List N° 20. *International Journal of Systematic Bacteriology*, 1986, **36**, 354-356.

- Sharpe, M.E., Latham, M.J., Garvie, E.I., Zirngibl, J. and Kandler, O.** (1973). Two new species of *Lactobacillus* isolated from the bovine rumen, *Lactobacillus ruminis* sp. nov. and *Lactobacillus vitulinus* sp. nov. *Journal of General Microbiology*, **77**, 37-39.
- Shaw, B.G. and Harding, C.D.** (1989). *Leuconostoc gelidum* sp. nov. and *Leuconostoc carnosum* sp. nov. from chill-stored meats. *International Journal of Systematic Bacteriology*, **39**, 217-223.
- Sherman, J. M. and Wing., H.U.** (1937). *Streptococcus durans* n. sp. *Journal of Dairy Science*, **20** (3), 165-167.
- Shewmaker, P.L., Steigerwalt, A.G., Morey, R.E., Carvalho, M. da G.S., Elliot, J.A., Joyce, K., Barratt, T.J., Teixeira, L.M. and Facklam, R.R.** (2004). *Vagococcus carniphilus* sp. nov., isolated from ground beef. *International Journal of Systematic and Evolutionary Microbiology*, **54**, 1505-1510.
- Skaar, I., Gaustad, P., Tønjum, T., Holm, B. and Stenwig, H.** (1994). *Streptococcus phocae* sp. nov., a new species isolated from clinical specimens from seals. *International Journal of Systematic Bacteriology*, **44**, 646-650.
- Sleat, R., Mah, R.A. and Robinson, R.** (1985). *Acetoanaerobium noterae* gen. nov., sp. nov.: an anaerobic bacterium that forms acetate from H₂ and CO₂. *International Journal of Systematic Bacteriology*, **35**, 10-15.
- Smith, F.R. and Sherman, J.M.** (1939). The hemolytic streptococci of human feces. *Journal of Infectious Diseases*, **62**, 186-189.
- Stackebrandt, E., Schumann, P., Swiderski, J. and Weiss, N.** (1999). Reclassification of *Brevibacterium incertum* (Breed 1953) as *Desemzia incerta* gen. nov., comb. nov. *International Journal of Systematic Bacteriology*, **49**, 185-188.
- Steinhaus, E.** (1941). A study of the bacteria associated with thirty species of insects. *Journal of Bacteriology*, **42**, 757-790.
- Stetter, H. and Stetter, K.O.** (1980). *Lactobacillus bavaricus* sp. nov., a new species of the subgenus Streptobacterium. *Zentralbl. Bakteriolog. Parasitenkd. Infektionskr. Hyg. Abt. 1 Orig. Reihe C*, **1**, 70-74.
- Suzuki, K., Funahashi, W., Koyanagi, M. and Yamashita, H.** (2004). *Lactobacillus paracollinoides* sp. nov., isolated from brewery environments. *International Journal of Systematic and Evolutionary Microbiology*, **54**, 115-117.
- Svec, P., Devriese, L.A., Sedláček, I., Baele, M., Vancanneyt, M., Haesebrouck, F., Swings, J. and Doskar, J.** (2001). *Enterococcus haemoperoxidus* sp. nov. and *Enterococcus moraviensis* sp. nov., isolated from water. *International Journal of Systematic and Evolutionary Microbiology*, **51**, 1567-1574.
- Svec, P., Vancanneyt, M., Devriese, L.A., Naser, S.M., Snnauwaert, C., Lefebvre, K., Hoste, B. and Swings, J.** (2005). *Enterococcus aquimarinus* sp. nov., isolated from sea water. *International Journal of Systematic and Evolutionary Microbiology*, **55**, 2183-2187.
- Svec, P., Vancanneyt, M., Sedláček, I., Naser, S.M., Snnauwaert, C., Lefebvre, K., Hoste, B. and Swings, J.** (2006). *Enterococcus silesiacus* sp. nov. and *Enterococcus termitis* sp. nov. *International Journal of Systematic and Evolutionary Microbiology*, **56**, 577-581.
- Svec, P., Vncanneyt, M., Koort, J., Naser, S.M., Hoste, B., Vihavainen, E., Vandamme, P., Swings, J. and Björkroth, J.** (2005). *Enterococcus devriesei* sp. nov., associated with animal sources. *International Journal of Systematic and Evolutionary Microbiology*, **55**, 2479-2484.
- Swezey, J.L., Nakamura, L.K., Abbott, T.P. and Peterson, R.E.** (2000). *Lactobacillus arizonensis* sp. nov., isolated from jojoba meal. *International Journal of Systematic and Evolutionary Microbiology*, **50**, 1803-1809.
- Takizawa, S., Kojima, S., Tamura, S., Fujinaga, S., Benno, Y. and Nakase, T.** (1994). *Lactobacillus kefirgranum* sp. nov. and *Lactobacillus parakefir* sp. nov., two new species from kefir grains. *International Journal of Systematic Bacteriology*, **44**, 435-439.
- Tanasupawat, S., Shida, O., Okada, S. and Komagata, K.** (2000). *Lactobacillus acidipiscis* sp. nov., and *Weissella thailandensis* sp. nov., isolated from fermented fish in Thailand. *International Journal of Systematic and Evolutionary Microbiology*, **50**, 1479-1485.
- Teixeira, L.M., Carvalho, M. da G.G., Espinola, M.M.B., Steigerwalt, A.G., Douglas, M.P., Brenner, D.J. and Facklam, R.R.** (2001). *Enterococcus porcinus* sp. nov. and *Enterococcus ratti* sp. nov., associated with enteric disorders in animals. *International Journal of Systematic and Evolutionary Microbiology*, **51**, 1737-1743.

- Tong, H. and Dong, X.** (2005). *Lactobacillus concavus* sp. nov., isolated from the walls of a distilled spirit fermenting cellar in China. *International Journal of Systematic and Evolutionary Microbiology*, **55**, 2199-2202.
- Tong, H., Gao, X. and Dong, X.** (2003). *Streptococcus oligofermentans* sp. nov., a noval oral isolate from caries-free humans. *International Journal of Systematic and Evolutionary Microbiology*, **53**, 1101-1104.
- Torriani, S., Van Reenen, C.A., Klein, G., Reuter, G., Dellaglio, F. And Dicks, L.M.T.** (1996). *Lactobacillus curvatus* subsp. nov. and *Lactobacillus curvatus* subsp. melibiosus subsp. nov. and *Lactobacillus sake* subsp. *carosus* subsp. nov., new subspecies of *Lactobacillus curvatus* Abo-Elnaga and Kandler 1965 and *Lactobacillus sake* Katagiri, Kitahara and Fukami 1934 (Klein *et al.* 1996, emended descriptions), respectively. *International Journal of Systematic and Bacteriology*, **46**, 1158-1163.
- Troili-Petersson, G.** (1903). Studien über die Mikroorganismen des schwedischen güterkäses. *Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene. Abteilung II*, **11**, 120-143.
- Tsakalidou, E., Zoidou, E., Pot, B., Wassill, L., Ludwig, W., Devriese, L.A., Klantzopoulos, G., Schleiffer, K. H. and Kersters, K.** (1998). Identification of streptococci from Greek Kasser cheese and description of *Streptococcus macedonicus* sp. nov. *International Journal of Systematic Bacteriology*, **48**, 519-527.
- Tyrell, G.J., Turnbull, L., Teixeira, L.M., Lefebvre, J., Carvalho, M.G.S., Facklam, R.R. and Lovgren, M.** (2002). *Enterococcus gilvus* sp. nov. and *Enterococcus pallens* sp. nov. isolated from human clinical samples. *Journal of Clinical Microbiology*, **40**, 1140-1145.
- Tyrrell, G.J., Turnbull, L., Teixeira, L.M., Lefebvre, J., Carvalho, M.G.S., Facklam, R.R. and Lovgren, M.** (2002). *Enterococcus gilvus* sp. nov. and *Enterococcus pallens* sp. nov. isolated from human clinical specimens. *Journal of Clinical Microbiology*, **40**, 1140-1145.
- Valcheva, R., Ferchichi, M.F., Korakli, M., Ivanova, I., Gänzle, M.G., Vogel, R.F., Prévost, H., Onno, B. and Dousset, X.** (2006). *Lactobacillus nantensis* sp. nov., isolated from French wheat sourdough. *International Journal of Systematic and Evolutionary Microbiology*, **56**, 587-591.
- Valcheva, R., Korakli, M., Onno, B., Prévost, H., Ivanova, I., Ehrmann, M.A., Dousset, X., Gänzle, M. G. and Vogel, R.F.** (2005). *Lactobacillus hammesii* sp. nov., isolated from French sourdough. *International Journal of Systematic and Evolutionary Microbiology*, **55**, 763-767.
- Vancanneyt, M., Devriese, L.A., De Graef, E.M., Baele, M., Lefebvre, K., Snauwaert, C., Vandamme, P., Swings, J. and Haesebrouck, F.** (2004). *Streptococcus minor* sp. nov., from faecal samples and tonsils of domestic animals. *International Journal of Systematic and Evolutionary Microbiology*, **54**, 449-452.
- Vancanneyt, M., Mengaud, J., Cleenwerck, I., Vanhocker, K., Hoste, B., Dawyndt, P., Degivry, M. C., Ringuet, D., Janssens, D. and Swings, J.** (2004). Reclassification of *Lactobacillus kefiranofaciens* subsp. *kefirgranum* subsp. nov. and emended description of *L. kefiranofaciens* Fujisawa *et al.* 1988. *International Journal of Systematic and Evolutionary Microbiology*, **5**, 551-556.
- Vancanneyt, M., Naser, S.M., Engelbeen, K., De Wachter, M., Van Der Meulen, R., Cleenwerck, I., Hoste, B., De Vuyst, L. and Swings, J.** (2006). Reclassification of *Lactobacillus brevis* strains LMG 11494 and LMG 11984 as *Lactobacillus parabrevis* sp. nov. *International Journal of Systematic and Evolutionary Microbiology*, **56**, 1553-1557.
- Vancanneyt, M., Neysens, P., De Wachter, M., Engelbeen, K., Snauwaert, C., De Vuyst, L. and Swings, J.** (2005). *Lactobacillus acidifarinae* sp. nov. and *Lactobacillus zymae* sp. nov., from wheat sourdoughs. *International Journal of Systematic and Evolutionary Microbiology*, **55**, 615-620.
- Vancanneyt, M., Snauwaert, C., Cleenwerck, I., Baele, M., Descheemaeker, P., Goossens, H., Pot, B., Vandamme, P., Swings, J., Haesebrouck, F. and Devriese, L.A.** (2001). *Enterococcus villorum* sp. nov., an enteroadherent bacterium associated with diarrhea in piglets. *International Journal of Systematic and Evolutionary Microbiology*, **51**, 393-400.
- Vancanneyt, M., Zamfir, M., Devriese, L. A., Lefebvre, K., Engelbeen, K., Vandemeulebroecke, K., Amar, M., De Vuyst, L., Haesebrouck, F. and Swings, J.** (2004). *Enterococcus saccharominimus* sp. nov., from dairy products. *International Journal of Systematic and Evolutionary Microbiology*, **54**, 2175-2179.
- Vandamme, P., Homme, J., Snauwaert, C., Hoste, B., Cleenwerck, I., Lefebvre, K., Vancanneyt, M., Swings, J., Devriese, L.A. and Haesebrouck, F.** (2001). *Globicatella sulfidifaciens* sp. nov., isolated from purulent infections in domestic animals. *International Journal of Systematic and Evolutionary Microbiology*, **51**, 1745-1749.

- Vandamme, P., Pot, B., Falsen, E., Kersters, K. and Devriese, L.A.** (1996). Taxonomic study of Lancefield streptococcal groups C, G, and L (*Streptococcus dysgalactiae*) and proposal of *S. dysgalactiae* subsp. *equisimilis* subsp. nov. International Journal of Systematic Bacteriology, **46**, 774-781.
- Vela, A.I., Fernández, E., Lawson, P.A., Latre, M.V., Falsen, E., Domínguez, L., Collins, M.D. and Fernández-Garayzábal, J.F.** (2002). *Streptococcus entericus* sp. nov., isolated from cattle intestine. International Journal of Systematic and Evolutionary Microbiology, **52**, 665-669.
- Vogel, R.F., Bocker, G., Stolz, P., Ehrmann, M., Fanta, D., Ludwig, W., Pot, B., Kersters, K., Schleifer, K.H. and Hammes, W.P.** (1994). Identification of lactobacilli from sourdough and description of *Lactobacillus pontis* sp. nov. International Journal of Systematic Bacteriology, **44**, 223-229.
- Wallbanks, S., Martínez-Murcia, A.J., Fryer, J.L., Phillips, B.A. and Collins, M.D.** (1990). 16S rRNA sequence determination for members of the genus *Carnobacterium* and related lactic acid bacteria and description of *Vagococcus salmoninarum* sp. nov. International Journal of Systematic Bacteriology, **40**, 224-230.
- Weinberg, M., Nativelle, R. and Prévot, A.R.** (1937). Les Microbes Anaérobies. Paris: Masson.
- Weiss, N. and Schillinger, U.** (1984). *Lactobacillus sanfrancisco* sp. nov., nom. rev. Systematic and Applied Microbiology, **5**, 230-232.
- Weiss, N., Schillinger, U. and Kandler, O.** (1984). *Lactobacillus lactis*, *Lactobacillus leichmannii* and *Lactobacillus bulgaricus* subjective synonyms of *Lactobacillus delbrueckii* subsp. *bulgaricus* comb. nov. International Journal of Systematic Bacteriology, **34**, 270-271.
- Weiss, N., Schillinger, U., Laternser, M. and Kandler, O.** (1981). *Lactobacillus sharpeae* sp. nov. and *Lactobacillus agillis* sp. nov., two new species of homofermentative, meso-diaminopimelic acid- containing lactobacilli. Zentralbl. Mikrobiol Parasitenkd Infektionstr. Hyg. Abt. 1 Orig., C2, 242-253. Validation List N^o 8. International Journal of Systematic Bacteriology, 1982, **32**, 266-268.
- Whiley, R.A. and Hardie, J.M.** (1988). *Streptococcus vestibularis* sp. nov. from the human oral cavity. International Journal of Systematic Bacteriology, **38**, 335-339.
- Whiley, R.A., Hall, L.M.C., Hardie, J.M. and Beighton, D.** (1999). A study of small-colony β -haemolytic, Lancefield group C streptococci within the anginosus group: description of *Streptococcus constellatus* subsp. *pharyngis* subsp. nov., associated with the human throat and pharyngitis. International Journal of Systematic Bacteriology, **49**, 1443-1449.
- Whiley, R.A., Fraser, H.Y., Douglas, C.W.I., Hardie, J.M., Williams, A.M. and Collins, M.D.** (1990). *Streptococcus parasanguis* sp. nov., an atypical viridans *Streptococcus* from human clinical specimens. FEMS Microbiology Letters, **68**, 115-122.
- Whiley, R.A., Russell, R.R.B., Hardie, J.M. and Beighton, D.** (1988). *Streptococcus downei* sp. nov. for strains previously described as *Streptococcus mutans* serotype h. International Journal of Systematic Bacteriology, **38**, 25-29.
- White, J.C. and Niven Jr., C.F.** (1946). *Streptococcus* s.b.e.: a streptococcus associated with subacute bacterial endocarditis. Journal of Bacteriology, **51**, 717-722.
- Wiese, B.J., Strohmair, W., Rainey, F.A. and Diekmann, H.** (1996). *Lactobacillus panis* sp. nov., from sourdough with a long fermentation period. International Journal of Systematic Bacteriology, **46**, 449-453.
- Willcox, M.D.P., Zhu, H. and Knox, K.W.** (2001). *Streptococcus australis* sp. nov., a novel oral streptococcus. International Journal of Systematic and Evolutionary Microbiology, **51**, 1277-1281.
- Williams, A.M. and Collins, M.D.** (1990). Molecular taxonomic studies on *Streptococcus uberis* types I and II. Description of *Streptococcus parauberis* sp. nov. Journal of Applied Bacteriology, **68**, 485-490.
- Williams, A.M., Farrow, J.A.E. and Collins, D.** (1989). Reverse transcriptase sequencing of 16S ribosomal RNA from *Streptococcus cecorum*. Letters in Applied Microbiology, **8**, 185-189.
- Williams, A.M., Fryers, J.L. and Collins, M.D.** (1990). *Lactococcus piscium* sp. nov. a new *Lactococcus* species from salmonid fish. FEMS Microbiology Letters, **68**, 109-114.
- Williams, R.E.O., Hirsch, A. and Cowan, S.T.** (1953). *Aerococcus*, a new bacterial genus. Journal of General Microbiology, **8**, 475-480.

Woo, P.C.Y., Tam, D.M.W., Leung, K.W., Lau, S.K.P., Teng, J.L.L., Wong, M.K.M. and Yuen, K.Y. (2002). *Streptococcus sinensis* sp. nov., a novel species isolated from a patient with infective endocarditis. *Journal of Clinical Microbiology*, **40**, 805-810.

Yoon, J.H., Kang, S.S., Mheen, T.T., Ahn, J.S., Lee, H.J., Kim, T.K., Park, C.S., Kho, Y.H., Kang, K.H. and Park, Y.H. (2000). *Lactobacillus kimchii* sp. nov., a new species from *kimchi*. *International Journal of Systematic and Evolutionary Microbiology*, **50**, 1789-1795.

Yumoto, I., Hirota, K., Nodasaka, Y., Yokota, Y., Hoshino, T. and Nakajima, K. (2004). *Alkalibacterium psychrotolerans* sp. nov., a psychrotolerant obligate alkaliphile that reduces an indigo dye. *International Journal of Systematic and Evolutionary Microbiology*, **54**, 2379-2383.

Zanoni, P., Farrow, J.A.E., Phillips, B.A. and Collins, M.D. (1987). *Lactobacillus pentosus* (Fred, Peterson, and Anderson) sp. nov., nom. rev. *International Journal of Systematic Bacteriology*, **37**, 339-341.

Zhang, B., Tong, H. and Dong, X. (2005). *Pediococcus cellicola* sp. nov., a novel lactic acid coccus isolated from a distilled-spirit-fermenting cellar. *International Journal of Systematic and Evolutionary Microbiology*, **55**, 2167-2170.

Zhilina, T.N., Kotsyurbenko, O.R., Osipov, G.A., Kostrinka, N.A. and Zavarzin, G.A. (1997). *Ruminococcus palustris* sp. nov. In Validation of the Publication of New Names and New Combinations Previously Effectively Published Outside the IJSB, List no. 60. *International Journal of Systematic Bacteriology*, **47**, 242.

Zhu, H., Willcox, M.D.P. and Knox, K.W. (2000). A new species of oral *Streptococcus* isolated from Sprague-Dawley rats, *Streptococcus orisratti* sp. nov. *International Journal of Systematic and Evolutionary Microbiology*, **50**, 55-61.